


Ecology: how organisms interact with one another and with their environment.

Environment: abiotic and biotic features **Levels of Organization:** (some species) > Community (diff. species, biotic) > Ecosystem (community+abiotic) > Biosphere (portion of Earth w/living species) **Productivity** **NPP** = rate at which energy is stored as biomass by plants or other primary producers and made available to consumers in ecosystem; **GPP** = rate at which solar energy is captured in sugar molecules; **NPP = GPP - metabolism**; **Net production efficiency = NPP/GPP** **trophic level efficiency:** ratio of production of trophic level to lower level; green plants: 1% of solar energy; herbivores: < 1%; 10% is a lie because it only applies to managed eco efficiency; usually, over 90% of energy transfer between levels is lost as heat. **Microcosm:** small community **Competition:** intraspecific: same species; inter: diff; **Competitive exclusion:** 2 species competing for same limiting resources can't coexist in same place; **Ecological niche:** sum of a species' use of resources; **Resource partitioning:** allows ecologically similar species to coexist in a community if they have significant differences in niches; **Fundamental vs Realized niche:** potentially/actually occupied by a species; **Character displacement:** tendency for characteristics to diverge in more in sympatric (no physical separation) pops than in allopatric pops; **Predation:** (+/-) **Cryptic coloration:** camouflage, **Aposomatic coloration:** warning colors, **Batesian mimicry:** harmless species mimics harmful, **Müllerian mimicry:** multiple unpalatable species resemble each other, **Herbivory:** (+/-) **Organism eats plant or algae, defense:** toxins, spines/thorns; **Parasitism:** (+/-) **Endoparasites:** live in host; **Ectoparasites:** live outside host, -33 species on humans; **Mutualism:** (+/+) **Obligate mutualism:** one species has lost ability to survive without its partner, **Facultative mutualism:** both species can survive alone if necessary; **Commensalism:** (+/0); **Amensalism:** (-/0) **Diversity** **Alpha diversity** - diversity of each site **Beta** - differences in species composition among sites **Gamma** - diversity of entire landscape (regional species pool) **EX:** Yellowstone: Wolf eradication -> inc in elk herds -> dec in willows -> widening of river flow **Species Richness:** count of species **Simpson Index:** $D = \frac{1}{\sum [n(n-1)]} / [N(N-1)]$ probability that two individuals randomly selected from a sample will belong to same species, bigger value of D, LOWER diversity; ranges from 0-1 **Simpson Index of Diversity (1-D):** greater value, greater sample diversity, probability that two individuals randomly selected from a sample will belong to different species; **Reciprocal Index 1/D:** 1 = community containing only one species, higher value, greater diversity, maximum value = number of species in sample **Shannon-Weiner Index:** p_i is often proportion of individuals belonging to i th species in dataset of interest, quantifies uncertainty in predicting species identity of an individual that is taken at random from dataset $H = -\sum [P_i \times \ln(P_i)]$ (for each species) $P_i = \text{sample}/\text{sum}$ | Evenness = H/H_{max} | $H_{\text{max}} = \ln(S)$ where S is species richness(count) **Sorensen's Coefficient (CC)** = $2C/(S_1+S_2)$ Where C is the number of species the two communities have in common, S1 is the total number of species found in community 1, and S2 is the total number of species found in community 2 **Effective population** $N_e = 4 * N_m * N_f / (N_m + N_f)$ **Cohort Life Tables** - L_x is survivorship (prop. that survived to next stage), d_x is mortality (prop. that die during stage), q_x is mortality rate, $L_x m_x$ is avg number of offspring per female or contribution of each age class to overall pop. growth rate, $L_x m_x$, x = weights $L_x m_x$ by life stage, R_{T0} = sum of $L_x m_x$ = reproductive rate, or the average net number of offspring produced by an individual in its lifetime. **Stream Ecology Watersheds** (drainage basin, catchment area): a land area that channels rainfall and snowmelt to creeks, streams, and rivers, and eventually to outflow points such as reservoirs, bays, and the ocean **Riparian Zone** narrow area alongside a stream that has its own special vegetation; contributes nutrients, shade, organic materials for small organisms, soil stability, habitat **Keystone Species:** species whose functions are so intertwined with the lives of other animals that their removal can cause imbalance or collapse **Stream Anatomy** **Edgewater:** habitats may have emergent plants, sheltered overhangs with suspended root mats and leaf packs in quiet back eddies. The composition of macro-invertebrates will tend to differ from that in riffles. Animals survive best in places that provide protection, camouflage and food sources. **Riffles:** Shallow with fast, turbulent water running over rocks. Only animals that cling very well, such as net-winged midges, caddisflies, stoneflies, some mayflies, dace, and sculpins can spend much time here, and plant life is restricted to diatoms and small algae. Riffles are a good place for mayflies, stoneflies, and caddisflies to live because they offer plenty of cobbly gravel to hide in. **Runs:** Close to any pool or riffle is a run, which merely describes a main body of water that runs smoothly downstream. Fishes, like minnows, too small to compete for pools often end up in runs. **Pools:** When a stream meets up with a huge fallen log, or a set of boulders, the water pours over the top. The vertical force of the water falling down on the other side will carve out a pool in the stream. Preferred by trout, mollusks (like clams and snails) and worms. Benefits to slow-moving water is that organic debris settles out into it. Also you don't have to relocate to another area if the stream level starts to lower. **Floods** - Riparian zones depend on floods **Adaptions to flooding:** fishes wait for annual spring flood to start breeding, insect larvae lay eggs, hatch, or metamorphose, new food sources, increased fertility **Dams** changes ecology forever, habitats removed, fish die from turbines **Hydroelectric Systems Impact** Silt Loads; Water Temperatures; River Flow; Dissolved Oxygen. **Channelizing Streams** **Done** to protect property and roads from flooding **Stream** becomes poor in nutrients and habitat; without periodic flooding, riparian zone dies, native fish die; **More** channelization also decreases control over river, and erosion threatens buildings **When** a stream is allowed to meander, it pushes against banks and swirls, reducing energy of water, but when streams are channelized straight down a mountain, it has more energy **Development** **Urban** runoff - more oil, fertilizer, pesticides, herbicides end up in streams **Lack** of trees - takes away shade from stream, warming it up; bugs that feed food chain in stream will no longer fall into stream **Lack** of roots of vegetation will cause soil to erode away into the river (as it is no longer grounded in place with vegetation) **Logging** **Silt** clouds up rivers **Silt** that settles in the bottom of the river prevents eggs of some species (like Salmon and Trout) that reside in gravel in bottom of river from receiving dissolved oxygen from flowing water above **The silt** in between the gravel also destroys the habitat of many aquatic insects, and takes away the food source of fish as well **Mining** **introduces** heavy metal and radioactive waste into river **makes** river acidic **Requires** much water and reduces level of aquifer, drying streams



since there are very little detritus **Can** develop into eutrophic over time **Runoff** brings in mineral nutrients and sediments **Human** activities increase nutrient content of runoff due to fertilizers **Municipal** wastes dumped into lakes enriches N and P, so more phytoplankton **Algal** blooms and increased plant growth creates more detritus and can lead to oxygen depletion. **Meatrophs** increased production, accumulated organic matter, occasional algal bloom, good fishery **Eutrophication** very productive, can experience oxygen depletion, rough fish common **Eutrophication** The process by which a body of water develops a high concentration of nutrients, like nitrates and phosphates. Phosphorus and Nitrogen are both released from sources related to land use. Nutrients cause an increase in blue-green bacteria and algae, which can cover the surface of the lake in mats; this blocks sunlight to plants as algae and bacteria die, they decompose and BOD increases, decreasing DO. If lowered enough, fish will die. **Beign Exorhic basin** - lake where water constantly flows out under all climatic circumstances; usually freshwater since dissolved solids do not accumulate **Endorhic basin** - no water flows out. Limited drainage basin that normally retains water and allows no outflow to other external bodies of water, like rivers and oceans. **Lake Mixing Dimictic** - mixes twice a year, during spring and fall **Monomictic** - 1 mix/yr, mixed all winter and spring **Polyimictic** - many mixes/yr **Meromictic** (never totally mixed be of stagnant bottom layer) - stratified lakes that have two layers that do not mix. **Holomictic** is opposite (turns over completely). **Monolimnion** - Layer of bottom water remains stagnant and anoxic for years **Mixolimnion** - layer that does mix one year in this type of lake; separated from monolimnion by chemocline. Decomposing bacteria makes hypolimnion hypoxic. **Adding** salt to lake increases density, making it harder to overturn. **Aquatic Ecosystems** **Lake Zones** **Lenic Ecosystems** (STILL Water) **Ponds** bottom of the pond still receives light, unlike lakes. **Horizontal Lake Zones** **Littoral Zone:** Near the shoreline; Sunlight penetrates all the way to sediments; Allows for aquatic plants (macrophytes) to grow. **Limnetic Zone** open water, away from shore. **Vertical Lake Zones** **Depth** in which photosynthesis can occur. **Aphotic** Photosynthesis cannot occur; Most organisms are invertebrates. Productivity depends on the organic content of the sediment. **Lake Turnover** heated by the sun The deepest layer, the hypolimnion, is the coldest. The sun's radiation does not reach this cold, dark layer. **During** the fall, the warm surface water begins to cool. As water cools, it becomes more dense, causing it to sink. This dense water forces the water of the hypolimnion to rise, "turning over" the layers. The opposite happens during the spring. **Lenic Ecosystems** (Flowing Water) **Stream** **Order** or waterbody order is a positive whole number used in geomorphology and hydrology to indicate the level of branching in a river system. **Mixing and stratification** **Exhibit** significant vertical stratification with light penetration and temperature **Light** penetration stratification - Ponds or lakes are divided into two layers due to a decrease in light intensity with increasing depth - as light is absorbed by the water and suspended microorganisms. **Photic** zone upper layer where light is sufficient for photosynthesis. **Aphotic** zone bottom layer with little light, no photosynthesis **For** deeper ponds and lakes, temperature stratification occurs; sunlight warms the upper layer as far as it can penetrate **Thermocline** - narrow vertical zone between warm and cold layers where a rapid temp change occurs **Cycles Oxygen** Three main reservoirs: **atmosphere** (air), total content of **biological** matter within biosphere, & **Earth's crust** - Failures in O-cycle w/in hydrosphere (combined mass of water found on, under, & over surface) can result in development of **hypoxic zones**. Main driving factor of O-cycle is photosynthesis, which is responsible for modern Earth's atmosphere & life. **Nitrogen Fixation:** done by free-living (ex. Azotobacter) or symbiotic bacteria known as diazotrophs, which have nitrogenase enzyme that combines gaseous nitrogen with hydrogen to produce ammonia (which is converted into other organic compounds by bacteria). Most biological nitrogen fixation occurs by activity of Mo-nitrogenase (a complex two component enzyme that has multiple metal-containing prosthetic groups), found in variety of bacteria - some Archaea. Symbiotic nitrogen-fixing bacteria (ex. Rhizobium) usually live in root nodules of legumes (peas, alfalfa, & locust trees) & a few non-legumes, forming a mutualistic relationship with plant, producing ammonia in exchange for carbohydrates. Because of this, legumes often increase nitrogen content of nitrogen-poor soils. **Assimilation** - Plants take nitrogen from soil by absorption through their roots via root hairs as amino acids, nitrate ions, or ammonium ions. If nitrate is absorbed, it is first reduced to nitrite ions & then ammonium ions for incorporation into amino acids, nucleic acids, & chlorophyll. Plants that have a symbiotic relationship with rhizobia assimilate some nitrogen in form of (NH₄)⁺ directly from nodules. **Ammonification:** When a plant/animal dies or an animal excretes waste, nitrogen is initially organic. Bacteria or fungi convert organic N w/in remains back into ammonium (NH₄)⁺, a process called ammonification or mineralization **Denitrification:** reduction of nitrates back into nitrogen gas (N₂), completing N cycle, performed by bacterial species such as Pseudomonas & Clostridium in anaerobic conditions. Nitrate used as an electron acceptor in place of oxygen during respiration. These facultatively anaerobic bacteria can also live in aerobic conditions. Denitrification happens in anaerobic conditions ex. waterlogged soils. Denitrifying bacteria use nitrates in soil to carry out respiration & consequently produce N₂, which is inert & unavailable to plants **Nitrification:** Conversion of ammonium to nitrate performed primarily by soil-living & other nitrifying bacteria. In primary stage, oxidation of ammonium (NH₄)⁺ is performed by bacteria such as Nitrosomonas species, which converts ammonium to nitrites (NO₂). Other bacterial species, such as Nitrobacter, are responsible for oxidation of nitrites into nitrates (NO₃). Ammonia conversion to nitrites or nitrates is important because ammonia gas is toxic to plants. **Phosphorus** Essential nutrient for plants & animals essential in form of ions PO₄³⁻ & HPO₄²⁻; part of DNA, ATP, ADP, fats of cell membranes, building block of certain parts of animal & human bodies; doesn't enter atmosphere, remains mostly on land & in rock & soil minerals; liquid at normal temp & pressure; cycling through water, soil, & sediments; slow water cycle; Moves slowly from deposits on land & in sediments, to living organisms, & turn much more slowly back into soil & water sediment. Passes through plants & animals much faster than through rocks & sediments. Animals get it by eating plants or plant-eating animals; When plants & animals die, phosphates return soils or oceans, ending up in sediments or rock formations again, remaining there for millions of years. Eventually, it is released through weathering & it begins again; Most commonly found in rock formations & ocean sediments as phosphate salts. Phosphate salts released through rocks usually dissolve in soil water & will be absorbed by plants. Quantities of phosphorus in soil are small, often making them limiting factors for plant growth. Not very water-soluble, making them limiting factors for plant growth in marine ecosystems; Constant additions of phosphates by humans & exceeding natural concentrations disrupt P-cycle strongly. Phosphate can build up in rivers & lakes, causing excessive algae growth; Increasing phosphorus concentrations in surface waters raise growth of phosphate-dependent organisms, like algae & duckweed. These organisms use great amounts of oxygen & prevent sunlight from entering water, known as eutrophication. **Wastewater treatment** **Usable Water Treatment** 1 **Coagulation and Flocculation:** First step. Chemicals with a positive charge are added to the water to neutralize the negative charge of dirt and other dissolved particles. Then the particles bind with the chemicals and form larger particles, called **floc**. 2 **Sedimentation:** floc settles to the bottom of the water supply, due to its weight. 3 **Filtration:** once the floc has settled to the bottom of the water supply, the clear water on top will pass through filters of varying compositions (sand, gravel, and charcoal), in order to remove dissolved particles (dust, parasites, bacteria, viruses, chemicals). 4 **Disinfection:** a disinfectant (chlorine, chloramine) may be added in order to kill any remaining parasites, bacteria, and viruses, and to protect the water from germs when it is piped to homes and businesses. **Groundwater treatment technique:** **Air** sparging: inject oxygen into groundwater; When used in combination with soil vapor extraction (SVE), air bubbles carry vapor phase contaminants to a SVE system which removes them. **Bioreactors:** vessel in which a chemical process is carried out which involves organisms or biochemically active substances derived from such organisms Chemical Oxidation **Constructed wetland:** artificial wetland to treat municipal or industrial wastewater, greywater or stormwater runoff. It may also be designed for land reclamation after mining, or as a mitigation step for natural areas lost to land development. use natural functions vegetation, soil, and organisms to treat wastewater. Acts as a biofilter and removes pollutants. **Dual phase extraction:** uses a high-vacuum system to remove both contaminated groundwater and soil vapor. **Pump and treat:** treatment of pumped groundwater before it is released. **Phytoremediation:** direct use of living green plants for in situ, or in place, removal, degradation, or containment of contaminants

in soils, sludges, sediments, surface water and groundwater. **Septic Tank Disposal System:** A sewer line from the house leads to an underground septic tank designed to separate solids from liquid, digest and store organic matter, and allow the treated sewage to seep into the surrounding soil. As the wastewater moves through the soil, it is further treated by the natural processes of oxidation and filtering. **This** method can fail if the tank isn't pumped out when it's full of solids or if there is poor drainage in the surrounding soil. **Primary Treatment** removes 30-40% of BOD by volume, mainly in the form of suspended solids and organic matter. **Incoming** raw sewage passes in and is first passed through a series of screens to remove large floating organic material. **Sewage** next enters the grit chamber, where sand, small stones, and grit are removed **Then** primary sedimentation tank, where particulate matter settles out to form a sludge. Chemicals can be used to help the settling process **Sludge** is removed and transported for further processing **Secondary Treatment** **Activated sludge** (most common treatment) **Aeration tank:** wastewater is pumped with air and some sludge from final sedimentation tank; sludge contains aerobic bacteria that consume organic material; the wastewater enters the final sedimentation tank and sludge settles out; some of activated sludge is recycled with new air and wastewater; the rest of the sludge goes to digester, where **anaerobic bacteria** further break down sludge; **C H₄** is a product of this anaerobic breakdown, and is captured for fuel. **Wastewater** is finally disinfectant to remove pathogens, usually by chlorination **Advanced Wastewater Treatment:** There may still be nutrients, organic chemicals, and heavy metals that can be removed with sand filters, carbon filters, and special chemicals **Septic Tank:** The sewer line from the house leads to an underground septic tank, which is designed to separate solids from liquid, treat, and store organic matter through a period of detention, and allow the clarified liquid to discharge into the drain (absorption) field from piping through which the treated sewage seeps into the surrounding soil. As the wastewater moves through the soil, it is further treated by natural processes of oxidation and filtering. These may fail bc of failure to pump out the septic tank when it is full of solids, and poor soil drainage, which allows the effluent to rise to the surface in wet weather **Analysis:** Salinity, pH, alkalinity, phosphates, nitrates, turbidity, dissolved oxygen (DO), temperature, fecal coliform, total solids, and biological oxygen demand (BOD). **Salinity** Water can be classified by its salinity as such: fresh water has a ppt of < 0.5 which means that there are 0.5 molecules of dissolved salt for every 100 molecules of solution, or 1 molecule of salt per every 200 molecules of solution. Brackish water has a ppt between 0.5 and 3, saline water has a ppt between 3 and 30, and brine has a ppt of >30. The only water safe for human consumption is fresh water, and drinking water often achieves salinity levels as low as 0.1 ppt. In ocean water, total salt content makes up 3.5% (35 ppt) of ocean water, the other 96.5% being water. Of the dissolved salts in ocean water, 86% of the salts is sodium chloride (5% of all the water, 30 ppt). The other 15% of the total salts are other salt ions such as Magnesium, Strontium etc. (5% of total ocean water, 5 ppt). **Consequences** Salinity in rivers and lakes of the US has recently been increasing due to road salt and other salt de-icers in runoff. **It** is very expensive to remove salt from water, and thus it is expensive to create drinking water **pH** $pH = -\log[H^+]$ **The** normal pH of rivers in the United States is 6.5 to 8.5, and values between 6.0 and 9.0 can support life for fish and invertebrates. **Influenced** by **Human** processes like automobile/fossil fuel power plant emissions release nitrogen oxides and sulfur dioxide. **Cool** mine drainage can lead to iron sulfide mixing with water. **Natural:** Limestone is a base when dissolved in water, so it can neutralize the effects of acids and increase the pH of the water. **Alkalinity** The ability of a solution to neutralize an acid without changing the overall pH. From presence of buffers. **Effects** If alkalinity is too low, the ecosystem has low stability as it is susceptible to sudden pH changes from devices such as acid rain or other pollution, which can be harmful to the flora and fauna of the ecosystem. If alkalinity is too high, the buffer acids and bases in the buffer solution can render the ecosystem uninhabitable. **Phosphates** Phosphates can become a limiting nutrient in many systems, usually freshwater systems. Generally, the only negative effect of an overabundance of phosphates is **eutrophication**. Phosphates can contribute to total dissolved solids. Extremely high levels of phosphates in drinking water can cause digestive issues. **Nitrates** Nitrates can be a limiting nutrient, usually in marine systems rather than freshwater systems. However, high levels of nitrates in aquatic ecosystems can be detrimental to ecosystem health, inhibit the growth of some organisms, cause stress, and contribute to eutrophication. Nitrates do contribute to total dissolved solids and can be used as a water quality indicator. **Turbidity** **Increased** salinity leads to low turbidity because salt ions bunch suspended solids in the water together, making it more easy for suspended solids to settle at the bottom. So oceans and estuaries tend to be more clear than freshwater. **Turbidity** is a measurement of how much light the water scatters, so it gets affected by not only suspended solids, but also colored dissolved materials, or dyes in the water. **"Turbidity maximum zone":** caused by change in flow when freshwater stream enters saltwater estuary. **Clear** = under 20 mg/L (of total suspended solids) **Cloudy** > over 40 mg/L **temperature:** $\downarrow DO$; \uparrow temperature, stratification \rightarrow may create hypoxic zones in lower levels because this is where most decomposition occurs; harmful to living organisms \uparrow ; turbidity, \downarrow photosynthesis and plant growth, \downarrow DO (b/c less plants and decomposition causes loss of DO), \downarrow primary food source in food chain, \downarrow pop. Of org. **Dissolved Oxygen** **Surface** waters contain between 5 and 15 ppm of dissolved oxygen. If a stream or river has below 5 ppm of dissolved oxygen, then that can put aquatic life under stress, and below 1-2 ppm for a few hours can kill large fish living in the river. **Supersaturation** of oxygen (levels of DO over 100%) can occur naturally through photosynthetically active species. Supersaturation can also occur through rapid changes in the environment that occur too quickly for the system to reach equilibrium, giving rise to DO levels over 100% temporarily. **< 6 mg/L** harmful to pond life **As** temperature increases, DO decreases **Oxygen** level changes result from streamflow, temperature, and run-off **At** sea level, **typical DO concentrations** in 100-percent saturated fresh water will range from 7.56 mg/L (or 7.56 parts oxygen in 1,000,000 parts water) at 30°C to 14.62 mg/L at zero 0°C. **Measured** by the **Winkler Test** **Biochemical Oxygen Demand** **(BOD)** measures how fast organisms use up the oxygen in the water. Aerobic microbes use oxygen to oxidize the organic matter in the water, using the energy that is released in the process for growth and reproduction and creating a demand for DO. **Most** pristine rivers should have a 5-day BOD below 1 mg/L. Moderately polluted rivers may have a BOD between 2 and 8 mg/L. **Municipal** sewage that is treated with a three-stage process would have a BOD of about 20 mg/L. Untreated sewage has varying BOD but averages about 200 mg/L **Temperature** **Affects** the amount of gases such as oxygen that can be dissolved in the water - cold water holds more oxygen than warm water \rightarrow Increases the metabolic rates of aquatic organisms. **Affects** the rate of photosynthesis by aquatic plants and algae. **Increases** the sensitivity of organisms to disease, parasites and pollution. **Small** chronic temperature changes can adversely affect the reproductive systems of aquatic organisms. **Raising** water temperature increases decomposition rate of organic matter, depleting DO. **Types** of temperature changes include natural seasonal changes, man's activities, industrial thermal pollution as discharge of cooling water, stormwater runoff from heated surfaces as streets, roofs, parking lots, soil erosion increasing water turbidity which warms the water, removal of shade trees from along the shores. **Tidal Solids** Dissolved solids in the body of water and is subdivided into two categories: total suspended solids (TSS) and total dissolved solids (TDS). Tested by conductivity and dehydration, measures the suspended and dissolved solids in water; suspended solids retain in water filters and sink to bottom of column (clay, silt, plankton, organic waste, and inorganic precipitates); dissolved solids pass through water filters (calcium, bicarbonate, nitrogen, phosphorus, iron, and sulfur and other ions in the water); concentration of dissolved solids determines flow of water in and out of cells of aquatic organisms; many dissolved solids are essential nutrients of life; low concentration of dissolved solids limit aquatic growth; elevated levels lead to accelerated eutrophication, increased turbidity, and decreased water quality; sources of elevated levels of total solids include runoff from agricultural activities, dredging, mining, salt from winter streets, fertilizers from lawns, water treatment plants, plant materials, soil particles and erosion, and decaying organic matter; high concentrations of suspended solids lead to increased turbidity, reduced photosynthesis, and increased water temperature due to increased absorption of light and binding of toxic chemicals and/or heavy metals, **Invasive Species**

Triploid grass carp may be the most cost-effective method that currently exists for Brazilian waterweed eradication, as well as other aquatic plants. **Black Carp** - Young eat mostly zooplankton, later insect larvae + organic debris. Adults eat mollusks. Introduced to control snails in aquaculture ponds. **Silver "Flying" carp** - eat phytoplankton. Introduced to control plankton in aquaculture. **Grass Carp** - Aquatic plants, but also can eat small organic debris, insects, small fish, earthworms, other vertebrates if plant material is not available. Introduced to control plants. Carp need constant feeding bc no true stomach; **bioinvasions:** Organisms that are used to monitor the health of an ecosystem. Some are tolerant to pollution and some are not. Finding few or only those that are tolerant indicates the water is polluted or something else has changed about the stream. **Wetlands** region of land that holds a great deal of water for significant periods of time, and that contains specialized plants able to grow in these wet conditions. Among richest and most valuable of biomes. Diverse invertebrate community, which supports a wide variety of birds. Variety of herbivorous species consume algae, detritus, plants. Provide water storage basins that reduce intensity of flooding. Improve water quality by filtering pollutants. **Basin wetlands** - develop in shallow basins ranging from upland depressions to lakes and ponds that have filled in **Riverine wetlands** - develop along shallow and periodically flooded banks of streams and rivers **Fringe wetlands** - along coasts of large lakes and seas where rising lake levels or tides cause water to flow back and forth **River Continuum** Upper reach - narrow, shallow, fast, CPOM input from riparian, sunlight influence minimal, shredders+collectors dominate Middle reach - open canopy so sunlight, algae+vascular plants, FPOM drift from upper reach, collectors+grazers dominate Lower reach - wider, deeper, slower, less input from riparian, sunlight matters but depth of bed is big; slow water lets FPOM settle, collectors+predators dominate CPOM - coarse particulate organic matter, FPOM - fine POM **Embeddedness:** Degree to which fine sediments surround a coarse substrate in a stream bed. More embedded means interstitial space is filled, means less room for critters. **Factors determining river bank:** Runoff from city streets, lack of riparian zone/vegetation, different soil types. **From Lic:** Aquatic Ecology: there are abiotic factors(water, salinity, light, pressure, temperature, dissolved gases, pH, tides, currents, waves, stratum, nutrient supply, exposure to air) and biotic factors. Zonation- major division of marine ecosystems. The pelagic zone includes the neritic zone(productive coastal waters) and oceanic zone(deep waters of open ocean). Benthic zone extends from seashore to deeper parts of sea. Estuaries-salt-wedge(mouth of river, wedge-shaped lower salt section), fjord(u-shaped salty basin), slightly stratified(partially mixed), vertically stratified(well mixed). Zoning include wetland pools, salt flat, salt marsh, mangrove zone, intertidal mudflat, and subtidal zone. Oyster reefs, kelp forests, rocky/soft shore, submerged vegetation, mangrove, swamp, mud/sand flat, tidal stream, barrier beach, salt marsh.

