

Midterm

Question 1 (A)

Given:

$(P \vee Q) \rightarrow R$ and $(P \rightarrow R) \wedge (Q \rightarrow R)$

Claim: They are logically equivalent.

Proof:

(\Rightarrow direction):

Assume $(P \vee Q) \rightarrow R$.

If P is true, then $P \vee Q$ is true $\rightarrow R$ is true $\rightarrow P \rightarrow R$ holds.

If Q is true, then $P \vee Q$ is true $\rightarrow R$ is true $\rightarrow Q \rightarrow R$ holds.

So, $(P \rightarrow R) \wedge (Q \rightarrow R)$.

(\Leftarrow direction):

Assume $(P \rightarrow R) \wedge (Q \rightarrow R)$.

If $P \vee Q$ is true, then either P or Q is true.

If P , then R (since $P \rightarrow R$)

if Q , then R

So,

$(P \vee Q) \rightarrow R$ holds.

Question 1 (B)

A, C, and D are tautologies.

Question 1 (C)

$P \rightarrow (Q \rightarrow R) \equiv P \rightarrow (\neg Q \vee R) \equiv \neg P \vee \neg Q \vee R \equiv \neg(P \wedge Q) \vee R \equiv (P \wedge Q) \rightarrow R$

So both expressions are logically equivalent.

Question 1 (D)

Let:

- $C(x)$: "x owns a cat"
- $L(x)$: "x loves animals"

Then:

- Premise 1: $\forall x (C(x) \rightarrow L(x))$
- Premise 2: $\exists x C(x)$
- Conclusion: $\exists x L(x)$

Valid by universal instantiation and modus ponens .

Question 2 (A)

If $\frac{1}{f(x)}$ is not continuous, then $f(x) = 0$ at some point

Because if $f(x) \neq 0$ everywhere and is continuous, then $\frac{1}{f(x)}$ is continuous.

It's Proved via contrapositive.

Question 2 (B)

Proof of contradiction:

Assume $\sqrt{2} * \sqrt{3} = r$, where r is rational and $\sqrt{2}$ is irrational

Then:

$$\sqrt{3} = \frac{r}{\sqrt{2}}$$

Since r is rational and $\sqrt{2}$ is irrational, $\frac{r}{\sqrt{2}}$ is irrational,

-> contradiction: $\frac{r}{\sqrt{3}}$ can't be both rational and irrational

So, the product must be irrational

Question 2 (C)

Using inclusion-exclusion:

Total solutions without constraints:

$$\frac{(12+3-1)}{2} = \frac{(14)}{2} = 91$$

Subtract:

- $x > 5 \rightarrow \frac{8}{2} = 28$
- $y > 6 \rightarrow \frac{7}{2} = 21$
- $z > 7 \rightarrow \frac{6}{2} = 15$

$$91 - (28 + 21 + 15) = 27$$

Question 2 (D)

Total circular arrangements of 6 people: $(6 - 1)! = 120$

bad cases (two specific people together)

$$(5 - 1)! \cdot 2 = 48$$

$$120 - 48 = 72$$

Question 3 (A)

365 possible birthdays

Need at least 366 people to ensure a shared birthday

The answer is 366

Question 3 (B)

- Total 8-letter passwords with no repeated letters

$$\text{Total (no restriction)} = P(26, 8) = 26 \times 25 \times 24 \times 23 \times 22 \times 21 \times 20 \times 19$$

- We must choose and arrange 8 distinct consonants from the 21 available.

$$\text{All consonants (no vowels)} = P(21, 8) = 21 \times 20 \times 19 \times 18 \times 17 \times 16 \times 15 \times 14$$

- Subtract to get number of valid passwords (at least one vowel)

$$\text{Valid passwords} = P(26, 8) - P(21, 8)$$

$$26 \times 25 \times 24 \times 23 \times 22 \times 21 \times 20 \times 19 - 21 \times 20 \times 19 \times 18 \times 17 \times 16 \times 15 \times 14$$

Which is

$$54,786,211,200$$

Question 3 (C)

$$\text{Given: } x_1 + x_2 + x_3 + x_4 = 20$$

Each $x_i > 1$ and $x_i \neq 2, 3$

Let $x_i \geq 4$: so define $y_i = x_i - 4 \rightarrow y_i \geq 0$

$$y_1 + y_2 + y_3 + y_4 \Rightarrow \frac{4+4-1}{3} = \frac{7}{3} = 35$$

Question 4 (A) i

DFA for accepting binary strings containing "101" :

- States: q_0 (start), q_1 , q_2 , q_3 (accepting)
- Alphabet: $\{0, 1\}$
- Transitions:
 - $q_0 \xrightarrow{1} q_1$, $q_0 \xrightarrow{0} q_0$
 - $q_1 \xrightarrow{0} q_2$, $q_1 \xrightarrow{1} q_1$
 - $q_2 \xrightarrow{1} q_3$, $q_2 \xrightarrow{0} q_0$

- $q_3 \xrightarrow{0/1} q_3$ (stay in accepting)
- Accepting State: q_3
- Accepts strings containing "101" as a substring.

Question 4 (A) ii

String: "1101"

- Loops on q_1 with extra 1s
- Exits to accept via "01"

Question 4 (A) iii

Modify DFA to accept even number of 1s.

Track parity: even \leftrightarrow odd \leftrightarrow even.

New string accepted: "1100"

Not accepted under old DFA (no "101").

Question 4 (B)

Regex for: no "11", ends in "0":

$(0+10)^* 0$

All binary strings that do not contain "11" and end with

Question 5 (A)

- abc123
- A1b2C3
- z9

Question 5 (B)

Regex: $(0 + 1)^* 101 (0 + 1)^*$

Then filter out strings containing "111"

Question 5 (C)

Use Pumping Lemma to show:

$L = \{a^p \mid p \text{ is prime}\}$ is not regular.

Assume regular \rightarrow pumping length p

Pick $\{a^p \mid p \text{ is prime}\}$ is not regular

Assume regular \rightarrow pumping length p

Pick a^q , where $q > p, q$ prime

Split $a^q = xyz$, pump y : get length non-prime \rightarrow contradiction.

Not regular

Question 5 (D)

Let:

- $G^1 : S \rightarrow aSb \mid \epsilon$
- $G^2 : S \rightarrow aS \mid b$

String "aabb" $\in G_1$ (balanced recursion)

Not $\in G_2$ (unbalanced structure)

"aabb" $\in L(G_1)$, but not $\in L(G_2)$