



Hierarchical  
Models  
1/ 10

D.E. Brown

Hierarchical  
Modeling  
Example

# Hierarchical Models

Donald E. Brown

School of Data Science  
University of Virginia  
Charlottesville, VA 22904



# Agenda

Hierarchical  
Models  
2/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- 1 Hierarchical Modeling
- 2 Radon Testing Example



# Bayesian Multilevel Models

Hierarchical  
Models  
3/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- Multilevel models are generalizations of regression models where the parameters vary by group



# Bayesian Multilevel Models

Hierarchical  
Models  
3/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- Multilevel models are generalizations of regression models where the parameters vary by group
- In Bayesian multilevel models, the parameters are random with distributions



# Bayesian Multilevel Models

Hierarchical  
Models  
3/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- Multilevel models are generalizations of regression models where the parameters vary by group
- In Bayesian multilevel models, the parameters are random with distributions
- Hierarchical models are special cases of multilevel models where the groups are nested



# Bayesian Multilevel Models

Hierarchical  
Models  
3/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- Multilevel models are generalizations of regression models where the parameters vary by group
- In Bayesian multilevel models, the parameters are random with distributions
- Hierarchical models are special cases of multilevel models where the groups are nested
- Multilevel models allow us to answer questions about grouped data where there are varying sizes to the groups, e.g.,



# Bayesian Multilevel Models

Hierarchical  
Models  
3/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- Multilevel models are generalizations of regression models where the parameters vary by group
- In Bayesian multilevel models, the parameters are random with distributions
- Hierarchical models are special cases of multilevel models where the groups are nested
- Multilevel models allow us to answer questions about grouped data where there are varying sizes to the groups, e.g.,
  - Schools and student or teacher scores



# Bayesian Multilevel Models

Hierarchical  
Models  
3/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- Multilevel models are generalizations of regression models where the parameters vary by group
- In Bayesian multilevel models, the parameters are random with distributions
- Hierarchical models are special cases of multilevel models where the groups are nested
- Multilevel models allow us to answer questions about grouped data where there are varying sizes to the groups, e.g.,
  - Schools, and student or teacher scores
  - Hospitals, and patient or caregiver results





# Bayesian Multilevel Models

Hierarchical  
Models  
3/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- Multilevel models are generalizations of regression models where the parameters vary by group
- In Bayesian multilevel models, the parameters are random with distributions
- Hierarchical models are special cases of multilevel models where the groups are nested
- Multilevel models allow us to answer questions about grouped data where there are varying sizes to the groups, e.g.,
  - Schools, and student or teacher scores
  - Hospitals, and patient or caregiver results
  - Political and geographic units, and economic or social outcomes



# Problem Structure for Hierarchical Models

Hierarchical  
Models  
4/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- Suppose we have  $N$  observations of a target variable  $y_i, i = 1, \dots, N$  with predictors in an  $N \times k$  matrix  $\mathbf{X}$ .



# Problem Structure for Hierarchical Models

Hierarchical  
Models  
4/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- Suppose we have  $N$  observations of a target variable  $y_i, i = 1, \dots, N$  with predictors in an  $N \times k$  matrix  $\mathbf{X}$ .
- Let  $X_i$  be the  $i^{\text{th}}$  row of  $\mathbf{X}$ .



# Problem Structure for Hierarchical Models

Hierarchical  
Models  
4/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- Suppose we have  $N$  observations of a target variable  $y_i, i = 1, \dots, N$  with predictors in an  $N \times k$  matrix  $\mathbf{X}$ .
- Let  $X_i$  be the  $i^{\text{th}}$  row of  $\mathbf{X}$ .
- Suppose we have  $J$  groups indexed by  $j = 1, \dots, J$



# Problem Structure for Hierarchical Models

Hierarchical  
Models  
4/10

D.E. Brown

Hierarchical  
Modeling

Example

- Suppose we have  $N$  observations of a target variable  $y_i, i = 1, \dots, N$  with predictors in an  $N \times k$  matrix  $\mathbf{X}$ .
- Let  $X_i$  be the  $i^{\text{th}}$  row of  $\mathbf{X}$ .
- Suppose we have  $J$  groups indexed by  $j = 1, \dots, J$
- Traditional regression model with dummy variables,  $z_1, \dots, z_{J-1}$

$$y_i = \alpha + \boldsymbol{\theta} \mathbf{X}_i + z_1 + \dots + z_{J-1} + \epsilon_i$$



# Pooling Models

Hierarchical  
Models  
5/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- A complete pooling model - ignore the groups

$$y_i = \alpha + \boldsymbol{\theta} \mathbf{X}_i + \epsilon_i$$



# Pooling Models

Hierarchical  
Models  
5/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- A complete pooling model - ignore the groups

$$y_i = \alpha + \boldsymbol{\theta} \mathbf{X}_i + \epsilon_i$$

- A no pooling model - one equation for each group,  
 $j = 1, \dots, J$

$$y_{j[i]} = \alpha_{j[i]} + \boldsymbol{\theta}_{j[i]} \mathbf{X}_{j[i]} + \epsilon_{j[i]}$$



# Partial Pooling

Hierarchical  
Models  
6/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- Partial pooling - no predictors

$$\hat{\alpha}_j \approx \frac{\frac{N_j}{\sigma_y^2} \bar{y}_j + \frac{1}{\sigma_\alpha^2} \bar{y}_{\text{all}}}{\frac{n_j}{\sigma_y^2} + \frac{1}{\sigma_\alpha^2}}$$

where  $\sigma_y^2$  is the within -group variance computed using either an average or a value for each group (i.e.,  $\sigma_{y,j}^2$ ) and  $\sigma_\alpha^2$  is the variance of the average values,  $\bar{y}_j$ , in the groups.  $\bar{y}_{\text{all}}$  is the overall average.





# Partial Pooling

Hierarchical  
Models  
6/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- Partial pooling - no predictors

$$\hat{\alpha}_j \approx \frac{\frac{N_j}{\sigma_y^2} \bar{y}_j + \frac{1}{\sigma_\alpha^2} \bar{y}_{\text{all}}}{\frac{n_j}{\sigma_y^2} + \frac{1}{\sigma_\alpha^2}}$$

where  $\sigma_y^2$  is the within- group variance computed using either an average or a value for each group (i.e.,  $\sigma_{y,j}^2$ ) and  $\sigma_\alpha^2$  is the variance of the average values,  $\bar{y}_j$ , in the groups.  $\bar{y}_{\text{all}}$  is the overall average.

- Averages from groups with less observations carry less weight and will “shrink” toward the overall average



# Partial Pooling

Hierarchical  
Models  
6/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- Partial pooling - no predictors

$$\hat{\alpha}_j \approx \frac{\frac{N_j}{\sigma_y^2} \bar{y}_j + \frac{1}{\sigma_\alpha^2} \bar{y}_{\text{all}}}{\frac{n_j}{\sigma_y^2} + \frac{1}{\sigma_\alpha^2}}$$

where  $\sigma_y^2$  is the within-group variance computed using either an average or a value for each group (i.e.,  $\sigma_{y,j}^2$ ) and  $\sigma_\alpha^2$  is the variance of the average values,  $\bar{y}_j$ , in the groups.  $\bar{y}_{\text{all}}$  is the overall average.

- Averages from groups with less observations carry less weight and will “shrink” toward the overall average
- Averages from bigger groups carry more weight and are closer to their unpooled estimates



# Basic Hierarchical Models

Hierarchical  
Models  
7/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- A basic hierarchical model with varying intercepts and varying slopes is given by

$$y_i = \alpha_{j[i]} + \boldsymbol{\theta}_{j[i]} \mathbf{X}_i + \epsilon_i$$

where  $\alpha_{j[i]}$ ,  $\boldsymbol{\theta}_{j[i]}$  are the parameters for the group, say  $j$ , associated with observation  $i$ .



# Summary

Hierarchical  
Models  
8/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- Complete pooling cannot provide estimates for groups in data



# Summary

Hierarchical  
Models  
8/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- Complete pooling cannot provide estimates for groups in data
- No pooling gives poor and possible extreme estimates for groups with a small size



# Summary

Hierarchical  
Models  
8/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- Complete pooling cannot provide estimates for groups in data
- No pooling gives poor and possible extreme estimates for groups with a small size
- Partial pooling captures differences in groups and better estimates smaller-sized groups but does not use predictor variables



# Summary

Hierarchical  
Models  
8/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- Complete pooling cannot provide estimates for groups in data
- No pooling gives poor and possible extreme estimates for groups with a small size
- Partial pooling captures differences in groups and better estimates smaller-sized groups but does not use predictor variables
- Hierarchical modeling uses the available information in the data, i.e., predictor variables, to better predict group target or response values, even in small groups



# Agenda

Hierarchical  
Models  
9/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- 1 Hierarchical Modeling
- 2 Radon Testing Example





# Example - Radon Testing (Gellman & Hill, 2006)

Hierarchical  
Models  
10/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- Radon is a radioactive gas that enters homes through contact points with the ground.



# Example - Radon Testing (Gellman & Hill, 2006)

Hierarchical  
Models  
10/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- Radon is a radioactive gas that enters homes through contact points with the ground.
- It is a carcinogen that is the primary cause of lung cancer in non-smokers.



# Example - Radon Testing (Gellman & Hill, 2006)

Hierarchical  
Models  
10/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- Radon is a radioactive gas that enters homes through contact points with the ground.
- It is a carcinogen that is the primary cause of lung cancer in non-smokers.
- Varying levels between houses.



# Example - Radon Testing (Gellman & Hill, 2006)

Hierarchical  
Models  
10/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- Radon is a radioactive gas that enters homes through contact points with the ground.
- It is a carcinogen that is the primary cause of lung cancer in non-smokers.
- Varying levels between houses.
- EPA measured levels in 80k houses - example data just from Minnesota



# Example - Radon Testing (Gellman & Hill, 2006)

Hierarchical  
Models  
10/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- Radon is a radioactive gas that enters homes through contact points with the ground.
- It is a carcinogen that is the primary cause of lung cancer in non-smokers.
- Varying levels between houses.
- EPA measured levels in 80k houses - example data just from Minnesota
- Two important predictors:



# Example - Radon Testing (Gellman & Hill, 2006)

Hierarchical  
Models  
10/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- Radon is a radioactive gas that enters homes through contact points with the ground.
- It is a carcinogen that is the primary cause of lung cancer in non-smokers.
- Varying levels between houses.
- EPA measured levels in 80k houses - example data just from Minnesota
- Two important predictors:
  - Floor - basement or first



# Example - Radon Testing (Gellman & Hill, 2006)

Hierarchical  
Models  
10/ 10

D.E. Brown

Hierarchical  
Modeling

Example

- Radon is a radioactive gas that enters homes through contact points with the ground.
- It is a carcinogen that is the primary cause of lung cancer in non-smokers.
- Varying levels in between houses.
- EPA measured levels in 80k houses - example data just from Minnesota
- Two important predictors:
  - Floor - basement or first
  - County uranium levels