



History and Physical:

Historical Findings	Physical Findings
<ul style="list-style-type: none">Known historyNew onset	<ul style="list-style-type: none">BGL < 70 mg/dLAMS, tremors, weakness, nausea/vomiting, intense hunger, diaphoresisBGL < 40 mg/dL should be treated, regardless of presence of other signs/symptoms

Assessment:

- Medical Assessment
- Differential Diagnoses: Infection, Infarction, Medication-related, Toxins, Drugs/Alcohol

Clinical Management Options:

Interventions	Pharmacology
<ul style="list-style-type: none">Oxygen therapy as appropriateVascular Access	<ul style="list-style-type: none">Oral glucose 15-31 G PO<ul style="list-style-type: none">Only patients with intact gag reflex and able to follow commands to swallow medicationThiamine 100mg IV/IO/IM<ul style="list-style-type: none">Prior to D50% administrationOnly with evidence of malnutritionDextrose 25-50 G IV/IO<ul style="list-style-type: none">May be repeated x1The following doses equate to 25 G<ul style="list-style-type: none">50ml Dextrose 50%250ml Dextrose 10%500ml Dextrose 5%Glucagon 1 mg IM/SQ if unable to obtain vascular access<ul style="list-style-type: none">Should be followed by Oral Glucose once patient can follow commands to swallow