

Rigid Analytic Geometry Week 2 Solution

So Murata

SoSe 26, University of Bonn

Problem 1

Let $e_1, \dots, e_m$ be the elements of the standard basis. Then

$$\|e_i|K^m\| = 1, x = \sum_{i=1}^m x_i e_i, x_i \in K.$$

Let $C \in K$ be such that

$$C = \max\{\|e_i\|_1 \mid i = 1, \dots, m\}.$$

Then,

$$\|x\|_1 \leq C \max\{|x_i|_K \mid i = 1, \dots, m\} = C\|x|K^m\|.$$

Problem 2

Suppose for all $C > 0$ there is $x \in K^m$ such that

$$\|x|K^m\| > n\|x\|_1.$$

Take a sequence $(x_n)_{n \in \mathbb{N}}$ in K^m such that

$$\|x_n|K^m\| > n\|x_n\|_1.$$

Then set $(y_n)_{n \in \mathbb{N}}$ be such that

$$\forall n \in \mathbb{N}, y_n = \frac{x_n}{x_n^{n_i}},$$

where $x_n^{n_i}$ is the coefficient of x_n in the standard basis with the largest absolute value. Since $(y_n)_{n \in \mathbb{N}}$ is a bounded sequence and K^m is Banach, there is a convergent subsequence of it with its limit y . Then, taking the limit in

$$1 > \frac{\|x\|_1}{\|x|K^m\|} n = \|y_n\|_1 n.$$

we see that

$$1 > \|y\|_1 \lim_{n \rightarrow \infty} n.$$

But $\|y\|_1 \neq 0$ since

$$1 = \lim_{n \rightarrow \infty} \|y_n|K^m\| = \|y|K^m\|.$$

Thus $\|y\|_1 \neq 0$ as $y \neq 0$. This is a contradiction.

Problem 3

Consider p and q -adic norm and $\mathbb{Q}$ where p, q are distinct primes. Then for any $n \in \mathbb{N}$, consider

$$\left| \left(\frac{1}{p} \right)^n \right|_p = p^n, \left| \left(\frac{1}{p} \right)^n \right|_q = 1.$$

Thus $|\cdot|_p, |\cdot|_q$ are not equivalent.

Problem 4

For $x \in L$, consider,

$$m_x(y) = xy.$$

Then m_x is a linear operator between Banach algebras thus bounded. Also

$$T : L \ni x \mapsto m_x \in \text{Hom}(L, L)$$

is also a linear operator in finite dimensional Banach spaces thus bounded. We conclude,

$$\|m_x\|_{\text{op}} \leq C\|x\|.$$

Thus,

$$\|xy\| \leq \|m_x\|\|y\| \leq C\|x\|\|y\|.$$

Problem 5

Let $x \in \mathfrak{g}_{\mathfrak{U}}$, then

$$\{\mathfrak{p} \in \text{Spec } A \mid x \in \mathfrak{p}\} \in \mathfrak{U}.$$

Thus,

$$\{\mathfrak{p} \in \text{Spec } A \mid x \in \mathfrak{p}\} \subseteq \{\mathfrak{p} \in \text{Spec } A \mid ax \in \mathfrak{p}\}.$$

As $\mathfrak{U}$ is a filter, $\{\mathfrak{p} \in \text{Spec } A \mid ax \in \mathfrak{p}\} \in \mathfrak{U}$. Thus $ax \in \mathfrak{g}_{\mathfrak{U}}$.

Problem 6

Let $X = \text{Sper } A$. Then X is T_0 since for any distance $x, y \in X$, there exists an element a such that $a \in x$ but $b \notin y$. Thus $\mathcal{P}(-a)$ will separate them.

Let Z be an irreducible closed subset of X . Consider

$$C_Z = \bigcup_{C \in Z} C.$$

Then C_Z is clearly a proper cone. To see this is prime, note that if there is $a \in A$ such that $a \in C, -a \notin C, a \notin C', -a \in C'$ for some $C, C' \in Z$. Then we have two non-empty open subsets $\mathcal{P}(a), \mathcal{P}(-a)$ of Z whose intersection is empty. This contradicts to the fact that Z is irreducible. Thus C_Z is prime. Since $C_Z \subseteq C$ for any $C \in Z$, we conclude it is a generic point. In particular X is sober. Couldn't figure out how to show that the topology base contains only quasi-compact sets.

Problem 7

Sobriety is clear as all closed subsets are finites set of points not containing η .

Thus each irreducible closed subset admits a unique generic point which is itself. Also all open sets are quasi-compact as for any open covering (U_i) , let

$$U = \bigcup_i U_i$$

then $X \setminus U$ is finite. Thus take any U_1 and as $X \setminus U_1$ is finite and contains $X \setminus U$, we only have to take at most $|X \setminus U_1| - |X \setminus U|$ many more open sets to cover U .

Problem 8

Since all open sets in X are quasi-compact, and in particular $X \setminus \{x\}$ is quasi-compact. Therefore, $\{x\}$ is open in X_{con} thus the topology on X_{con} is discrete in particular $\{\{x\}\}$ is a neighborhood basis.