

# Space, time, and episodic memory: The hippocampus is all over the cognitive map

Arne D. Ekstrom<sup>1,2,3\*</sup> | Charan Ranganath<sup>1,2,3\*</sup> 

<sup>1</sup>Center for Neuroscience, University of California, Davis, 1544 Newton Court, Davis, California

<sup>2</sup>Department of Psychology, University of California, Davis, Davis, California

<sup>3</sup>Neuroscience Graduate Group, University of California, Davis, Davis, California

## Correspondence

Charan Ranganath, Center for Neuroscience, University of California, Davis, 1544 Newton Court, Davis, CA.  
Email: cranganath@ucdavis.edu

## Funding information

NIH/NINDS, Grant/Award Number: NS076856 and NS093052; NSF, Grant/Award Number: BCS-1630296; Office of Naval Research, Grant/Award Number: N00014-15-1-0033

## Abstract

In recent years, the field has reached an impasse between models suggesting that the hippocampus is fundamentally involved in spatial processing and models suggesting that the hippocampus automatically encodes all dimensions of experience in the service of memory. Here, we consider key conceptual issues that have impeded progress in our understanding of hippocampal function, and we review findings that establish the scope and limits of hippocampal involvement in navigation and memory. We argue that space and time serve as a primary scaffold to break up experiences into specific contexts, and to organize multimodal input that is to be associated within a context. However, the hippocampus is clearly capable of incorporating additional dimensions into the scaffold if they are determined to be relevant in the event-defined context. Conceiving of the hippocampal representation as constrained by immediate task demands—yet preferring axes that involve space and time—helps to reconcile an otherwise disparate set of findings on the core function of the hippocampus.

## KEYWORDS

allocentric, cognitive map, egocentric, hippocampus, humans, path integration, spatial navigation

## 1 | "I AM A TRAVELER OF BOTH TIME AND SPACE, TO BE WHERE I HAVE BEEN." ROBERT PLANT

Our understanding of the hippocampus, ranging from analysis of cellular properties up to its contributions to behavior, can be seen as one of the greatest success stories in modern neuroscience. Neuroscience research has tied the role of the hippocampus to spatial navigation, and there are compelling links between studies illustrating the properties of place cells, functional imaging, and neurophysiological recordings in humans during virtual navigation. Such findings provide considerable support for the idea of the hippocampus as representing a cognitive map of the current environment (O'Keefe & Nadel, 1978).

There is also considerable evidence suggesting that the hippocampus contributes to a broad range of behaviors beyond spatial cognition. In rodents and primates, these include cells selective for odors, spatial views of landmarks, auditory cues, and conjunctions of task-specific variables, such as spatial context, landmarks, goals, and odors (Aronov, Nevers, & Tank, 2017; Ekstrom et al., 2003; O'Keefe & Speakman,

1987; Rolls & O'Mara, 1995; Wirth, Baraduc, Plante, Pinede, & Duhamel, 2017; Wood, Dudchenko, & Eichenbaum, 1999). In humans, there is near-consensus agreement that the hippocampus is necessary for episodic memory, and in particular, its contextual components (i.e., "when and where did I eat dinner last night?") (Davachi and Dobbins, 2008; Eichenbaum, Yonelinas, & Ranganath, 2007; Ranganath, 2010). It is unclear, although, whether or how episodic memory is related to a broader "spatial" function within the framework of a cognitive map.

It is also clear that many extrahippocampal regions in humans and (to a lesser extent) in rodents, are sufficient to support spatial navigation and memory. For example, hippocampal lesions result in subtle deficits in spatial navigation in humans compared to the observed effects on episodic memory (Hirshhorn, Grady, Rosenbaum, Winocur, & Moscovitch, 2012; Kolarik et al., 2016; Maguire, Nannery, & Spiers, 2006). Lesions to areas outside of the hippocampus—like precuneus, retrosplenial cortex, parahippocampal cortex, and posterior parietal cortex, appear to have much more profound affects on our ability to navigate (Barrash, 1998; Barrash, Damasio, Adolphs, & Tranel, 2000; Mimori et al., 1998; Takahashi, Kawamura, Shiota, Kasahata, & Hirayama, 1997). Rodents with hippocampal lesions, while severely impaired at solving the Morris Water Maze (a classic task for

\*Each author contributed equally to this work

navigational integrity), show some sparing of navigation, including inference of new navigational trajectories (Day, Weisand, Sutherland, & Schallert, 1999; Pearce, Roberts, & Good, 1998). Thus, space appears to involve a broader array of brain areas than the hippocampus, and this begs the question of what unique role the hippocampus plays in processing space more generally (Eichenbaum, 2017).

It is fair to say that we are at an impasse regarding the function of the hippocampus. Proponents of the cognitive map theory have used evidence from a wide range of tasks, ranging from scene perception to virtual navigation, and even prospective planning, to say that the hippocampus is preferentially or exclusively involved in “spatial processing” (Banta-Lavenex, Colombo, Ribordy-Lambert, & Lavenex, 2014). The strongest version of this view that space is fundamental to all forms of cognition, and thus seemingly non-spatial tasks are really supported by spatial processing (Maguire & Mullally, 2013). On the other side of the spectrum, evidence of “non-spatial” processing in the hippocampus, and extra-hippocampal contributions to space, have been used to refute the cognitive map theory of hippocampal function (Eichenbaum & Cohen, 2014). The strongest version of this view is that the hippocampus is involved in all aspects of declarative memory, and that space is just one of a nearly infinite set of variables that is automatically encoded by the hippocampus (Squire & Cave, 1991).

To move forward, we think it is useful to think about some basic questions that have gotten lost in the debate between the spatial and nonspatial perspectives and attempt to reconceptualize the function of the hippocampus more broadly in cognition. In this review, we will consider some significant questions about the role of the hippocampus in spatial and nonspatial cognition and we will review findings that establish the scope and limits of hippocampal involvement in navigation and memory. We argue that the hippocampus “maps” multimodal input in reference to a 4-D continuous topological representation involving space (3-D) and time (1-D). Critically, this mapping process is not rigid—the specific hippocampal context map that is selected, the relative weighting of spatial and temporal dimensions, and the relevance of other “nonspatial” dimensions depend critically on behavioral relevance and task demands.

## 2 | WHAT IS THE DIFFERENCE BETWEEN “SPATIAL” AND “NONSPATIAL”?

One key stumbling block in the ongoing debate about hippocampal function is that authors rarely clarify what they mean when they say that a behavior, task, or representation, is “spatial” or “nonspatial.” Although physicists can agree on a definition of space, most mammalian species do not have sensory receptors that can directly transduce information about space. As pointed out centuries ago by Immanuel Kant, space must be constructed in relation to a reference frame. Depending on how we navigate, we can consider the positions of objects in relation to our immediate peri-personal space, our larger, unseen environment (i.e., the town we are in), or even larger geographic context (i.e., the state or country we are in), which we may never actually navigate but can readily see on a map (Montello, 1993).

Objects typically have a position in space, which we can define relative to our current body position (termed “egocentric”), or relative to one another (termed “allocentric”). Space also can also be topological, rather than Cartesian, meaning that it can preserve the relative, rather than absolute positions of objects to each other (Poucet, 1993).

Tolman proposed that we construct a cognitive map of space, which refers to a metric, viewer-independent representation of the locations of objects (Tolman, 1948). O’Keefe and Nadel (1978) subsequently argued that the hippocampus accomplishes this basic function via neurons that fire at specific spatial locations during navigation (i.e., “place cells”). O’Keefe and Nadel’s (1978) monograph provides a detailed account of the cognitive map model, which has been helpful in detailing predictions about space. However, one potential limitation with current theoretical models of the hippocampus is a slight tendency to for circularity, such that spatial processing is operationally defined in reference to the variables that happen to be related to hippocampal involvement in a task. For example, some studies have found that the hippocampus shows increased activity during perception, encoding, imagery, or working memory maintenance of scenes (Aly et al., 2013; Barense et al., 2010; Bird et al., 2010; Hassabis et al., 2007a; Preston et al., 2010; Stern et al., 2001; Summerfield et al., 2010), and patients with hippocampal damage appear to be deficient at imagining scenes (Hassabis et al., 2007b; see also Hurley et al., 2011; Mullally et al., 2014). Based on such findings, Maguire and colleagues (Maguire & Mullally, 2013; Mullally & Maguire, 2014) proposed “scene construction theory” (“SCT”), which argues that, “...the hippocampus primarily acts to facilitate the construction of atemporal scenes and in doing so allows the event details of episodic memories and imagined future experiences a foundation on which to reside. In this way, hippocampal-dependent scene construction processes are held to underpin and support episodic memory, predicting the future, spatial navigation, and perhaps even dreaming and mind-wandering...” (Mullally and Maguire, 2014). Such studies of scene encoding and imagery are often seen as supporting the Cognitive Map theory and might be considered extensions of Cognitive Map Theory.

The potential issue here is the attempt to accommodate human episodic memory and prospective thinking within the broader umbrella of spatial processing and thus as fundamentally supported by the hippocampus. Specifically, in the attempt to accommodate a wide range of behaviors, theories like SCT run the potential risk of reframing space in a manner that is not obviously consistent with the original idea of the cognitive map. Whereas Cognitive Map Theory specifically refers to the idea that the hippocampus supports a metric, Euclidean, allocentric representation of space, there is no reason to assume that scene perception, recognition, or imagery should require an allocentric representation. Scenes are routinely considered as “items” in memory studies (Canli et al., 2000; Ryan et al., 2000), and in many studies, scenes include multiple objects. Scenes are also typically experienced egocentrically, and there is no requirement that they maintain metric/Euclidean properties during encoding, retrieval, or visual imagery. For these reasons, Burgess, Maguire, & O’Keefe (2002) stated, “A fundamental distinction exists between simple iconic representations of

single objects or 2D scenes and representations that include knowledge of the 3D locations of the elements of a scene. For example, either might suffice for recognition of a scene as familiar, but the latter would be needed to decide upon a novel shortcut or appreciate what the scene would look like from another point of view." Burgess et al. (2002) went on to expand on Cognitive Map Theory by proposing that, "the parahippocampus supports processing of the spatial information present in visual scenes [and] the hippocampus appears only to be activated in more complex navigational situations." Even here, though, the potential issue is that terms like "simple" (to describe processing within parahippocampal cortex) and "complex" (to describe processing within the hippocampus) are underspecified and while useful in some contexts, lack of the specificity needed for detailed empirical testing of hippocampal function at this stage, given current theoretical and empirical advances in the field.

Of course, theories regarding space, episodic memory, and the hippocampus do not have to be consistent with Cognitive Map Theory, but one potential risk is that, in attempting to accommodate novel findings, such theories become under specified and potentially difficult to falsify. For example, as simply one example, we believe that theories such as SCT are vague about the concept of "space" with regard to scenes. Scenes are by no means a homogenous category, and a picture of a natural scene can differ in countless ways from a picture of an object. Given that SCT does not specify, mechanistically, what the hippocampus does during scene construction, and that it does not provide an explicit definition of a scene, it then becomes challenging to test aspects of it as it relates specifically to space and the hippocampus. More generally, specificity about space and representations that the hippocampus may be involved in are a significant issue in the interpretation of research linking the hippocampus to perception, encoding, and imagination of scenes. There is certainly evidence linking the hippocampus to processing of scenes (e.g., Aly et al., 2013; Barense et al., 2010; Bird et al., 2010; Hassabis et al., 2007a; Preston et al., 2010; Stern et al., 2001; Summerfield et al., 2010), but we believe that mechanistic hypotheses about how the hippocampus represents information in scenes, and space specifically, are critical to moving the field forward. One idea that potentially bridges scene processing with the Cognitive Map Theory is that the hippocampus could encode information in scenes in reference to a location in a spatial context (Georges-Francois et al., 1999), and we will explore this idea further in the next section.

### 3 | HOW IS SPACE MAPPED BY THE HIPPOCAMPUS?

When considering exactly what the hippocampus does, it is useful to think about the characteristics of place cells. Although there are good reasons to draw analogies between place cells and a cognitive map, overwhelming evidence suggests that place cells do not map space in a manner that is akin to a GPS. For instance, many studies have shown that when a rat runs on a linear track, place cells remap (i.e., the place fields for individual cells differ) when a rat runs in one direction compared to another (Gothard, Skaggs, Moore, & McNaughton, 1996;

Redish, Rosenzweig, Bohanick, McNaughton, & Barnes, 2000). Perhaps more fundamentally problematic for the idea of a global Cartesian map of space, place cells also change their preferred location of firing depending on the future goal of the animal (Ainge, Tamosiunaite, Woer-goetter, & Dudchenko, 2007; Ferbinteanu and Shapiro, 2003; Lee, Grif-fin, Zilli, Eichenbaum, & Hasselmo, 2006; Wood, Dudchenko, Robitsek, & Eichenbaum, 2000). When an animal navigates toward a goal location on a T-maze, for instance, different place cells fire depending on whether the goal is on the right or left side of the maze. The results from T-maze and linear track paradigms demonstrate that the same location in space can be referenced to competing hippocampal place cell maps, depending on the behavioral context. Similar findings have also been shown in humans: place cells change where they fire depending on the store that a patient is searching for in virtual reality, and also fire to conjunctions of viewing landmarks, searching for goals, and spatial locations (Ekstrom et al., 2003). These findings make clear that the hippocampus does not indicate precise cartographic location within the world. Instead, it indicates topologically accurate coordinates within a multidimensional context space, which are shaped strongly by the ability of the animal to infer the variables necessary for solving the task.

Even studies of place cell activity during simple random foraging reveal that hippocampal place cell maps are anchored to specific spatial contexts (Muller & Kubie, 1987; Skaggs & McNaughton, 1998; Wilson & McNaughton, 1993). Place cells can indicate the location of an animal in a particular spatial context, but the same cells would have different place fields in a different, nearby context. This raises an interesting question—if place cell codes are bound to specific spatiotemporal contexts rather than global spatial coordinates, how can the hippocampus map relative locations across context boundaries? For instance, if you ring your neighbor's doorbell, and you hear her walk to the door, you might know that she is in close proximity. However, the hippocampal place cell code could not be used to figure this out, because she is outside of your current context. Otherwise, there would be immediate interference between the current location mandated by the hippocampus and the imagined locations. Moreover, it is not clear that the hippocampal place code could be used to determine that your house is closer to your neighbor's house than to a house on the opposite side of the planet.

Our example illustrates the point that the hippocampus does not necessarily represent a global cognitive map, but rather that it represents a series of maps, each tied to a different spatial context. In the case of the T-maze and linear track paradigms, competing maps can even be tied to the same spatial context (see also McKenzie, Robinson, Herrera, Churchill, & Eichenbaum, 2014; McKenzie et al., 2013). Critically, if we conceptualize the hippocampal place code as contextually-driven, rather than globally accurate, we solve the problem of otherwise catastrophic interference of competing place cell maps. This point accords with an idea that is well established in the human spatial navigation literature—we often construct representations of disconnected "microenvironments" without updating our position relative to the global "macroenvironment" (Han & Becker, 2013; Wang & Brockmole, 2003a; Wang & Brockmole, 2003b). In other words, when we navigate

in a building, we encode the locations and features that are useful for remembering locations within the building, with little memory or knowledge of how the landmarks and directions we have chosen relate to outdoor locations or other buildings we have recently explored. The idea that a microcontext—be it a spatial goal or nearby environment that we explore, can remain unconnected with other neighboring contexts is entirely consistent with findings suggesting goal and context remapping in the hippocampus. Rather than providing a systematic map of known space, the hippocampus constructs multiple maps with overlapping content, and the selection of the appropriate map is based on many factors beyond space. In the next section, we focus on the most important factor—time (Ranganath & Hsieh, 2016).

#### 4 | WHAT DOES COGNITIVE MAPPING HAVE TO DO WITH EPISODIC MEMORY?

Just as there is a general consensus that the hippocampus is critical for allocentric spatial memory in the rodent brain, there is near consensus that it is critical for episodic memory in the human brain. Virtually every paper on hippocampal representation of space begins with a perfunctory reference to episodic memory (Bird & Burgess, 2008; Burgess et al., 2002; Buzsaki & Moser, 2013). Buzsaki and Moser (2013) went so far as to equate path integration with episodic memory (despite the fact that the very idea of “integration” implies something greater than a single episode). In most instances, however, the exact relationship between a cognitive map and episodic memory is left to the reader’s imagination. This is probably because it is not clear how a purely spatial code would be sufficient to support episodic memory tasks, such as remembering where you left your glasses *last night*, or where you parked your car *this morning*. These memories would be nearly impossible to retrieve due to interference with countless other memories for the locations of your glasses and keys.

O’Keefe and Nadel (1978), considered this issue, and in accord with Tulving’s (1972) definition of episodic memory, they stated that the hippocampus supports, “memory for items or events within a spatio-temporal context.” Studies of hippocampal “time cells” are consistent with this idea (Eichenbaum, 2014). Much as a population of place cells provides a map of a spatial context, a population of time cells maps the progression of time through a situational context. More specifically, time cells fire at specific time points as an animal experiences a predictable sequence of events, even when the animal remains in the same location (Kraus et al., 2015; MacDonald, Carrow, Place, & Eichenbaum, 2013; MacDonald, LePage, Eden, & Eichenbaum, 2011; Naya and Suzuki, 2011; Pastalkova, Itskov, Amarasingham, & Buzsaki, 2008; Wang, Romani, Lustig, Leonardo, & Pastalkova, 2015). Moreover, just as place cells remap when an animal is moved to a new spatial context, hippocampal time cells “retime” when the temporal structure of a task, or the current behavioral context is changed (MacDonald et al., 2013; MacDonald et al., 2011; Sakon, Naya, Wirth, & Suzuki, 2014). These findings demonstrate that hippocampal time and place cells “map” experiences in relation to a context that is bounded in time and space. Consistent with the data from rodents, our labs have repeatedly shown that


activity patterns in the human hippocampus carry information about the spatiotemporal context in which an object was encountered (Dimsdale-Zucker, Ritchey, Ekstrom, Yonelinas, & Ranganath, 2017; Ekstrom, Copara, Isham, Wang, & Yonelinas, 2011; Hsieh, Gruber, Jenkins, & Ranganath, 2014; Kyle, Smuda, Hassan, & Ekstrom, 2015; Libby, Ragland, & Ranganath, 2016; Lieberman, Kyle, Schedlbauer, Stokes, & Ekstrom, 2017; Zhang & Ekstrom, 2013).

With our current understanding of time cells, we can see how the hippocampus might represent experiences as they unfold within the boundaries of an episode. But what is an “episode”? Research on event segmentation provides one answer to this question. Zacks and colleagues have shown that, when watching a movie or reading a narrative, people tend to parse temporally continuous information into discrete events (Zacks & Swallow, 2007). Just as a large border (such as a wall) leads one to assume a boundary between two spatial contexts, large changes in the continuity of incoming information over time appear to trigger boundaries between events (Speer & Zacks, 2005; Zacks et al., 2001). Interestingly, both event boundaries (Axmacher et al., 2010; Baldassano et al., 2016; Ezzyat and Davachi, 2014; Hsieh et al., 2014) and spatial boundaries (Bird et al., 2010) modulate hippocampal activity. These studies demonstrate how hippocampus often uses temporal information, along with spatial information, to represent task-specific aspects of context (Watrous & Ekstrom, 2014).

Putting together what we have reviewed so far, we propose that the hippocampus defines contexts as chunks of experience that have low variance across space and time (Fuhs & Touretzky, 2007; Zucker & Ranganath, 2015). When the variance across experiences exceeds a threshold, we form a new context by differentiating unique temporal epochs (Fuhs & Touretzky, 2007). According to this view, we can expect the hippocampus to treat any cue that remains stable over a chunk of time spent in a given place as an indicator of the spatial context, and sufficient change in one of these cues can lead one to infer the existence of a new context. As an example, if the configuration of furniture remains the same every time you walk into your office, then those factors will become indicators of the context (or “landmarks”), and hippocampal cells would likely map your ongoing experiences relative to the boundaries of your office (Figure 1a). If, one day, the furniture configuration has changed in your office, the change would trigger the hippocampus to assign distinct context representations to the “old” vs. “new” office configurations (Figure 1b). Additionally, if some pieces of furniture, such as your office chair, change locations continuously, these would be unlikely to serve as indicators of your current spatial context. Instead, the chair would be treated as an item to be mapped in the office context (Figure 1c).

#### 5 | WHEN DOES HIPPOCAMPUS GO BEYOND SPACE AND TIME?

Until now, we have presented a view in which the hippocampus maps people and things to spatial and temporal positions within the boundaries of an event context (Eichenbaum, 2013; O’Keefe & Nadel, 1978; Zucker & Ranganath, 2015). The implication is that hippocampal maps



**FIGURE 1** Factors that can influence hippocampal representations of spatial contexts [Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

are limited to four dimensions (3-D space + 1-D time). Recent studies, however, have shown that hippocampal cells encode dimensional information that extends beyond space and time—as just one example, Aronov et al. (2017) trained rats to perform complex mappings between sounds and manual responses (Aronov et al., 2017). They found that hippocampal cells formed discrete firing fields at particular sound frequencies. These findings, along with findings of selectivity for other nonspatial features (e.g., odors) clearly indicate that the hippocampus can add dimensions beyond space and time to the cognitive map. Functional imaging studies in humans have also indicated that the hippocampus can encode abstract spaces such as social hierarchies (Tavares et al., 2015). Does this mean that time and space are no different than any other variable? Should we assume that the hippocampus simply assigns dimensional axes to all continuous variables, such that a hippocampal event representation has almost infinite dimensionality?

An alternative way of thinking about the issue relates back to our earlier discussion. The hippocampus has a strong bias to encode features within a 4-D spatiotemporal framework, but it also encodes variables that are behaviorally relevant, given the current context. Space and time have a privileged status because they are used to define and learn about the current context. For example, rats in the Aronov et al. experiment were trained to perform a “sound manipulation task” during sustained chunks of time in an operant conditioning chamber. We think it is reasonable to speculate that hippocampal cells encoded information about the spatial context and the temporal structure of task trials from the very beginning, whereas auditory selectivity became apparent only after the animal learned that sound frequencies are behaviorally relevant in the context. The critical point is that the hippocampus uses regularities in the environment across time and space in order to define a context (Eichenbaum, 2013; O’Keefe & Nadel, 1978; Ranganath and Hsieh, 2016), and this initial representation must be established in order to learn what is behaviorally relevant in that context. Once the behaviorally relevant variables are determined, then these dimensions are added to the cognitive map.

Although laboratory rats with limited experience of the world must rely on gradual learning to incorporate new dimensions in the hippocampal map, we suspect that, in adult humans, the process is likely to be much faster. To understand why this might be the case, it is helpful to consider the information that gets *in* to the hippocampus.

Considerable evidence suggests that the hippocampus, along with the diencephalon, closely interacts with a posterior medial network that includes the parahippocampal, retrosplenial, and medial entorhinal cortex (Bergmann, Zur, Bershady, & Kahn, 2016; Kahn, Andrews-Hanna, Vincent, Snyder, & Buckner, 2008; Libby, Ekstrom, Ragland, & Ranganath, 2012; Maass, Berron, Libby, Ranganath, & Duzel, 2015). The posterior medial network, like the hippocampus, is extensively engaged in spatial navigation and episodic memory. Ranganath and Ritchey (2012) and Ritchey, Libby, and Ranganath (2015) proposed that the posterior medial network encodes schemas that specify the gist of the spatial, temporal, and causal relationships that apply within a particular event context. Similar proposals regarding a network basis for spatial navigation have also been proposed (Ekstrom, Arnold, & Iaria, 2014; Ekstrom & Watrous, 2014). There is considerable evidence suggesting that ventromedial prefrontal cortex, possibly via the nucleus reunions (Ito, Zhang, Witter, Moser, & Moser, 2015), plays a critical role in regulating hippocampal encoding of information beyond space and time (Gruber et al., 2017; Navawongse & Eichenbaum, 2013; Place, Farovik, Brockmann, & Eichenbaum, 2016; Young & Shapiro, 2011). Notably, the ventromedial prefrontal cortex is extensively interconnected with the posterior medial network, and prefrontal-hippocampal interactions may be critical for both incorporating new information into neocortical event schema representations (Preston & Eichenbaum, 2013; van Kesteren, Ruiter, Fernandez, & Henson, 2012; Zeithamova & Preston, 2010).

The more general point is that the hippocampus is situated between large networks of brain regions, and its role in cognition can be dynamically configured dependent on the relevant extra-hippocampal brain areas with which it interacts. Interactions with neocortical network “hubs” might be a mechanism for the hippocampus to preferentially emphasize any dimension of information coding. In this way, extra-hippocampal cortical brain hubs specify the dimensions beyond space and time that are relevant for representation by the hippocampus (Schedlbauer, Copara, Watrous, & Ekstrom, 2014; Zhang & Ekstrom, 2013).

## 6 | CLOSING THOUGHTS

At the outset of this article, we stated that the field has reached an impasse between two views of the hippocampus—one that places



spatial mapping as its primary function and another that sees the hippocampus as indiscriminately mapping all dimensions of experience. Both of these views capture essential points—any viable theory of hippocampal function needs to account for the critical role of the hippocampus in spatial cognition and its role in episodic memory (Schiller et al., 2015). Spatial and temporal information are used to break up our experiences into specific events that occurred at specific places (Eichenbaum, 2013; O'Keefe & Nadel, 1978; Ranganath & Hsieh, 2016). Although the hippocampus uses space and time as a primary scaffold for defining contexts, and for organizing incoming information within a context representation (O'Keefe & Nadel, 1978), other dimensions can be incorporated into the scaffold (Eichenbaum & Cohen, 2014) if they are determined to be relevant in the event-defined context. Conceiving of the hippocampal representation as constrained by immediate task demands – yet preferring axes that involve space and time – helps to reconcile an otherwise disparate set of findings on the core function of the hippocampus.

## ACKNOWLEDGMENTS

ADE wishes to acknowledge funding from NIH/NINDS grants NS076856, NS093052 (ADE), and NSF BCS-1630296 and CR acknowledges funding from a Vannevar Bush Fellowship (Office of Naval Research Grant N00014-15-1-0033). Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the Office of Naval Research or the U.S. Department of Defense.

## REFERENCES

- Ainge, J. A., Tamosiunaite, M., Woergoetter, F., & Dudchenko, P. A. (2007). Hippocampal CA1 place cells encode intended destination on a maze with multiple choice points. *Journal of Neuroscience*, 27(36), 9769–9779.
- Aly, M., Ranganath, C., & Yonelinas, A. P. (2013). Detecting changes in scenes: the hippocampus is critical for strength-based perception. *Neuron*, 78(6), 1127–1137.
- Aronov, D., Nevers, R., & Tank, D. W. (2017). Mapping of a non-spatial dimension by the hippocampal-entorhinal circuit. *Nature*, 543(7647), 719–722.
- Axmacher, N., Cohen, M. X., Fell, J., Haupt, S., Dumpelmann, M., Elger, C. E., . . . Ranganath, C. (2010). Intracranial EEG correlates of expectancy and memory formation in the human hippocampus and nucleus accumbens. *Neuron*, 65(4), 541–549.
- Baldassano, C., Chen, J., Zadbood, A., Pillow, J. W., Hasson, U., & Norman, K. A. (2016). Discovering event structure in continuous narrative perception and memory. *bioRxiv*.
- Banta-Lavenex, P., Colombo, F., Ribordy-Lambert, F., & Lavenex, P. (2014). The human hippocampus beyond the cognitive map: Evidence from a densely amnesic patient. *Frontiers in Human Neuroscience*, 8(711), 1–18.
- Barense, M. D., Henson, R. N., Lee, A. C., & Graham, K. S. (2010). Medial temporal lobe activity during complex discrimination of faces, objects, and scenes: Effects of viewpoint. *Hippocampus*, 20(3), 389–401.
- Barrash, J. (1998). A historical review of topographical disorientation and its neuroanatomical correlates. *Journal of Clinical and Experimental Neuropsychology*, 20(6), 807–827.
- Barrash, J., Damasio, H., Adolphs, R., & Tranel, D. (2000). The neuroanatomical correlates of route learning impairment. *Neuropsychologia*, 38(6), 820–836.
- Bergmann, E., Zur, G., Bershadsky, G., & Kahn, I. (2016). The Organization of Mouse and human cortico-hippocampal networks estimated by intrinsic functional connectivity. *Cereb Cortex*, 26(12), 4497–4512.
- Bird, C. M., & Burgess, N. (2008). The hippocampus and memory: insights from spatial processing. *Nature Reviews Neuroscience*, 9(3), 182–194.
- Bird, C. M., Capponi, C., King, J. A., Doeller, C. F., & Burgess, N. (2010). Establishing the boundaries: the hippocampal contribution to imagining scenes. *Journal of Neuroscience*, 30(35), 11688–11695.
- Burgess, N., Maguire, E. A., & O'Keefe, J. (2002). The human hippocampus and spatial and episodic memory. *Neuron*, 35(4), 625–641.
- Buzsaki, G., & Moser, E. I. (2013). Memory, navigation and theta rhythm in the hippocampal-entorhinal system. *Nature Neuroscience*, 16(2), 130–138.
- Canli, T., Zhao, Z., Brewer, J., Gabrieli, J. D., & Cahill, L. (2000). Event-related activation in the human amygdala associates with later memory for individual emotional experience. *Journal of Neuroscience*, 20(19), RC99.
- Davachi, L., & Dobbins, I. G. (2008). Declarative memory. *Current Directions in Psychological Science*, 17(2), 112–118.
- Day, L. B., Weisand, M., Sutherland, R. J., & Schallert, T. (1999). The hippocampus is not necessary for a place response but may be necessary for pliancy. *Behavioral Neuroscience*, 113(5), 914–924.
- Dimsdale-Zucker, H. R., Ritchey, M., Ekstrom, A. D., Yonelinas, A. P., & Ranganath, C. (2017). CA1 And CA3 Differentially Support Spontaneous Retrieval Of Episodic Contexts Within Human Hippocampal Subfields. *bioRxiv*.
- Eichenbaum, H. (2013). Memory on time. *Trends in Cognitive Sciences*, 17(2), 81–88.
- Eichenbaum, H. (2014). Time cells in the hippocampus: a new dimension for mapping memories. *Nature Reviews Neuroscience*, 15(11), 732–744.
- Eichenbaum, H. (2017). The role of the hippocampus in navigation is memory. *Journal of Neurophysiology*, 117(4), 1785–1796.
- Eichenbaum, H., & Cohen, N. J. (2014). Can we reconcile the declarative memory and spatial navigation views on hippocampal function? *Neuron*, 83(4), 764–770.
- Eichenbaum, H., Yonelinas, A. P., & Ranganath, C. (2007). The medial temporal lobe and recognition memory. *Annual Review of Neuroscience*, 30, 123–152.
- Ekstrom, A. D., Arnold, A. E., & Iaria, G. (2014). A critical review of the allocentric spatial representation and its neural underpinnings: Toward a network-based perspective. *Frontiers in Human Neuroscience*, 1–15.
- Ekstrom, A. D., Copara, M. S., Isham, E. A., Wang, W. C., & Yonelinas, A. P. (2011). Dissociable networks involved in spatial and temporal order source retrieval. *Neuroimage*, 56(3), 1803–1813.
- Ekstrom, A. D., Kahana, M. J., Caplan, J. B., Fields, T. A., Isham, E. A., Newman, E. L., & Fried, I. (2003). Cellular networks underlying human spatial navigation. *Nature*, 425(6954), 184–188.
- Ekstrom, A. D., & Watrous, A. J. (2014). Multifaceted roles for low-frequency oscillations in bottom-up and top-down processing during navigation and memory. *Neuroimage*, 85(2), 667–677.
- Ezzyat, Y., & Davachi, L. (2014). Similarity breeds proximity: Pattern similarity within and across contexts is related to later mnemonic judgments of temporal proximity. *Neuron*, 81(5), 1179–1189.
- Ferbinteanu, J., & Shapiro, M. L. (2003). Prospective and retrospective memory coding in the hippocampus. *Neuron*, 40(6), 1227–1239.

- Fuhs, M. C., & Touretzky, D. S. (2007). Context learning in the rodent hippocampus. *Neural Comput*, 19(12), 3173–3215.
- Georges-Francois, P., Rolls, E. T., & Robertson, R. G. (1999). Spatial view cells in the primate hippocampus: Allocentric view not head direction or eye position or place. *Cereb Cortex*, 9(3), 197–212.
- Gothard, K. M., Skaggs, W. E., Moore, K. M., & McNaughton, B. L. (1996). Binding of hippocampal CA1 neural activity to multiple reference frames in a landmark-based navigation task. *J Neurosci*, 16(2), 823–835.
- Gruber, M. J., Hsieh, L.-T., Staesina, B., Elger, C., Fell, J., Axmacher, N., & Ranganath, C. (2017). Theta Phase Synchronization Between The Human Hippocampus And The Prefrontal Cortex Supports Learning Of Unexpected Information. *bioRxiv*.
- Han, X., & Becker, S. (2013). One spatial map or many? Spatial coding of connected environments. *Journal of Experimental Psychology: Learning, Memory, and Cognition*.
- Hassabis, D., Kumaran, D., & Maguire, E. A. (2007a). Using imagination to understand the neural basis of episodic memory. *Journal of Neuroscience*, 27(52), 14365–14374.
- Hassabis, D., Kumaran, D., Vann, S. D., & Maguire, E. A. (2007b). Patients with hippocampal amnesia cannot imagine new experiences. *Proceedings of the National Academy of Sciences of the United States of America*, 104(5), 1726–1731.
- Hassabis, D., & Maguire, E. A. (2007). Deconstructing episodic memory with construction. *Trends in Cognitive Sciences*, 11(7), 299–306.
- Hirshhorn, M., Grady, C., Rosenbaum, R. S., Winocur, G., & Moscovitch, M. (2012). The hippocampus is involved in mental navigation for a recently learned, but not a highly familiar environment: A longitudinal fMRI study. *Hippocampus*, 22(4), 842–852.
- Hsieh, L. T., Gruber, M. J., Jenkins, L. J., & Ranganath, C. (2014). Hippocampal activity patterns carry information about objects in temporal context. *Neuron*, 81(5), 1165–1178.
- Hurley, N. C., Maguire, E. A., & Vargha-Khadem, F. (2011). Patient HC with developmental amnesia can construct future scenarios. *Neuropsychologia*, 49(13), 3620–3628.
- Ito, H. T., Zhang, S. J., Witter, M. P., Moser, E. I., & Moser, M. B. (2015). A prefrontal-thalamo-hippocampal circuit for goal-directed spatial navigation. *Nature*, 50–55.
- Kahn, I., Andrews-Hanna, J. R., Vincent, J. L., Snyder, A. Z., & Buckner, R. L. (2008). Distinct cortical anatomy linked to subregions of the medial temporal lobe revealed by intrinsic functional connectivity. *Journal of Neurophysiology*, 100(1), 129–139.
- Kolarik, B. S., Shahlaie, K., Hassan, B., Borders, A. A., Kaufman, K., Gurkoff, G., ... Ekstrom, A. D. (2016). Impairments in precision, rather than spatial strategy, characterize performance on the virtual Morris Water Maze: A case study. *Neuropsychologia*, 80, 90–101.
- Kraus, B. J., Brandon, M. P., Robinson, R. J., 2nd, Connerney, M. A., Hasselmo, M. E., & Eichenbaum, H. (2015). During running in place, grid cells integrate elapsed time and distance run. *Neuron*, 88(3), 578–589.
- Kyle, C. T., Smuda, D. N., Hassan, A. S., & Ekstrom, A. D. (2015). Roles of human hippocampal subfields in retrieval of spatial and temporal context. *Behavioural Brain Research*, 278C, 549–558.
- Lee, I., Griffin, A. L., Zilli, E. A., Eichenbaum, H., & Hasselmo, M. E. (2006). Gradual translocation of spatial correlates of neuronal firing in the hippocampus toward prospective reward locations. *Neuron*, 51(5), 639–650.
- Libby, L. A., Ekstrom, A. D., Ragland, J. D., & Ranganath, C. (2012). Differential connectivity of perirhinal and parahippocampal cortices within human hippocampal subregions revealed by high-resolution functional imaging. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 32(19), 6550–6560.
- Libby, L. A., Ragland, J. D., & Ranganath, C. (2016). The Hippocampus Generalizes Across Memories that Share Item and Context Information. *bioRxiv*.
- Lieberman, J. S., Kyle, C. T., Schedlbauer, A., Stokes, J., & Ekstrom, A. D. (2017). A tale of two temporal coding strategies: Common and dissociable brain regions involved in recency versus associative temporal order retrieval strategies. *Journal of Cognitive Neuroscience*, 29(4), 739–754.
- Maass, A., Berron, D., Libby, L. A., Ranganath, C., & Duzel, E. (2015). Functional subregions of the human entorhinal cortex. *Elife*, 4, 1–20.
- MacDonald, C. J., Carrow, S., Place, R., & Eichenbaum, H. (2013). Distinct hippocampal time cell sequences represent odor memories in immobilized rats. *Journal of Neuroscience*, 33(36), 14607–14616.
- MacDonald, C. J., Lepage, K. Q., Eden, U. T., & Eichenbaum, H. (2011). Hippocampal “time cells” bridge the gap in memory for discontinuous events. *Neuron*, 71(4), 737–749.
- Maguire, E. A., & Mullally, S. L. (2013). The hippocampus: a manifesto for change. *J Exp Psychol Gen.*, 142(4), 1180–1189.
- Maguire, E. A., Nannery, R., & Spiers, H. J. (2006). Navigation around London by a taxi driver with bilateral hippocampal lesions. *Brain*, 129(Pt 11), 2894–2907.
- McKenzie, S., Frank, A. J., Kinsky, N. R., Porter, B., Riviere, P. D., & Eichenbaum, H. (2014). Hippocampal representation of related and opposing memories develop within distinct, hierarchically organized neural schemas. *Neuron*, 83(1), 202–215.
- McKenzie, S., Robinson, N. T., Herrera, L., Churchill, J. C., & Eichenbaum, H. (2013). Learning causes reorganization of neuronal firing patterns to represent related experiences within a hippocampal schema. *Journal of Neuroscience*, 33(25), 10243–10256.
- Mimori, T., Tsuzuki, Y., Zeki, S., Yamaguchi, K., Hayashi, K., Takata, A., & Kuramochi, S. (1998). A case of polyarteritis nodosa who developed severe pneumonia [clinical conference]. *The Keio Journal of Medicine*, 47(4), 223–233.
- Montello, D. R. (1993). Scale and multiple psychologies of space. In: A. U. Frank & I. Campari (Eds.), *Spatial Information Theory: A Theoretical Basis for GIS*. Berlin: Springer-Verlag Lecture Notes in Computer Science.
- Mullally, S. L., & Maguire, E. A. (2014). Memory, imagination, and predicting the future: A common brain mechanism? *Neuroscientist*, 20(3), 220–234.
- Mullally, S. L., Vargha-Khadem, F., & Maguire, E. A. (2014). Scene construction in developmental amnesia: an fMRI study. *Neuropsychologia*, 52, 1–10.
- Muller, R. U., & Kubie, J. L. (1987). The effects of changes in the environment on the spatial firing of hippocampal complex-spike cells. *Journal of Neuroscience*, 7(7), 1951–1968.
- Navawongse, R., & Eichenbaum, H. (2013). Distinct pathways for rule-based retrieval and spatial mapping of memory representations in hippocampal neurons. *Journal of Neuroscience*, 33(3), 1002–1013.
- Naya, Y., & Suzuki, W. A. (2011). Integrating what and when across the primate medial temporal lobe. *Science*, 333(6043), 773–776.
- O’Keefe, J., & Nadel, L. (1978). *The Hippocampus as a Cognitive Map*. London: Oxford University Press.
- O’Keefe, J., & Speakman, A. (1987). Single unit activity in the rat hippocampus during a spatial memory task. *Experimental Brain Research*, 68(1), 1–27.
- Pastalkova, E., Itskov, V., Amarasingham, A., & Buzsaki, G. (2008). Internally generated cell assembly sequences in the rat hippocampus. *Science*, 321(5894), 1322–1327.

- Pearce, J. M., Roberts, A. D., & Good, M. (1998). Hippocampal lesions disrupt navigation based on cognitive maps but not heading vectors. *Nature*, 396(6706), 75–77.
- Place, R., Farovik, A., Brockmann, M., & Eichenbaum, H. (2016). Bidirectional prefrontal-hippocampal interactions support context-guided memory. *Nature Neuroscience*, 19(8), 992–994.
- Poucet, B. (1993). Spatial cognitive maps in animals: New hypotheses on their structure and neural mechanisms. *Psychol Rev*, 100(2), 163–182.
- Preston, A. R., Bornstein, A. M., Hutchinson, J. B., Gaare, M. E., Glover, G. H., & Wagner, A. D. (2010). High-resolution fMRI of content-sensitive subsequent memory responses in human medial temporal lobe. *Journal of Cognitive Neuroscience*, 22(1), 156–173.
- Preston, A. R., & Eichenbaum, H. (2013). Interplay of hippocampus and prefrontal cortex in memory. *Current Biology*, 23(17), R764–R773.
- Ranganath, C. (2010). A unified framework for the functional organization of the medial temporal lobes and the phenomenology of episodic memory. *Hippocampus*, 20(11), 1263–1290.
- Ranganath, C., & Hsieh, L. T. (2016). The hippocampus: A special place for time. *Annals of the New York Academy of Sciences*, 1369(1), 93–110.
- Ranganath, C., & Ritchey, M. (2012). Two cortical systems for memory-guided behaviour. *Nature Reviews Neuroscience*, 13(10), 713–726.
- Redish, A. D., Rosenzweig, E. S., Bohanick, J. D., McNaughton, B. L., & Barnes, C. A. (2000). Dynamics of hippocampal ensemble activity realignment: time versus space. *Journal of Neuroscience*, 20(24), 9298–9309.
- Ritchey, M., Libby, L. A., & Ranganath, C. (2015). Cortico-hippocampal systems involved in memory and cognition: The PMAT framework. *Progress in Brain Research*, 219, 45–64.
- Rolls, E. T., & O'mara, S. M. (1995). View-responsive neurons in the primate hippocampal complex. *Hippocampus*, 5(5), 409–424.
- Ryan, J. D., Althoff, R. R., Whitlow, S., & Cohen, N. J. (2000). Amnesia is a deficit in relational memory. *Psychological Science*, 11(6), 454–461.
- Sakon, J. J., Naya, Y., Wirth, S., & Suzuki, W. A. (2014). Context-dependent incremental timing cells in the primate hippocampus. *Proceedings of the National Academy of Sciences of the United States of America*, 111(51), 18351–18356.
- Schedlbauer, A. M., Copara, M. S., Watrous, A. J., & Ekstrom, A. D. (2014). Multiple interacting brain areas underlie successful spatiotemporal memory retrieval in humans. *Sci Rep*, 4, 6431.
- Schiller, D., Eichenbaum, H., Buffalo, E. A., Davachi, L., Foster, D. J., Leutgeb, S., & Ranganath, C. (2015). Memory and space: Towards an understanding of the cognitive map. *Journal of Neuroscience*, 35(41), 13904–13911.
- Skaggs, W. E., & McNaughton, B. L. (1998). Spatial firing properties of hippocampal CA1 populations in an environment containing two visually identical regions. *Journal of Neuroscience*, 18(20), 8455–8466.
- Speer, N. K., & Zacks, J. A. (2005). Temporal changes as event boundaries: Processing and memory consequences of narrative time shifts. *Journal of Memory and Language*, 53(1), 125–140.
- Squire, L. R., & Cave, C. B. (1991). The hippocampus, memory, and space. *Hippocampus*, 1(3), 269–271.
- Stern, C. E., Sherman, S. J., Kirchoff, B. A., & Hasselmo, M. E. (2001). Medial temporal and prefrontal contributions to working memory tasks with novel and familiar stimuli. *Hippocampus*, 11, 337–346.
- Summerfield, J. J., Hassabis, D., & Maguire, E. A. (2010). Differential engagement of brain regions within a "core" network during scene construction. *Neuropsychologia*, 48(5), 1501–1509.
- Takahashi, N., Kawamura, M., Shiota, J., Kasahata, N., & Hirayama, K. (1997). Pure topographic disorientation due to right retrosplenial lesion. *Neurology*, 49(2), 464–469.
- Tavares, R. M., Mendelsohn, A., Grossman, Y., Williams, C. H., Shapiro, M., Trope, Y., & Schiller, D. (2015). A map for social navigation in the human brain. *Neuron*, 87(1), 231–243.
- Tolman, E. C. (1948). Cognitive maps in rats and men. *Psychological Review*, 55, 189–208.
- Tulving, E. (1972). Episodic and semantic memory. In: E. Tulving & W. Donaldson (Eds.), *Organization of Memory* New York (pp. 382–402). New York: Academic Press.
- van Kesteren, M. T., Ruiters, D. J., Fernandez, G., & Henson, R. N. (2012). How schema and novelty augment memory formation. *Trends Neuroscience*, 35(4), 211–219.
- Wang, R. F., & Brockmole, J. R. (2003a). Simultaneous spatial updating in nested environments. *Psychonomic Bulletin & Review*, 10(4), 981–986.
- Wang, R. X. F., & Brockmole, J. R. (2003b). Human navigation in nested environments. *Journal of Experimental Psychology-Learning Memory and Cognition*, 29(3), 398–404.
- Wang, Y., Romani, S., Lustig, B., Leonardo, A., & Pastalkova, E. (2015). Theta sequences are essential for internally generated hippocampal firing fields. *Nature Neuroscience*, 18(2), 282–288.
- Watrous, A. J., & Ekstrom, A. D. (2014). The spectro-contextual encoding and retrieval theory of episodic memory. *Frontiers in Human Neuroscience*, 8, 1–15.
- Wilson, M. A., & McNaughton, B. L. (1993). Dynamics of the hippocampal ensemble code for space. *Science*, 261(5124), 1055–1058.
- Wirth, S., Baraduc, P., Plante, A., Pinede, S., & Duhamel, J. R. (2017). Gaze-informed, task-situated representation of space in primate hippocampus during virtual navigation. *PLoS Biology*, 15(2), e2001045.
- Wood, E. R., Dudchenko, P. A., & Eichenbaum, H. (1999). The global record of memory in hippocampal neuronal activity. *Nature*, 397(6720), 613–616.
- Wood, E. R., Dudchenko, P. A., Robitsek, R. J., & Eichenbaum, H. (2000). Hippocampal neurons encode information about different types of memory episodes occurring in the same location. *Neuron*, 27(3), 623–633.
- Young, J. J., & Shapiro, M. L. (2011). Dynamic coding of goal-directed paths by orbital prefrontal cortex. *J Neurosci*, 31(16), 5989–6000.
- Zacks, J. M., Braver, T. S., Sheridan, M. A., Donaldson, D. I., Snyder, A. Z., Ollinger, J. M., ... Raichle, M. E. (2001). Human brain activity time-locked to perceptual event boundaries. *Nature Neuroscience*, 4(6), 651–655.
- Zacks, J. M., & Swallow, K. M. (2007). Event segmentation. *Current Directions in Psychological Science*, 16(2), 80–84.
- Zeithamova, D., & Preston, A. R. (2010). Flexible memories: Differential roles for medial temporal lobe and prefrontal cortex in cross-episode binding. *Journal of Neuroscience*, 30(44), 14676–14684.
- Zhang, H., & Ekstrom, A. (2013). Human neural systems underlying rigid and flexible forms of allocentric spatial representation. *Human Brain Mapping*, 34(5), 1070–1087.
- Zucker, H. R., & Ranganath, C. (2015). Navigating the human hippocampus without a GPS. *Hippocampus*, 25(6), 697–703.

**How to cite this article:** Ekstrom AD, Ranganath C. Space, time, and episodic memory: The hippocampus is all over the cognitive map. *Hippocampus*. 2018;28:680–687. <https://doi.org/10.1002/hipo.22750>