

Orthopolity as a resource-allocation hypothesis

Reference measures, counterexamples, and frozen-forecast tests

Research manuscript, revised 3 October 2026. The analyses of Sections 3 and 4 are exploratory or governed by a locally recorded plan. Section 5 distinguishes forecasts frozen before outcome decoding from a plant transfer study frozen after raw outcome exposure. None is an externally preregistered study. Authorship and submission declarations require agreement before submission. Historical drafts remain in Git history. Computational details accompany the manuscript in `evidence.md` and `source-audit.md`.

Abstract

Orthopolity proposes that equally allocated resources can produce unequal abundances: classes whose objects require more resource contain fewer objects. We reconstruct this idea as a conditional allocation hypothesis and distinguish it from a universal explanation of power laws. The defining accounting identity depends on a reference measure. With resource cost proportional to size raised to power d , equal resource per logarithmic size interval implies a count-density exponent $d + 1$; equal resource per linear interval implies exponent d . Conservation alone selects neither allocation. Exploratory assessments of earthquake and solar-flare catalogues, an ocean biomass reconstruction and 1,300 aquatic size-spectrum estimates contradict the specified physical proxies and find aquatic slopes near the logarithmic prediction without establishing equivalence. Centred slopes, fitted dispersion and conservation cannot establish ensemble equipartition. We then test frozen forecasts, disclosing prior outcome exposure. In an engineered CPU-allocation system, independently measured non-unit costs predict allocation profiles and their response to a restriction. On an unused solar year, logarithmic neutrality predicts counts better than linear neutrality, but neutrality itself remains unresolved. In 30 size-selected algal lineages regrown in one medium, carrying capacity in biovolume is constant across a 10-fold range of cell volume, so abundance scales inversely with volume, as budget closure with a geometric cost predicts. In grazed food-web chemostats, a two-budget cost-ratio rule fails to predict pulse-driven redistribution. In directly weighed plant censuses, a flat logarithmic biomass template ranks second of five transferred models, while all three trained models predict counts better. Finite-census calibration distinguishes normalized expected stocks from expected normalized census shares; ecological neutrality remains unresolved. Resource symmetry supplies a possible physical postulate for equal expected allocations. The evidence supports conditional budget-closure forecasts under declared resource conditions and a framework for testing allocation; it does not establish neutrality across coexisting natural size classes or a new natural or cosmological law.

1. Introduction

An allocation can be equal when counted in resources and unequal when counted in objects. For example, equal biomass in successive body-mass classes requires fewer large organisms. This observation motivates the term *orthopolity* in Fabbri’s 2024 essay, following an earlier manuscript by Fabbri and Oliveira Jr. (2017). The earlier manuscript assumes uniform resource allocation across component-wealth values; the later essay proposes a much broader natural and

cosmological principle. The useful question is which allocation statement can be defined and tested independently of the abundance distribution it seeks to explain.

The ecological precedent is substantial. Sheldon, Prakash and Sutcliffe (1972) studied ocean particle size distributions; Gaedke (1993) explicitly related equal biomass in logarithmic classes to normalized biomass spectra; Hatton et al. (2021) reconstructed the ocean spectrum from bacteria to whales. Arranz et al. (2022) studied variation among hundreds of lake fish communities and its ecological correlates. Variation among individual systems is therefore already a research topic. Energetic equivalence (Damuth, 1981) is related, but its species-population sampling unit differs from community abundance per size interval. Species richness across sizes prevents automatic interchange of those statements.

Power laws arise under different generative mechanisms, and apparent straight lines in logarithmic plots do not establish a power-law distribution (Newman, 2005; Clauset, Shalizi and Newman, 2009). Rewriting a distribution as inverse resource cost does not identify a new mechanism. Our contribution is a critical reconstruction: we specify the measure and resource, derive their conditional consequences, give counterexamples to stronger interpretations, assess what the repository’s existing data can support, and test frozen forecasts with their exposure history retained. We make no priority claim for the accounting identity or for ecological biomass equivalence.

2. Mathematical formulation

2.1 Objects, resources, and a reference measure

Let k be a positive size coordinate on a finite domain $D = [a, b]$, with $0 < a < b < \infty$. Choose a reference measure μ , with $0 < \mu(D) < \infty$, for the classes being compared. Let $n_\mu(k) = dN/d\mu$ denote object abundance and $\bar{q}(k) = E[q \mid k] > 0$ the conditional arithmetic mean of an additive resource per object. Then

$$\mathcal{O}_\mu(k) = \frac{dR}{d\mu} = \bar{q}(k)n_\mu(k). \quad (1)$$

For an observed class, the corresponding identity is simply $\sum_i q_i = N_{\text{class}}\bar{q}_{\text{class}}$. Resources may be a standing stock, such as biomass, or an accumulated event quantity over a declared observation period. The latter is not a stock resident in a size class, and neither is a flux through size classes. Additivity is needed for (1); a dynamical conservation law is not.

Define **orthopolity relative to μ** , denoted $\mathcal{O}_\mu$, by

$$\mathcal{O}_\mu(k) = C > 0 \quad \text{for } \mu\text{-almost every } k \in D. \quad (2)$$

This is an assumption about allocation. It gives $n_\mu = C/\bar{q}$. Equivalently, if an object is sampled uniformly from the population, its size density with respect to μ is

$$p_N(k) = \frac{1}{Z\bar{q}(k)}, \quad Z = \int_D \frac{d\mu(k)}{\bar{q}(k)} < \infty. \quad (3)$$

Sampling instead in proportion to the resource held gives $p_R(k) = \bar{q}(k)p_N(k)/E_N[q] = 1/\mu(D)$. Thus uniform resource-weighted size and inverse-cost object abundance are equivalent statements, under the same reference measure. This is an accounting equivalence, not a derivation of the allocation assumption. The count, resource total, and constants also satisfy $N = CZ$ and $R = C\mu(D)$. For a finite observed population, the identity is exact for class totals; a smooth density is a population model or approximation.

2.2 The measure changes the exponent

Write $u = \ln(k/k_0)$ for a fixed reference size k_0 . For $\bar{q}(k) = q_0(k/k_0)^d$, with $d > 0$, two different hypotheses give:

Equal resource with respect to	Resource spectrum	Count density with respect to dk
Linear size, $d\mu = dk$	$\bar{q}(k)dN/dk = C$	$dN/dk \propto k^{-d}$
Logarithmic size, $d\mu = du = dk/k$	$\bar{q}(k)dN/du = C$	$dN/dk \propto k^{-(d+1)}$

The second row follows because $dN/du = k dN/dk$. For biomass with $q = k = m$, the predictions are consequently density exponents 1 and 2, respectively. The 2017 manuscript's explicit uniform density over a linear wealth interval is not interchangeable with the logarithmic convention used in the empirical analyses below. Adopting logarithmic classes is a substantive specification, motivated by size-spectrum practice.

Logarithmic allocation is unchanged by changing size units or logarithm base, apart from a constant density multiplier. It also survives $z = (k/k_0)^c$ for constant $c > 0$, since $d \ln z = c d \ln k$. It is not invariant under arbitrary nonlinear coordinates. In general, $dR/dv = (dR/du)|du/dv|$. Choosing a transformation after observing R can therefore manufacture flatness; the coordinate and measure must be fixed independently.

Under $O_{\log}$ and the power-cost assumption, the slope of abundance per log interval is $-d$, the ordinary density exponent is $d + 1$, and the survival function on $[a, b]$ is

$$P(K \geq k) = \frac{k^{-d} - b^{-d}}{a^{-d} - b^{-d}}. \quad (4)$$

The familiar survival exponent d and rank-size exponent $1/d$ are approximations away from the upper boundary, not exact finite-domain identities. For an unbounded domain with $d > 0$, count can be finite while logarithmically uniform resource has infinite total. Finite boundaries are essential to a finite-budget interpretation.

2.3 Allocation postulates and limits of the accounting

Resource symmetry as a candidate law. The proposed rationale is that, absent restrictions favouring particular resource costs, allocation should not distinguish those costs. A natural law can adopt this as a physical postulate without a deeper mechanistic derivation. One precise sufficient formulation uses K declared classes with fixed positive per-object costs q_i , allocations $R_i = q_i N_i$, and fixed total $\sum_i R_i = R > 0$. If the probability law of $(R_1, \dots, R_K)$ is invariant under exchanging allocations across the fixed classes, then symmetry and the sum give

$$E[R_i] = R/K, \quad E[N_i] = R/(K q_i).$$

This postulate selects resource allocations as the symmetric quantities; symmetry of counts would give a different prediction. It is stronger than conservation and is not implied by an unspecified absence of constraints. Indivisible objects may require an approximation where exact exchangeability is infeasible. The class measure also matters: with cost itself as coordinate, equal resource per dq predicts $dE[N]/dq \propto q^{-1}$, whereas equal resource per $d \ln q$ predicts $dE[N]/dq \propto q^{-2}$. Transformation-based treatments of indifference likewise require a specified symmetry (Jaynes, 1968); an inferential rule does not itself establish a physical law.

Equal expectation does not establish equal snapshots or relaxation toward equality. In a divisible-resource idealization, assigning all resource to one uniformly chosen class has the required symmetry and unequal allocation in every realization. Time-average equality needs an additional condition connecting time averages to expectations or its own empirical postulate. A test must therefore specify what "tends to allocate" means and identify eligible systems through conditions independent of their observed allocation. A deeper mechanism is optional for an empirical law; the scope and predictive evidence are not.

Conservation does not imply equal allocation. On any bounded logarithmic domain, $\mathcal{O}(u) = A \exp(\beta u)$ can be normalized to the same resource total for every β . Conservation fixes the integral and leaves its shape undetermined. Likewise, maximum entropy over finitely many equally weighted object classes, constrained by a mean cost, yields $p_j \propto \exp(-\lambda q_j)$. Different reference measures or constraints give different distributions (Jaynes, 1957; Visser, 2013).

Equipartition does not imply a power law without a cost law. If $\bar{q}(k) = q_0 \exp(k/k_0)$, then $O_{\log}$ gives $dN/du \propto \exp(-k/k_0)$ and $dN/dk \propto \exp(-k/k_0)/k$. Neither is a power law. Geometric dimension supplies d only in a specified geometric model such as similar objects with volume proportional to length cubed. It does not establish the number of independent resource inputs in an arbitrary system.

Independent inputs do not generally add tail exponents. For example, two independent unit-scale Pareto variables with densities hx^{-h-1} , $x \geq 1$, $h > 0$, have product density $h^2 z^{-h-1} \ln z$, $z \geq 1$. Integration over the possible factorizations introduces a logarithmic factor. This limits the independent-input argument in the 2017 essay.

The resource must not be fitted to force the conclusion. For any positive abundance, defining $\bar{q} = C/n_\mu$ makes (2) hold identically. A test must instead specify the resource through measurements, definitions, or an independently justified cost model. An unrestricted residual multiplier called "friction" is equally uninformative unless its form or parameters are predicted independently.

2.4 What an ensemble statement would mean

Suppose a system has a power-shaped resource spectrum $\mathcal{O}_i(u) = A_i \exp(\beta_i u)$. A proposed ensemble condition $E[\beta_i] = 0$ concerns mean slopes. It does not imply $E[\mathcal{O}_i(u)]$ is constant. With equal $A_i = A$ and $\beta_i \sim N(0, \tau^2)$,

$$E[\mathcal{O}_i(u)] = A \exp(\tau^2 u^2 / 2), \quad (5)$$

which is curved whenever $\tau > 0$. Equal-total normalization changes this formula but does not make mean-zero slopes sufficient: averaging normalized spectra proportional to $\exp(cu)$ and $\exp(-cu)$ on a symmetric domain produces a nonconstant hyperbolic cosine. A median slope is a different estimand again. A claim about average resource occupancy requires spectra, a common domain, and an explicit weighting of systems; slopes alone do not identify it.

More generally, let $H(u) = \sum_{i=1}^n w_i A_i e^{\beta_i u}$ on a common log-domain, with fixed positive weights and amplitudes. Then

$$H''(u) = \sum_i w_i A_i \beta_i^2 e^{\beta_i u} \geq 0.$$

If any slope is nonzero, the derivative is strictly positive, so the arithmetic mixture cannot be constant on an open interval. Exact flatness requires every contributing slope to be zero. With $p_i(u) = w_i A_i e^{\beta_i u} / H(u)$, its log-curvature is $d^2 \ln H / du^2 = \text{Var}_{p(u)}(\beta)$. Per-system total normalization preserves these conclusions. They are standard convexity consequences for exact power spectra; arbitrary curved profiles can cancel to a flat mean. They therefore neither exclude

a general resource-symmetry law nor convert noisy fitted slopes into a profile test. The scientific assessment gives a quantitative bound for approximate flatness and its weighting restrictions.

Mean and variance also do not require a Gaussian distribution. Gaussianity is an additional maximum-entropy modelling choice, and selecting it by AIC is not a goodness-of-fit test. If $\beta_i \sim N(\mu_\beta, \tau^2)$ and reported slope error is independent $N(0, \sigma_i^2)$, the predicted probability that the *reported slope* lies in $[-c_i, c_i]$ is

$$\Phi_N\left(\frac{c_i - \mu_\beta}{\sqrt{\tau^2 + \sigma_i^2}}\right) - \Phi_N\left(\frac{-c_i - \mu_\beta}{\sqrt{\tau^2 + \sigma_i^2}}\right). \quad (6)$$

The expression $2\Phi_N(c_i/\tau) - 1$ applies to zero-centred *latent* slopes. Comparing it to observed estimates without their errors is a different calculation. Fitting τ on the same slopes makes either comparison an internal diagnostic, not independent validation.

There is a valid simultaneous-resource restriction for an individual system: $\mathcal{O}_1/\mathcal{O}_2 = \bar{q}_1/\bar{q}_2$. Both spectra can be exactly flat only if their mean resource ratio is size-independent. Across systems, however, $\beta_{1i} - \beta_{2i} = d_{1i} - d_{2i}$ for power cost laws. Equal variances and correlation one follow only if that difference is constant and slope variance is nonzero, whether or not equipartition holds. They are not a general test of ensemble equipartition.




Figure 1. Analytic illustrations, not empirical data. (A) With resource equal to size, linear and logarithmic resource equipartition produce different count exponents. Both curves show unit-total resource density in the logarithmic coordinate. (B) Two unit-total resource profiles on a common domain have opposite log-slopes and therefore zero mean log-slope, but their arithmetic mean is not flat. Generated by `run_theory.py`; a vector version is available.

3. Data and methods

3.1 Sources and scope

Data source	Analysed material	Resource and size coordinate
NOAA GOES XRS flare reports, 2022–2024	10,501 catalogue events; 1,168 paired events in the primary evaluation domain	End fluence in J/m ² ; peak irradiance in W/m ²
USGS ComCat, 2010–2024	6,639 events after retaining moment-magnitude types and rounding to 0.1 at threshold 5.5	Magnitude-derived energy proxy as both resource and coordinate
Hatton et al. (2021)	23 one-decade biomass bins, upper 200 m and full water column	Body mass for both
GLOSSAQUA (Ersoy et al., 2025)	1,300 normalized biomass-spectrum estimates from 16 study identifiers	Body mass for both

These are four data sources; the two ocean domains are related reconstructions. GLOSSAQUA contains published estimates rather than organism-level observations. Study identifiers do not guarantee independent authors, methods, sites, or measurement errors. The sources are convenience cases for evaluating a hypothesis, not a representative sample of natural systems. Input bytes are fixed by SHA-256 manifests; provenance and source notices accompany the data.

3.2 Analysis-plan provenance

The pilot configuration explicitly labels itself exploratory. The subsequent aquatic/ocean plan is retained as prereg_2026-09-09.json. In local history, commit 6df364f contains the plan and raw slope data, preceding analysis commit a0c9973. This records an order of commits. It does not establish an immutable public registration, independently verified blinding, or when values were first viewed. Statements of blinding in the historical file are author reports. The current corrections are retrospective, and the later ensemble, stratification, and variance analyses are exploratory. No claim of confirmatory preregistration is needed for the descriptive and mathematical results here.

3.3 Accounting and uncertainty

For bins B_j of logarithmic width w_j , compute $\hat{\mathcal{O}}_j = \sum_{i \in B_j} q_i / w_j$ and $\hat{C} = \sum_j \sum_{i \in B_j} q_i / \sum_j w_j$. The normalized profile $\phi_j = \hat{\mathcal{O}}_j / \hat{C}$ has width-weighted mean one by construction. This is not evidence of flatness. Empty bins remain in the profile, the final boundary is included, and missing resources require an explicit exclusion policy.

A factor- F endpoint drift across log-domain width L corresponds, for a power-shaped spectrum, to slope tolerance $c = \ln(F)/L$. This differs from requiring every $\phi_j \in [1/F, F]$: the latter permits a maximum-to-minimum ratio of F^2 . Both slope and profile deviations matter. A slope equivalence interval combined with a point-estimate profile screen is not full simultaneous equivalence; that would require uncertainty for the whole profile. Failure to establish equivalence is not, by itself, evidence of a nonzero departure.

For flares, the arithmetic mean resource exponent is fitted by Gamma quasi-likelihood on 2022 data, with bounded count-density fitting on 2023–2024 over $[10^{-5}, 10^{-3}]$ W/m². Log-resource least squares would target a different conditional quantity unless an appropriate retransformation were justified. Pilot intervals use 600 calendar-month block resamples for flares and calendar-year

blocks for earthquakes. These do not capture all dependence or selection bias. The temporal split separates estimation samples but does not make the exploratory domain choice a prospective prediction.

Power-law diagnostics use maximum likelihood and a refitted parametric-bootstrap KS statistic on declared support, following the approach of Clauset et al. (2009). The rounded earthquake magnitudes are tested on their integer grid: for $K = (M - M_0)/\Delta$, $P(K = k) = (1 - r)r^k$ with $r = 10^{-b\Delta}$. Continuous-reference KS calibration is unsuitable for these ties. Parametric-bootstrap probabilities remain conditional on an independent-event working model; catalogue clustering is a limitation.

3.4 Aquatic estimates and reconstruction uncertainty

The primary GLOSSAQUA subset requires body mass on the size axis, finite slopes and positive ordered size limits, and the normalized biomass-spectrum convention. For biomass per unit linear mass $B(m) \propto m^t$, equal biomass per log mass requires $t = -1$. We examine $\beta = t + 1$ and each estimate's reported size span. The historical plan uses $F = 1.25$ and $F = 2$, a 2,000-replicate study-block bootstrap for the pooled median, and a joint criterion involving its interval and the fraction of slopes inside their tolerances. That fraction ignores individual slope uncertainty and must not be interpreted as formal individual equivalence. Slope summaries cannot detect within-spectrum curvature. A post-audit sensitivity explicitly excludes StudyID_07 from range-dependent assessment because its reported bounds are physically implausible; no replacement bounds are imputed (§4.4).

For the ocean reconstruction, 20,000 simulated profiles propagate reported multiplicative uncertainty factors using lognormal draws centred on the published log estimates, with log-scale standard deviation $\ln(f)/1.96$. The published bounds match estimate divided and multiplied by f . Independent errors across groups and bins are a working assumption, not a property established by those bounds. The reconstruction is not a sample of independent organisms, and its plateau was selected after inspecting the upper-ocean estimates.

Exploratory random-effects and latent-distribution fits use reported errors where available. Intervals converted to standard errors assume their advertised coverage and approximate normality. Size-spectrum estimators can have biased estimates and miscalibrated intervals (Edwards et al., 2017, 2020). Repeated spectra within studies and estimated error variances further limit inference; independence-based pooled intervals are not primary evidence of a population mean.

4. Results

4.1 Earthquakes contradict the specified energy-proxy allocation

Gutenberg–Richter scaling $N(\geq M) \propto 10^{-bM}$ and proxy $E \propto 10^{\gamma M}$ give resource per log energy proportional to $E^{1-b/\gamma}$. Flatness therefore requires $b = \gamma$. At threshold 5.5, the fitted $b = 0.998$ has year-block 95% interval $[0.973, 1.024]$, far below the chosen $\gamma = 1.5$. Thresholds 6.0 and 6.5 give 0.984 and 0.977. Using conversion exponent 1.44 also leaves a substantial discrepancy.

At threshold 5.5 the discrete KS statistic is approximately 0.0077, with a parametric-bootstrap probability about 0.26. This is non-rejection of the fitted geometric model, not proof that the model is exact. A power-like tail can thus coexist with a resource-proxy allocation that disagrees with $O_{\log}$. No discrete lognormal comparison was conducted. Magnitude-derived energy is not independently measured radiated energy; the finding is restricted to the specified proxy and catalogue, without declustering or a regional completeness model.

4.2 Flares disagree with the joint prediction on the selected domain

The fitted resource exponent is $d = 0.858$ with 95% interval $[0.697, 1.054]$. The implied density exponent is 1.858, compared with 2.239 $[2.085, 2.399]$ in the paired evaluation sample. The estimated gap is 0.382 $[0.125, 0.620]$. A rise-phase fluence sensitivity gives gap 0.411 $[0.208, 0.604]$. This alternate resource has no missing values in the domain, but does not prove the primary end-fluence analysis free of missingness bias.

For all 1,306 in-domain evaluation events, peak irradiance has fitted exponent 2.274 and KS statistic about 0.032; its bootstrap probability is about 0.02. The power law is therefore inadequate under this working test. Its likelihood cannot clearly distinguish it from the fitted lognormal. Exponent mismatch tests the conjunction of equipartition, power-shaped mean cost, transfer between years, and the count model; it cannot isolate equipartition from all those assumptions.

The gaps at four lower cutoffs are -0.480 , 0.191 , 0.382 , and 0.960 . This substantial sensitivity discourages treating one selected range as a universal scaling regime. Fluence is instrument-band irradiance integrated over an event, not total flare energy.

4.3 The ocean reconstruction is compatible with a broad, uneven plateau

The upper-ocean summary reproduces the published abundance slope at approximately -1.039 . Its maximum-to-minimum biomass ratio is 38.8 across all 23 bins, and 1.7 within the selected plateau of 16 bins. The corresponding full-water-column plateau ratio is 4.3. The 23 one-decade bins span 23 decades between outer edges; their centres are 22 decades apart. Similarly, the plateau covers 16 decades between edges and 15 between centres.

Under the stated uncertainty propagation, only 2 of 23 upper-ocean bin intervals lie wholly outside the factor-1.25 band; the corresponding full-column count is 3. These are pointwise intervals without multiplicity adjustment. The remaining intervals can include both near-flat and materially non-flat values, so their overlap with the band establishes neither equality nor absence of departures. Correlated reconstruction errors and post hoc range selection further limit the assessment. These results re-express Hatton et al.’s reconstruction; they do not independently replicate it.

The full upper-ocean range has a fitted biomass log-slope of -0.0392 and a propagated 90% interval approximately $[-0.058, -0.026]$, outside the factor-1.25 slope tolerance of 0.00421. Thus the assumed uncertainty model indicates a systematic gradient even though most individual bin intervals overlap the pointwise band. Pointwise overlap does not make the whole-profile question uninformative; its interpretation remains conditional on the reconstruction error model.

4.4 Aquatic slopes are close in location, without established equivalence

Among 1,300 normalized biomass spectra from 16 study identifiers, the pooled median slope is -1.015 . The study-block 95% interval for its departure from -1 is $[-0.100, 0.010]$. The median slope tolerance is 0.0317 at $F = 1.25$ and 0.0984 at $F = 2$. This interval is not wholly within either band. Under the historical joint criteria, the result is **not supported**.

The reported-slope fractions inside their individual drift tolerances are 0.07846 ($F = 1.25$) and 0.24231 ($F = 2$), counting boundary cases inclusively. These are descriptive compatibility fractions. Their complements are not estimates of the fraction of true spectra violating equipartition, because measurement error and within-spectrum shape are not accounted for.

These historical fractions also depend on a metadata defect: all 377 StudyID_07 records report bounds of 2×10^{-8} to 2×10^{27} pg C, a 35-decade range with a physically implausible upper body

mass of 2×10^{12} kg C. Excluding those records from this range-dependent calculation leaves 923 slopes from 15 studies. Their point-slope compatibility fractions are 9.32% at $F = 1.25$ and 28.93% at $F = 2$. The source-derived replacement bounds are unknown; the original slopes remain in the location summary. The remaining records also require a methods and units audit. Consequently neither version of the pass fraction should be treated as a validated estimate of ecological prevalence.

Two studies supply 1,016 of the 1,300 estimates, approximately 78%. A central estimate near -1 is worth investigating, but it neither establishes a mean of -1 in a defined population nor establishes equal average biomass occupancy. The stronger ensemble interpretation fails on both statistical and mathematical grounds (§2.4).

4.5 Exploratory diagnostics identify further limits

Latent-shape fits and random-effects summaries describe appreciable variation in the error-reporting subset. Their fitted dispersion includes ecological differences, study methods, and potentially unmodelled dependence. An I^2 estimate is not a measured fraction of variance caused by ecosystem differences. AIC rankings among Gaussian, Laplace, and Student families do not establish adequacy or identify ecological classes. Error-aware, matched-sample pass fractions are internal model checks, not independent predictions of individual failure.

The cross-study comparison of 639 lake-fish estimates with 377 records associated with Gaedke’s Lake Constance study does not partition spatial and temporal variance. Different taxa, sampling designs, fitting methods, and measurement uncertainties prevent treating their variance ratio as a bound on one population. Reproducing Arranz et al.’s reported summary statistics from the same underlying observations checks extraction, not independent replication of a dispersion parameter.

Exponent metadata also need source verification. Counts in equal-log bins and abundance divided by linear bin width have slopes differing by one under power scaling. The Perkins et al. (2018) count-bin method illustrates why a database category alone may not identify the estimand. Choosing a mapping because it makes a slope closer to -1 or -2 would bias the test. Secondary conventions remain sensitivity analyses; a complete study-level methods audit is needed before pooling them. Checks of two dominant primary studies do not validate every record in the primary subset.

5. Frozen-forecast tests

The analyses of Section 4 assess data that had already been inspected. We therefore added tests in which the resource, its cost law, the forecasts and the decision rules are fixed before the evaluated outcomes are decoded, and in which whole units are held out: launches, a calendar year, vessels and selection treatments. Each executed study retains its inputs, frozen protocol, archived algorithms, forecasts and results in an append-only run registry. Three kinds of evidence are kept separate: measurements on engineered systems, an unused observation period of a natural catalogue, and published biological experiments reanalysed retrospectively. Published summaries of those experiments had been read, so their reanalyses are retrospective validations, not blinded tests. A further plant study uses fixed scoring and a development-only geographic split after accidental raw outcome exposure. Its freeze cannot establish that those outcomes were unseen.

5.1 Controlled allocation measurements

In a controlled workload experiment, separately measured memory and CPU costs of matrix tasks predicted completion profiles under assigned quota pairs within four tolerances frozen before 672 validation tasks (workload pilot). A fresh launch exposed a transfer boundary: the original forecast failed its unchanged tolerance in two of four conditions, and local recalibration recovered one (workload transfer). The quotas were acceptance criteria, so these tests concern the transfer of cost models, not allocation.

An allocation test requires the system, not the experimenter, to divide a resource. In a Python runtime whose global interpreter lock serializes execution, worker threads running matrix kernels of five sizes competed for one execution resource (dimensionality intervention). Separate calibration gave CPU cost degrees 1.870 and 2.686 for the two kernels. The calibrated mean-cost curves predicted complete CPU and job profiles before 16 allocation trials, with maximum CPU-share errors of 0.0035–0.0063 and job-share total-variation errors of 0.0021–0.0124. Replacing the largest-class thread with a second smallest-class thread produced the predicted CPU-share change within 0.0078. A unit-cost alternative had count errors of 0.126–0.273. This success concerns one engineered allocation mechanism; it identifies no geometric dimension and no tendency of natural systems.

5.2 Calibrated decisions and an unused observation period

Complete-profile equivalence decisions were calibrated on 180,000 independently generated datasets in 90 conditions (profile calibration). Simultaneous bands met the declared gate for dense, bounded, independent blocks at 48 blocks, with minimum coverage 93.95%, but failed at 12 and 24 blocks. With twelve serially dependent blocks, wrong-departure rates reached 43.75%. These failures define observation designs for which no neutrality verdict is available, whatever a nominal interval suggests.

For solar flares, measured rise-phase fluence costs, six logarithmic classes, three forecasts, a tolerance and an application gate were committed before the 2025 NOAA records were acquired (solar validation). Of 3,277 events, 356 met the domain rules. Logarithmic neutrality predicted the class counts much better than linear neutrality, with count total variation 0.125 against 0.573; historical counts did better still, at 0.022. The point profile departed from the $\ln 1.5$ margin, but the calibrated decision remained unresolved, as declared before acquisition, and the instrument mix changed between periods.

5.3 Published biological measurements

Study	Held out	Resource cost	and	Frozen comparison	Outcome
<i>Synechococcus</i> quotas (Harcourt et al., 2024)	Ten cultures at 25 °C	C, N and P per cell against measured diameter		Fixed cube, free power, strain means, temperature trends	Fixed cube best for C; strain means best for N and P
Size-selected <i>Dunaliella</i> (Malerba et al., 2018)	Each selection treatment	Biovolume at carrying capacity in one shared medium		Equal biovolume ($d = 1$), fitted size law, assigned exponents, equal cell number	Equal biovolume best; cells at capacity scale as $V^{-1.02}$
Food-web chemostats (Wojcik et al., 2025)	Twelve polyculture vessels	Algal N stock: biovolume times separately assayed N per volume		Persistence, development response, no-herbivore response, equal stock, two-budget cost ratio	No model beat persistence; the cost-ratio rule failed
Harvested plants (Dillon et al., 2019)	Five Colorado plots; prior raw exposure	Direct above-ground dry mass; $q(m) = m$ by definition		Log and linear neutrality, Ohio histogram, bounded Pareto and Weibull	Pareto best for biomass; trained models better for counts; ecological neutrality unresolved

Published elemental quotas of four *Synechococcus* strains tested whether a size-based cost transfers to an unused temperature (cost transfer). Trained on 16–22 °C cultures, a cubic diameter law with one fitted intercept predicted held-out 25 °C carbon quotas within a typical factor of 1.14. For nitrogen and phosphorus, strain means beat both size laws, and a fitted exponent never beat the fixed cube. Extrapolated temperature trends were worst for every element. A cost calibration can therefore transfer for one resource while failing as a size law for another resource in the same cells.

The accounting identity behind orthopolity, $Nq(V) = R$, predicts how abundance scales with per-object cost when a budget is shared. Thirty *Dunaliella tertiolecta* lineages, artificially selected for about 280 generations into small, control and large classes, were regrown separately in one F/2 medium after replete, N-deprived or P-deprived histories (size budget). The protocol was committed before any small- or large-selected outcome was decoded; each held-out selection treatment was then predicted from the other two. Across a 10.4-fold range of mean cell volume, replete carrying capacity in total biovolume was constant to within 3%, so cell number at capacity scaled as $V^{-1.02}$ (Figure 2). The frozen equal-biovolume law ($d = 1$) had held-out errors of 0.124 and 0.179 log units. It outperformed a size law fitted within part of the range, whose fitted cost dimension was 0.74 in one fold and 1.12 in the other, and an equal-cell-number law, with errors of 1.48–2.08. A carbon cost degree assigned from the *Synechococcus* study ($d = 0.91$) performed comparably; its nitrogen degree ($d = 0.80$) did not. Regrowth after N deprivation nearly restored capacity, whereas P deprivation left an overshoot of 25–31% in control and large lineages but not in small ones. Restoring the medium therefore did not guarantee a return to the replete profile.

In 24 food-web chemostats, a nitrogen pulse redistributed algal resource composition (chemostat response). Pre-pulse N and C per cell volume from separate monocultures converted three algal groups’ biovolumes into resource stocks. Forecasts for twelve held-out polyculture vessels were frozen before their post-pulse values were decoded. The pulse moved resource composition by 0.25 total variation on average, about 1.5 times the pre-pulse variation around baseline, yet no

forecast improved appreciably on persistence: 0.243 for the development response against 0.247. A two-budget allocation rule, in which a binding secondary carbon budget reweights each group's share by its measured C:N ratio, failed on three counts. Its attainable redistribution, 0.05–0.12, was below every observed departure, 0.29–0.49. Its predicted direction held in 4 of 12 vessels, the chance count. It did not beat persistence. No held-out vessel recovered its pre-pulse composition within 12 days.




Figure 2. Size-selected *Dunaliella* lineages regrown in one shared medium, from the data of Malerba et al. (2018). (a) Replete carrying capacity in total biovolume against mean cell volume, with the two cross-fitted size laws and the equal-biovolume law. (b) Implied cell number at capacity. (c) Held-out errors of the frozen forecasts. (d) Capacity after N or P deprivation against replete capacity. Generated from retained outputs by `report_dunaliella_size_budget.py`.

5.4 Directly weighed coexisting plant stocks

Ten harvested herbaceous plots provide direct aboveground dry masses for coexisting ramets or stem clusters (Dillon et al., 2019; plant study). Forest and desert allometric masses are excluded. Five Ohio plots supply development fits; five Colorado plots supply evaluation. The protocol, algorithms and forecasts were committed before formal scoring, but raw rows had already been exposed during source inspection. This is a retrospective transfer comparison. It identifies neither a limiting nutrient nor an opportunity budget; $q(m) = m$ defines the measured stock rather than testing an independent cost law.

The frozen domain is 0.01–100 g, split into eight half-decade bins with empty classes retained. The upper bound depends only on Ohio masses. Thirty missing mass records, 144 subthreshold ramets and one above-domain ramet remain in the evaluation membership ledger. The latter weighs 112.55 g and accounts for 24.50% of one plot's known biomass. In-domain mass coverage

is 75.48% there and 99.89–99.97% in the other four plots. Scores therefore describe the declared domain; the excluded mass and unknown missing mass cannot be silently absorbed.

The primary score is equal-plot mean total variation between realized normalized biomass and each expected-stock or empirical template. Bounded Pareto scores 0.470, logarithmic neutrality 0.493, linear neutrality 0.507, the development histogram 0.510 and bounded Weibull 0.531 (Figure 3). Pareto beats logarithmic neutrality in three of five plots; large opposing plot differences leave a mean improvement of only 0.023. Count profiles provide a different comparison: the development histogram, Pareto and Weibull score 0.260–0.271, versus 0.412 for linear and 0.495 for logarithmic neutrality. Their binned count log losses are about 1.83, versus 2.08 and 2.42. These are point discrepancies, without a calibrated ecological superiority or equivalence verdict. They also do not contradict the original study’s superior within-site Weibull fits: the models here transfer across locales and share a development-fixed domain.

Synthetic calibration fixes census size and draws masses from m^{-2} on the same domain, so each bin has equal expected biomass before measurement rounding. At 160 independent ramets, mean realized stock TV is 0.441 and every simulated census has an empty bin. The largest bin’s average normalized share is 3.13%, although its normalized expected stock is 12.50%. At 1,280 ramets these values are 0.244, 87.2% and 9.12%. Thus $E[R_i / \sum_j R_j]$ differs from $E[R_i] / \sum_j E[R_j]$. A separately generated iid 95% TV envelope has exceedance rates 6.2% and 5.2%, but repeating masses in blocks of twenty raises these to 97.8% and 99.9%. This stress test is not an ecological dependence model. Field dependence, inclusion and independent regional replication remain unidentified, so the ecological neutrality decision is unresolved. An uneven small census alone cannot settle a claim about expected allocation.

5.5 What the tests establish

The positive results concern conditional budget closure: a separately measured or geometric cost, declared shared resource conditions, and abundance or resource shares predicted from them. The runtime experiment and the algal lineages are the clearest cases; in the lineages the closure holds with cost proportional to volume, so the abundance gradient carries no information beyond geometry. A rule that goes beyond closure, by predicting how a restriction redistributes resource among coexisting classes, failed in the one natural community tested, where grazing and fast dynamics dominated. The plant study now observes coexisting class stocks directly, but it does not establish equal expected allocation: the flat template is not best for biomass or counts, and the ecological observation law is uncalibrated. No natural system here has independently identified neutral eligibility and a design sufficient for the expected-allocation verdict. That remains the decisive missing test.

6. Discussion

The conditional identity is useful because it forces a researcher to name the resource, sampling unit, coordinate, and reference measure before interpreting a slope. It also separates a distributional question from an allocation question: a power-law model can fit without the specified resource being equally allocated. Conversely, equal allocation can hold with a non-power cost law and yield a different abundance shape.

Neither this identity nor the current case studies establishes a universal tendency. The original linear-allocation assumption and the ecological logarithmic-allocation assumption must be treated separately. The physical examples contradict selected proxy versions, while the ecological summaries contain suggestive locations and substantial uncertainty. No result here explains why a dynamical system should approach equal allocation, proves an equilibrium or time arrow, or

Measured plant dry mass: retrospective profile transfer and census calibration




Figure 3. Directly weighed plant profiles and synthetic observation limits. (a) Equal-plot mean Colorado biomass shares and transferred stock templates. (b) Count shares and retained count forecasts. (c) Each plot’s biomass discrepancy and the five means. (d) Mean normalized census shares under iid logarithmic neutrality, compared with normalized expected stocks. All panels read retained outputs; `report_plant_biomass_profile.py` performs presentation only. Prior raw exposure and excluded biomass are retained.

quantifies the work needed to change social inequality. No spacetime model or cosmological prediction has been supplied.

A narrower empirical claim remains possible. It would specify a population of systems, one independently justified resource and measure, a sampling frame, a common or explicitly varying size domain, and an equivalence tolerance with scientific meaning. Raw resource totals would permit testing profile shape as well as slope. A design with multiple sites and repeat visits could separate sources of variation using a model appropriate to its sampling structure. A new dataset could test a frozen conditional prediction, including its uncertainty; numerical proximity to a dispersion fitted on the original sample is not sufficient. Section 5 reports such tests. Their positive results concern budget closure, not neutrality. The plant application additionally shows why normalized snapshots cannot be interchanged with normalized expected stocks without an observation model.

The principal limitation is the absence of independently specified conditions identifying the proposed neutral regime and of data/design sufficient to test its generalization. An empirically supported allocation law need not explain its symmetry through a deeper mechanism. Its predictions must nevertheless specify the resource, class measure, and statistical target. An experiment that weakens an independently measured source of bias could test approach to the neutral prediction, if that convergence is part of the declared hypothesis. Departures cannot simply be relabelled as previously unspecified restrictions. The tests of Section 5 sharpen this limitation. Budget closure with a geometric cost is confirmed in separately grown lineages that share one medium. A cost-ratio rule for redistribution among coexisting classes failed in a grazed community.

Prior ecological work already supplies models and environmental explanations (Cuesta, Delius and Law, 2018; Arranz et al., 2022). Marshall et al. (2022) independently measured metabolic scaling and predicted maximum density and biovolume in evolved bacteria. Cost-to-capacity prediction therefore has direct experimental precedents. Dillon et al. (2019) already established curvature and superior Weibull fits in the plant data; the present application adds transferred stock/count comparisons and observation limits. Further orthopolity research needs a demonstrable contribution relative to those accounts, such as an independently validated conditional prediction or a calibrated diagnostic comparison. A resource-symmetry law could coexist with mechanisms explaining when it applies and how it is approached.

7. Conclusion

Orthopolity can be stated as a mathematically coherent, falsifiable hypothesis once its resource, measure, and domain are fixed. Its inverse-cost consequence is an accounting identity shared with established size-spectrum theory. Resource symmetry provides a coherent postulate for equal expected allocations; a deeper mechanism is not required for it to become an empirically established law. The stronger universal, cosmological, and ensemble-confirmation claims are not justified by the available arguments or evidence. A scientific document is warranted as a critical synthesis with reproducible exploratory tests and explicit counterexamples. Frozen forecasts support the budget-closure form of the identity in an engineered allocation system and in algal lineages sharing one medium. They reject a two-budget cost-ratio rule for short-term redistribution in grazed food webs. Direct coexisting plant stocks give mixed template transfer and an unresolved expected-neutrality verdict. The current original contribution is modest and methodological; transparent execution does not substitute for novelty or discriminating natural-system evidence. A paper announcing a new natural law is not warranted.

Data and code availability

Code, frozen inputs, configurations, and generated results are in this repository. The make all target verifies input checksums, runs tests, and regenerates the empirical analyses; see the README for setup. Pilot estimates originate in results.json; distribution diagnostics in gof.json; aquatic median, compatibility fractions, and ocean intervals in independent.json. Remaining JSON files contain explicitly exploratory diagnostics. The studies of Section 5 are registered in an append-only run registry with their inputs, frozen protocols, archived algorithms and outputs. Plant sources are pinned to author commit defccc3dcbbbf3ba57ff1572377de88fba83ff7f, with acquisition checksums and complete inclusion ledgers. The registry-verify target audits the registry offline, and each study report gives its replay command; source errata found during those audits are recorded there. Source-specific conditions are recorded in data/NOTICE.md; there is no blanket licence for all inputs. The supplied 2017 manuscript was inspected privately and is not redistributed.

References

- Arranz, I., Fournier, B., Lester, N. P., Shuter, B. J., and Peres-Neto, P. R. (2022). Species compositions mediate biomass conservation: The case of lake fish communities. *Ecology*, 103, e3608. <https://doi.org/10.1002/ecy.3608>.
- Clauset, A., Shalizi, C. R., and Newman, M. E. J. (2009). Power-law distributions in empirical data. *SIAM Review*, 51, 661–703. <https://doi.org/10.1137/070710111>.
- Cuesta, J. A., Delius, G. W., and Law, R. (2018). Sheldon spectrum and the plankton paradox: two sides of the same coin—a trait-based plankton size-spectrum model. *Journal of Mathematical Biology*, 76, 67–96. <https://arxiv.org/abs/1607.04158>.
- Damuth, J. (1981). Population density and body size in mammals. *Nature*, 290, 699–700. <https://doi.org/10.1038/290699a0>.
- Dillon, K. T., et al. (2019). On the relationships between size and abundance in plants: beyond forest communities. *Ecosphere*, 10, e02856. <https://doi.org/10.1002/ecs2.2856>.
- Edwards, A. M., Robinson, J. P. W., Plank, M. J., Baum, J. K., and Blanchard, J. L. (2017). Testing and recommending methods for fitting size spectra to data. *Methods in Ecology and Evolution*, 8, 57–67. <https://doi.org/10.1111/2041-210X.12641>.
- Edwards, A. M., Robinson, J. P. W., Blanchard, J. L., Baum, J. K., and Plank, M. J. (2020). Accounting for the bin structure of data removes bias when fitting size spectra. *Marine Ecology Progress Series*, 636, 19–33. <https://doi.org/10.3354/meps13230>.
- Ersoy, Z., et al. (2025). GLOSSAQUA: A global dataset of size spectra across aquatic ecosystems. *Ecology*, 106, e70050. <https://doi.org/10.1002/ecy.70050>.
- Fabbri, R., and Oliveira Jr., O. N. (2017). *A simple model that explains why inequality is ubiquitous*. Supplied manuscript dated 17 March 2017, 13 pp.; publication status unverified.
- Fabbri, R. (2024). The Orthopolity cosmological principle and the Natural distribution law. Author’s essay, 14 August. <https://ttm.github.io/2024/08/14/power.html>.
- Gaedke, U. (1993). Ecosystem analysis based on biomass size distributions: A case study of a plankton community in a large lake. *Limnology and Oceanography*, 38, 112–127. <https://doi.org/10.4319/lo.1993.38.1.0112>.
- Harcourt, R., Garcia, N. S., and Martiny, A. C. (2024). *Synechococcus* batch culture data (cell quotas and ratios (C, N, P), size, and diameter) from laboratory experiments in 2021 to 2022 with related isolates cultured across a range of temperatures. BCO-DMO dataset 926311, version 1. <https://doi.org/10.26008/1912/bco-dmo.926311.1>.
- Hatton, I. A., Heneghan, R. F., Bar-On, Y. M., and Galbraith, E. D. (2021). The global ocean size spectrum from bacteria to whales. *Science Advances*, 7, eabh3732. <https://doi.org/10.1126/>

sciadv.abh3732.

- Jaynes, E. T. (1957). Information theory and statistical mechanics. *Physical Review*, 106, 620–630. <https://doi.org/10.1103/PhysRev.106.620>.
- Jaynes, E. T. (1968). Prior probabilities. *IEEE Transactions on Systems Science and Cybernetics*, 4(3), 227–241. <https://bayes.wustl.edu/etj/articles/prior.pdf>.
- Malerba, M. E., Palacios, M. M., and Marshall, D. J. (2018). Do larger individuals cope with resource fluctuations better? An artificial selection approach. *Proceedings of the Royal Society B*, 285, 20181347. <https://doi.org/10.1098/rspb.2018.1347>. Data: <https://doi.org/10.5061/dryad.4mh47r7>.
- Marshall, D. J., et al. (2022). Long-term experimental evolution decouples size and production costs in *Escherichia coli*. *Proceedings of the National Academy of Sciences*, 119, e2200713119. <https://doi.org/10.1073/pnas.2200713119>.
- Newman, M. E. J. (2005). Power laws, Pareto distributions and Zipf’s law. *Contemporary Physics*, 46, 323–351. <https://arxiv.org/abs/cond-mat/0412004>.
- Perkins, D. M., et al. (2018). Bending the rules: exploitation of allochthonous resources by a top-predator modifies size-abundance scaling in stream food webs. *Ecology Letters*, 21, 1771–1780. <https://doi.org/10.1111/ele.13147>.
- Sheldon, R. W., Prakash, A., and Sutcliffe, W. H., Jr. (1972). The size distribution of particles in the ocean. *Limnology and Oceanography*, 17, 327–340. <https://doi.org/10.4319/lo.1972.17.3.0327>.
- Visser, M. (2013). Zipf’s law, power laws and maximum entropy. *New Journal of Physics*, 15, 043021. <https://arxiv.org/abs/1212.5567>.
- Wojcik, L. A. M., Pfennig, A., Flamm, S., Klauschies, T., Rosenbaum, B., Weithoff, G., and Gaedke, U. (2025). Top-down control and species composition influence nonlinearly the short-term response of experimental food webs to a nutrient pulse perturbation. *Proceedings of the Royal Society B*. <https://doi.org/10.1098/rspb.2025.1969>. Data: <https://doi.org/10.5061/dryad.51c59zwj5>.