

$$\begin{array}{l}
\boxed{\vec{\nabla} = [\vec{\nabla}(r, \theta, \phi)] \bar{\partial}_o} \\
d = [dx \ dy \ dz] \vec{\nabla} = d\vec{l}^T \vec{\nabla} \\
d(r, \theta, \phi) = [dx \ dy \ dz] \vec{\nabla}(r, \theta, \phi) \\
\boxed{\partial \bar{l}_o^T = d\vec{l}^T \vec{\nabla}(r, \theta, \phi)} \\
\partial \bar{l}_o^T \bar{\partial}_o = d\vec{l}^T [\vec{\nabla}(r, \theta, \phi)] \bar{\partial}_o \\
\boxed{d = \partial \bar{l}_o^T \bar{\partial}_o = d\vec{l}^T \vec{\nabla}}
\end{array}
\left| \begin{array}{l}
\vec{\nabla} = \begin{bmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \\ \frac{\partial}{\partial z} \end{bmatrix} = \begin{bmatrix} | & | & | \\ \nabla_r & \nabla_\theta & \nabla_\phi \\ | & | & | \end{bmatrix} \begin{bmatrix} \frac{\partial}{\partial r} \\ \frac{\partial}{\partial \theta} \\ \frac{\partial}{\partial \phi} \end{bmatrix} \\
\partial \bar{l}_o = \begin{bmatrix} dr \\ d\theta \\ d\phi \end{bmatrix} = [\vec{\nabla}(r, \theta, \phi)]^T d\vec{l} \\
= \begin{bmatrix} -\nabla_r - \\ -\nabla_\theta - \\ -\nabla_\phi - \end{bmatrix} \begin{bmatrix} dx \\ dy \\ dz \end{bmatrix}
\end{array} \right|
\begin{array}{l}
\theta = \theta(x, y, z) \quad (x^2 + y^2 = z^2 \tan^2 \theta) \\
\phi = \phi(x, y, z) \quad (y = x \tan \phi) \\
\frac{\partial}{\partial x} = \frac{\partial r}{\partial x} \frac{\partial}{\partial r} + \frac{\partial \theta}{\partial x} \frac{\partial}{\partial \theta} + \frac{\partial \phi}{\partial x} \frac{\partial}{\partial \phi} \\
\frac{\partial}{\partial \theta} = \frac{\partial x}{\partial \theta} \frac{\partial}{\partial x} + \frac{\partial y}{\partial \theta} \frac{\partial}{\partial y} + \frac{\partial z}{\partial \theta} \frac{\partial}{\partial z} \\
d\theta = dx \frac{\partial \theta}{\partial x} + dy \frac{\partial \theta}{\partial y} + dz \frac{\partial \theta}{\partial z} \\
dy_{\vec{r}_o}(\vec{r}'_o)|_{t=0} = (\frac{\partial y}{\partial r} dr + \frac{\partial y}{\partial \theta} d\theta + \frac{\partial y}{\partial \phi} d\phi) \frac{1}{dt} |_{t=0}
\end{array}$$

$$\boxed{\vec{b}^i \cdot \vec{b}_i = \delta_{ij}} : \begin{array}{l} \vec{\nabla} \phi \cdot \frac{\partial \vec{r}}{\partial \phi} = 1 \\ \vec{\nabla} \phi \cdot \frac{\partial \vec{r}}{\partial \theta} = 0 \end{array} \Rightarrow \begin{bmatrix} -\vec{\nabla}_r - \\ -\vec{\nabla}_\theta - \\ -\vec{\nabla}_\phi - \end{bmatrix} \begin{bmatrix} \frac{\partial}{\partial r} & \frac{\partial}{\partial \theta} & \frac{\partial}{\partial \phi} \end{bmatrix} \vec{r} = \mathbb{1}_3 \Rightarrow \boxed{\frac{\partial y}{\partial \phi} = [0 \ 1 \ 0] \begin{bmatrix} -\vec{\nabla}_r - \\ -\vec{\nabla}_\theta - \\ -\vec{\nabla}_\phi - \end{bmatrix}^{-1} \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \frac{\partial y}{\partial \phi}^T}$$

$$\begin{aligned}
d &= dx \frac{\partial}{\partial x} + dy \frac{\partial}{\partial y} + dz \frac{\partial}{\partial z} = d\vec{l} \cdot \vec{\nabla} = \left[\frac{dr}{\|\nabla_r\|}, \frac{d\theta}{\|\nabla_\theta\|}, \frac{d\phi}{\|\nabla_\phi\|} \right] \left[\|\nabla_r\| \frac{\partial}{\partial r}, \|\nabla_\theta\| \frac{\partial}{\partial \theta}, \|\nabla_\phi\| \frac{\partial}{\partial \phi} \right]^T \\
&= dr \frac{\partial}{\partial r} + d\theta \frac{\partial}{\partial \theta} + d\phi \frac{\partial}{\partial \phi} = \partial \bar{l}_o^T \bar{\partial}_o = [dr, r d\theta, r \sin \theta d\phi] \left[\frac{\partial}{\partial r}, \frac{1}{r} \frac{\partial}{\partial \theta}, \frac{1}{r \sin \theta} \frac{\partial}{\partial \phi} \right]^T \\
&= \frac{dr}{\|\nabla_r\|} \|\nabla_r\| \frac{\partial}{\partial r} + \frac{d\theta}{\|\nabla_\theta\|} \|\nabla_\theta\| \frac{\partial}{\partial \theta} + \frac{d\phi}{\|\nabla_\phi\|} \|\nabla_\phi\| \frac{\partial}{\partial \phi} \dots = d\bar{l}_o^T \bar{\nabla}_o = d\vec{l}_o^T \vec{\nabla}_o.
\end{aligned}$$

$$\begin{array}{l}
d\vec{l} = d\vec{r} = [dr \frac{\partial}{\partial r} + d\theta \frac{\partial}{\partial \theta} + d\phi \frac{\partial}{\partial \phi}](x, y, z)^T \\
d(x, y, z) = \left[\frac{dr}{\|\nabla_r\|} \|\nabla_r\| \frac{\partial}{\partial r} + \frac{d\theta}{\|\nabla_\theta\|} \|\nabla_\theta\| \frac{\partial}{\partial \theta} + \frac{d\phi}{\|\nabla_\phi\|} \|\nabla_\phi\| \frac{\partial}{\partial \phi} \right] (x, y, z) \\
(dx, dy, dz) = dr \hat{r}^T + r d\theta \hat{\theta}^T + r \sin \theta d\phi \hat{\phi}^T
\end{array}
\left| \begin{array}{l}
(\hat{r}, \hat{\theta}, \hat{\phi}) \equiv \left(\|\nabla_r\| \frac{\partial \vec{r}}{\partial r}, \|\nabla_\theta\| \frac{\partial \vec{r}}{\partial \theta}, \|\nabla_\phi\| \frac{\partial \vec{r}}{\partial \phi} \right) \\
= \left(\frac{\partial}{\partial r}, \frac{1}{r} \frac{\partial}{\partial \theta}, \frac{1}{r \sin \theta} \frac{\partial}{\partial \phi} \right) \otimes (x, y, z)^T \\
= \vec{\nabla}_o^T \otimes \vec{r}
\end{array} \right.$$

$$\begin{array}{l}
\boxed{d\vec{l} = (dx, dy, dz) \cdot (\hat{x}, \hat{y}, \hat{z}) = (dr, r d\theta, r \sin \theta d\phi) \cdot (\hat{r}, \hat{\theta}, \hat{\phi}) = d\vec{l}_o = d\bar{l}_o^T \cdot (\hat{r}, \hat{\theta}, \hat{\phi})} \\
\boxed{\vec{\nabla} = (\hat{x}, \hat{y}, \hat{z}) \cdot \left(\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right) = (\hat{r}, \hat{\theta}, \hat{\phi}) \cdot \left(\frac{\partial}{\partial r}, \frac{1}{r} \frac{\partial}{\partial \theta}, \frac{1}{r \sin \theta} \frac{\partial}{\partial \phi} \right) = \vec{\nabla}_o = \left(\frac{\partial \vec{r}}{\partial r}, \frac{1}{r} \frac{\partial \vec{r}}{\partial \theta}, \frac{1}{r \sin \theta} \frac{\partial \vec{r}}{\partial \phi} \right) \vec{\nabla}_o^T}
\end{array}$$

$$\begin{aligned}
\vec{\nabla} &= [\vec{\nabla}_o^T \otimes \vec{r}] \vec{\nabla}_o = [\vec{\nabla}_o^T \otimes (x, y, z)^T] \begin{bmatrix} \frac{\partial}{\partial r} \\ \frac{1}{r} \frac{\partial}{\partial \theta} \\ \frac{1}{r \sin \theta} \frac{\partial}{\partial \phi} \end{bmatrix} \Rightarrow \frac{\partial}{\partial x} = \underline{\frac{\partial x}{\partial r}} \frac{\partial}{\partial r} + \underline{\|\nabla_\theta\|^2} \frac{\partial x}{\partial \theta} \frac{\partial}{\partial \theta} + \underline{\|\nabla_\phi\|^2} \frac{\partial x}{\partial \phi} \frac{\partial}{\partial \phi} \\
&= [\vec{\nabla}(r, \theta, \phi)] \bar{\partial}_o = [\vec{\nabla}(r, \theta, \phi)] \begin{bmatrix} \frac{\partial}{\partial r} \\ \frac{\partial}{\partial \theta} \\ \frac{\partial}{\partial \phi} \end{bmatrix} \Rightarrow \frac{\partial}{\partial x} = \underline{\frac{\partial r}{\partial x}} \frac{\partial}{\partial r} + \underline{\frac{\partial \theta}{\partial x}} \frac{\partial}{\partial \theta} + \underline{\frac{\partial \phi}{\partial x}} \frac{\partial}{\partial \phi} \Rightarrow \boxed{\frac{\partial \phi}{\partial y} = \frac{\partial y}{\partial \phi} \|\nabla_\phi\|^2}
\end{aligned}$$

		contravariant _i		(equal since orthog.)		covariant ⁱ	
$\hat{r} = (\hat{r}_x, \hat{r}_y, \hat{r}_z)$	$= \frac{\vec{r}}{r}$	$= \frac{\partial}{\partial r} \vec{r}$	$= \frac{\partial \vec{r}}{\partial r} \ \frac{\partial \vec{r}}{\partial r}\ ^{-1}$	$\stackrel{\rightarrow}{=} \ \nabla_r\ \frac{\partial \vec{r}}{\partial r}$	$\stackrel{\leftarrow}{=} \frac{\nabla_r}{\ \nabla_r\ }$	$= \nabla r$	
$\hat{\theta} = (\hat{\theta}_x, \hat{\theta}_y, \hat{\theta}_z)$	$= \frac{\partial \hat{r}}{\partial \theta}$	$= \frac{1}{r} \frac{\partial}{\partial \theta} \vec{r}$	$= \frac{\partial \vec{r}}{\partial \theta} \ \frac{\partial \vec{r}}{\partial \theta}\ ^{-1}$	$\stackrel{\rightarrow}{=} \ \nabla_\theta\ \frac{\partial \vec{r}}{\partial \theta}$	$\stackrel{\leftarrow}{=} \frac{\nabla_\theta}{\ \nabla_\theta\ }$	$= r \nabla \theta$	
$\hat{\phi} = (\hat{\phi}_x, \hat{\phi}_y, \hat{\phi}_z)$	$= \frac{1}{\sin \theta} \frac{\partial \hat{r}}{\partial \phi}$	$= \frac{1}{r \sin \theta} \frac{\partial}{\partial \phi} \vec{r}$	$= \frac{\partial \vec{r}}{\partial \phi} \ \frac{\partial \vec{r}}{\partial \phi}\ ^{-1}$	$\stackrel{\rightarrow}{=} \ \nabla_\phi\ \frac{\partial \vec{r}}{\partial \phi}$	$\stackrel{\leftarrow}{=} \frac{\nabla_\phi}{\ \nabla_\phi\ }$	$= r \sin \theta \nabla \phi$	

1 Frenet Equations

$(a \times b) \cdot (c \times d) = a \cdot b \times (c \times d)$ $= \begin{vmatrix} \vec{a} \cdot \vec{c} & \vec{a} \cdot \vec{d} \\ \vec{b} \cdot \vec{c} & \vec{b} \cdot \vec{d} \end{vmatrix}$ <div style="border: 1px solid black; padding: 2px; display: inline-block;"> $\frac{dt}{ds} = \frac{1}{v}$ </div>	$T = \hat{v} = \frac{\vec{v}}{v}$ $\frac{dT}{dt} = \frac{(\vec{v} \cdot \vec{v})\vec{a} - (\vec{v} \cdot \vec{a})\vec{v}}{v^3} = \frac{\vec{v} \times (\vec{a} \times \vec{v})}{v^3} = \frac{(\vec{v} \times \vec{a}) \times \vec{v}}{v^3}$ $\left\ \frac{dT}{dt} \right\ = \frac{\sqrt{v^2 a^2 - (\vec{v} \cdot \vec{a})^2}}{v^2} = \frac{\ \vec{a} \times \vec{v}\ }{v^2}, \quad \frac{dT}{ds} = k \hat{N}$ $\hat{N} = \frac{T'}{\ T'\ } = \frac{(\vec{v} \times \vec{a}) \times \vec{v}}{\ \vec{v} \times \vec{a}\ v} = \hat{B} \times \hat{v}$ $\hat{B} = \frac{\vec{v} \times \vec{a}}{\ \vec{v} \times \vec{a}\ } = \widehat{\vec{v} \times \vec{a}} = \hat{v} \times \hat{N} \quad (\hat{B} \cdot \vec{v} = 0)$ $\frac{d\hat{B}}{dt} = \frac{\vec{v} \times \dot{\vec{a}}}{\ \vec{v} \times \vec{a}\ } - \left[\frac{\vec{v} \times \dot{\vec{a}}}{\ \vec{v} \times \vec{a}\ } \cdot \hat{B} \right] \hat{B}, \quad \tau = \hat{N} \cdot \frac{d\hat{B}}{ds}$	<div style="border: 1px solid black; padding: 5px;"> $\vec{a} = a_T \hat{T} + a_N \hat{N}$ $a_T = \vec{a} \cdot \hat{v} = \frac{dv}{dt}$ $a_N = \frac{\ \vec{a} \times \vec{v}\ }{v} = \ \vec{a} \times \hat{v}\$ $a^2 = a_T^2 + a_N^2 = \left\ \frac{d\vec{v}}{dt} \right\ ^2$ </div>
--	--	--

Frenet Trihedron

<p>Differentiable (in this book) : C^∞</p> <p>No singular pts. Order 0 (Regular) : $\vec{v}(t) \neq 0$</p> <p>• $\ \vec{v}(t)\ = c \rightarrow 1 \Rightarrow \int_s \ \vec{v}(t)\ dt = t = \Delta s$ $\rightarrow s : \vec{x}(t) = \vec{x}(s)$</p> <p>• $\frac{1}{2} \frac{d}{dt} (\vec{v} \cdot \vec{v}) = \vec{v} \cdot \vec{a} = 0$</p> <p>No singular pts. Order 1 : $\vec{a}(t) \neq 0$</p> <p>• Curvature, $k \neq 0$ (see right) • Vertex, $k' = 0$</p>	<p>$1 = \ \vec{t}\ = \ \vec{n}\ = \ \vec{b}\ , \quad 0 = \vec{t} \cdot \vec{n} = \vec{n} \cdot \vec{b} = \vec{b} \cdot \vec{t}$</p> <p>• $\vec{v}(s) = \vec{t}(s) \quad (t = n \times b)$</p> <p>• $\vec{a}(s) = \vec{t}'(s) = k(s) \vec{n}(s), \quad k(s) \geq 0$ (can be L or R-handed) (can be neg. if in \mathbb{R}^2)</p> <p>* $k(s) > 0$ for well defined curve with \hat{n}</p> <p>• $\vec{b} = \vec{t} \times \vec{n}, \quad \frac{d}{dt} (\vec{b} \cdot \vec{b}) = \vec{b} \cdot \vec{b}' = 0, \quad * \vec{b}'(s) = \tau(s) \vec{n}(s)$</p> <p>• $\vec{n} = \vec{b} \times \vec{t}, \quad * \vec{n}'(s) = -k \vec{t} - \tau \vec{b}, \quad * \text{t-n pl.} = \text{osculating pl.}$</p>
--	--

<p>• $t''(s) = k'n - k^2 t - k\tau b$</p> <p>• $\tau = \ b'\$</p> <p>• $n \Rightarrow k, \tau : \quad * \ n'\ ^2 = k^2 + \tau^2$</p>	<p>• $b''(s) = \tau'n - \tau kt - \tau^2 b$</p> <p>• $\tau = -\frac{(t \times t') \cdot t''}{k^2} = -\frac{t \cdot (t' \times t'')}{\ t'\ ^2}$</p> <p>• $k = \ t'\ = \frac{(b \times b') \cdot b''}{\tau^2} = \frac{b \cdot (b' \times b'')}{\ b'\ ^2}$</p> <p>• $\frac{(n \times n') \cdot n''}{\ n'\ ^2} = \frac{k' \tau - k \tau'}{k^2 + \tau^2} = \frac{\frac{d}{ds}(k/\tau)}{(k/\tau)^2 + 1} = \frac{d}{ds} \arctan(k/\tau)$</p>
--	--

Indicatrix [of Tangents] :

- $\vec{t}(\theta(s)) = (\cos \theta, \sin \theta) = (x'(s), y'(s))$
- $\vec{t}'(\theta) = \theta'(s) (-\sin \theta, \cos \theta) = \underline{k(s)} \vec{n}$
- $\theta(s) = \arctan(y'/x')$
- $\int_0^l k(s) ds = \theta(s) \Big|_0^l = 2\pi I_{\text{rot. index}}$

Local Canonical Form at $t = 0$:

- $(\hat{t}, \hat{n}, \hat{b}) = (\hat{x}, \hat{y}, \hat{z})$
- $\vec{r}(s) - \vec{r}(0) \approx (s - \frac{k^2 s^3}{6}, \frac{k}{2} s^2 + \frac{k' s^3}{6}, \frac{-k\tau}{6} s^3)$
- $\tau < 0 \Rightarrow \frac{dz}{ds} > 0$

Isoperimetric Inequality : $0 \leq l^2 - 4\pi A$

Four-Vertex Theorem : A simple closed curve has ≥ 4 vertices

Cauchy-Crofton Formula (measure of number of times lines intersect a curve) :

- Tangent line at $(\rho, \theta) : x \cos \theta + y \sin \theta = \rho$
- Curve $c : y = 0, x \in (-l/2, l/2), \quad C = \sum c_i$
- $\int \text{Lines that cross } C = \int_0^{2\pi} \int_0^{|\cos \theta| l/2} d\rho d\theta = 2l \Rightarrow \int_0^{2\pi} \int_0^\infty n_C d\rho d\theta = 2l$

2 Jacobian/Differential, $dF_{\alpha(0)} : \mathbb{R}^n \rightarrow \mathbb{R}^m$

- $\boxed{\alpha(0) = \beta(0)} \Rightarrow \underline{F(t=0)} = F \circ \alpha|_{t=0} = F \circ \beta|_{t=0}$

- $\boxed{\alpha'(0) = \beta'(0)} \Rightarrow \frac{\partial x}{\partial \alpha_i}|_{t=0} = \frac{\partial x}{\partial \beta_i}|_{t=0} \cdot \frac{d\beta_i/dt}{d\alpha_i/dt}|_{t=0} \Rightarrow \underline{dF_{\alpha(0)}(\alpha'(0)) = dF_{\beta(0)}(\beta'(0))}$

- * $F = (x_0, x_1, \dots, x_m) \Rightarrow \underline{dF_{\alpha(0)}(\alpha'(0))} \equiv \frac{d}{dt}(F \circ \alpha)|_{t=0} = \begin{bmatrix} \frac{\partial x_1}{\partial \alpha_0} & \frac{\partial x_1}{\partial \alpha_1} & \dots \\ \frac{\partial x_2}{\partial \alpha_0} & \frac{\partial x_2}{\partial \alpha_1} & \dots \\ \vdots & & \end{bmatrix}_{t=0} \begin{bmatrix} \frac{d\alpha_0}{dt} \\ \frac{d\alpha_1}{dt} \\ \vdots \end{bmatrix}_{t=0} = \boxed{J_{F(0)} \cdot \alpha'(0)}$

- $d(G \circ F)_p = dG_{F(p)} \circ dF_p$

3 Del

$$\nabla F = \left(\hat{r} \frac{\partial}{\partial r} + \hat{\theta} \frac{1}{r} \frac{\partial}{\partial \theta} + \hat{\phi} \frac{1}{r \sin \theta} \frac{\partial}{\partial \phi} \right) F$$

$$= \begin{bmatrix} \hat{r} \\ \hat{\theta} \\ \hat{\phi} \end{bmatrix} \cdot \begin{bmatrix} \frac{\partial}{\partial r} \\ \frac{1}{r} \frac{\partial}{\partial \theta} \\ \frac{1}{r \sin \theta} \frac{\partial}{\partial \phi} \end{bmatrix} F = \begin{bmatrix} \cos \phi \sin \theta \hat{x} + \sin \phi \sin \theta \hat{y} + \cos \theta \hat{z} \\ \cos \phi \cos \theta \hat{x} + \sin \phi \cos \theta \hat{y} - \sin \theta \hat{z} \\ -\sin \phi \hat{x} + \cos \phi \hat{y} \end{bmatrix} \cdot \begin{bmatrix} \frac{\partial}{\partial r} \\ \frac{1}{r} \frac{\partial}{\partial \theta} \\ \frac{1}{r \sin \theta} \frac{\partial}{\partial \phi} \end{bmatrix} F$$

$$\begin{bmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \\ \frac{\partial}{\partial z} \end{bmatrix} F = \begin{bmatrix} \cos \phi \sin \theta \frac{\partial}{\partial r} + \frac{\cos \phi \cos \theta}{r} \frac{\partial}{\partial \theta} - \frac{\sin \phi}{r \sin \theta} \frac{\partial}{\partial \phi} \\ \sin \phi \sin \theta \frac{\partial}{\partial r} + \frac{\sin \phi \cos \theta}{r} \frac{\partial}{\partial \theta} + \frac{\cos \phi}{r \sin \theta} \frac{\partial}{\partial \phi} \\ \cos \theta \frac{\partial}{\partial r} - \frac{\sin \theta}{r} \frac{\partial}{\partial \theta} \end{bmatrix} F = \begin{bmatrix} \frac{\partial r}{\partial x} \frac{\partial}{\partial r} + \frac{\partial \theta}{\partial x} \frac{\partial}{\partial \theta} + \frac{\partial \phi}{\partial x} \frac{\partial}{\partial \phi} \\ \frac{\partial r}{\partial y} \frac{\partial}{\partial r} + \frac{\partial \theta}{\partial y} \frac{\partial}{\partial \theta} + \frac{\partial \phi}{\partial y} \frac{\partial}{\partial \phi} \\ \frac{\partial r}{\partial z} \frac{\partial}{\partial r} + \frac{\partial \theta}{\partial z} \frac{\partial}{\partial \theta} + \frac{\partial \phi}{\partial z} \frac{\partial}{\partial \phi} \end{bmatrix} F = \begin{bmatrix} \hat{x} \\ \hat{y} \\ \hat{z} \end{bmatrix} \cdot \begin{bmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \\ \frac{\partial}{\partial z} \end{bmatrix} F$$

$$\nabla \cdot \vec{A} = \frac{1}{r} \frac{1}{r \sin \theta} \left\langle \frac{\partial}{\partial r}, \frac{\partial}{\partial \theta}, \frac{\partial}{\partial \phi} \right\rangle \cdot [r \cdot r \sin \theta] \left\langle A_r, \frac{1}{r} A_\theta, \frac{1}{r \sin \theta} A_\phi \right\rangle$$

$$\nabla \times \vec{A} = \frac{1}{r} \frac{1}{r \sin \theta} \begin{vmatrix} \hat{r} & r\hat{\theta} & r \sin \theta \hat{\phi} \\ \frac{\partial}{\partial r} & \frac{\partial}{\partial \theta} & \frac{\partial}{\partial \phi} \\ A_r & r A_\theta & r \sin \theta A_\phi \end{vmatrix} = \begin{vmatrix} \frac{\partial \vec{r}}{\partial r} & \frac{\partial \vec{r}}{\partial \theta} & \frac{\partial \vec{r}}{\partial \phi} \\ \frac{\partial}{\partial r} & \frac{\partial}{\partial \theta} & \frac{\partial}{\partial \phi} \\ A_r & r A_\theta & r \sin \theta A_\phi \end{vmatrix}$$

$$[\vec{A} \times (\vec{B} \times \vec{C})]_i = \vec{A} \cdot (B_i \vec{C}) - \vec{A} \cdot (\vec{B} C_i)$$

$$\begin{aligned} \vec{A} \times (\vec{B} \times \vec{C}) &= \boxed{(\vec{A} \cdot (\vec{B} \otimes \vec{C})^T)^T - (\vec{A} \cdot \vec{B} \otimes \vec{C})^T} = (\vec{A} \otimes \vec{B}) \cdot \vec{C} - (\vec{A} \cdot \vec{B} \otimes \vec{C})^T \\ &= (A^T (BC^T)^T)^T - (A^T BC^T)^T = (AB^T)^T C - (A^T BC^T)^T \end{aligned}$$