

# Notes Section R.2 – Integer Exponents

## Lesson Objectives

1. Zero and negative exponents
2. Product, Quotient, and Power Properties for Exponents

### A. ZERO and NEGATIVE exponents

Zero and negative exponents may be better understood by reviewing place value with base 10:

It turns out that **any** base (except zero) that has a zero power is equal to 1.

**Property:**  $a^0 = 1$  ( $a \neq 0$ )

Negative exponents do not make negative numbers! They cause a

**Property:**  $a^{-n} = \frac{1}{a^n}$  ( $a \neq 0$ )

and  $\frac{1}{a^{-n}} = a^n$

for negative exponents.

Cross the line, ~~\_\_\_\_\_~~ !

Powers of 10			
Power of 10	Standard Form	Fractional Form	Place Value
$10^4$	10,000	$\frac{10,000}{1}$	ten thousands
$10^3$	1,000	$\frac{1,000}{1}$	thousands
$10^2$	100	$\frac{100}{1}$	hundreds
$10^1$	10	$\frac{10}{1}$	tens
$10^0$	1	$\frac{1}{1}$	ones
$10^{-1}$	0.1	$\frac{1}{10}$	tenths
$10^{-2}$	0.01	$\frac{1}{100}$	hundredths
$10^{-3}$	0.001	$\frac{1}{1,000}$	thousandths
$10^{-4}$	0.0001	$\frac{1}{10,000}$	ten thousandths

- Negative Exponents and Fractions – “take the stairs” (reciprocal)

**Property:**  $\frac{a^{-n}}{b^{-m}} = \frac{b^m}{a^n}$

**Property:**  $\left(\frac{a}{b}\right)^{-n} = \frac{b^n}{a^n}$

**Simplifying Exponents Tip #1:** Final answer should have **no** exponents.

### B. Product, Quotient, and Power Rules for Exponents

- **Product Rule:**  $a^m \cdot a^n = a^{m+n}$

When multiplying powers w/**same base**, add exponents.

- **Quotient Rule:**  $\frac{a^m}{a^n} = a^{m-n}$

When dividing powers w/**same base**, subtract exponents.

## Notes Section R.2 – Integer Exponents

### Simplifying Exponents Tip #2:

Final answer should have **no** **variables** –  
you should see each variable only .

- **Power Rule:**  $(a^m)^n =$

When raising a power to a power, exponents.

- **Product to Power Rule:**  $(ab)^n =$

The exponent applies to factor in the parentheses.

- **Quotient to Power Rule:**  $\left(\frac{a}{b}\right)^n =$

The exponent applies to factor in numerator and denominator.

### Simplifying Exponents Tip #3:

Final answer should have **no** .

### Simplifying Exponents Tip #4:

A factor with no visible exponent has an **understood** (or implied) **exponent of** . To reduce errors, consider writing in an exponent of 1 in these situations.

### Simplifying Exponents Tip #5:

Simplify exponents that have a constant (number) base.  
“ !” For example, don’t leave  $2^5$  – change it to 32.

### Other Tips for Success:

Remember to always reduce (simplify) .  
Use a for any numerical (non-variable) part.  
Watch out with the negatives!  
To get better with exponents, you must **PRACTICE!**

- **EXAMPLE:** Evaluate.  $(-29)^0$  [R.2.21]

Any base (except zero) raised to the power of 0 equals 1.  $(-29)^0 =$  **Answer**

## Notes Section R.2 – Integer Exponents

- EXAMPLE:** Simplify the following expression.  $6^{-3}$  [R.2.27]

There are two common **errors** with this:

1. Multiplying **error**  $6^{-3} = -18$  (INCORRECT!)
2. Sign **error**  $6^{-3} = -6^3 = -216$  (INCORRECT!)

NOTE: A negative exponent does **not** make a negative number!  
A negative exponent means (flip it!).

$6^{-3}$	$= \frac{6^{-3}}{1}$	$=$	$=$
Write it as a	Negative exponent means “ ” (reciprocal). BASE (6) is the ; EXPONENT (– 3) changes sign.	Simplify – “ .” $6^3 = 216$	<b>Answer</b>

Note also that any time you are dealing with constants (no variables), you can verify the result using your calculator:

```
6^(-3)
.0046296296
Ans>Frac
1/216
```

```
6^-3
.0046296296
Ans>Frac
1/216
```

- EXAMPLE:** Use the quotient rule to simplify the expression.  $\frac{6^{-5}}{7^{-2}}$  [R.2.33]

$\left( \frac{6^{-5}}{7^{-2}} \right)$	$=$	$=$
Negative exponent means “ <b>take the stairs</b> ” (reciprocal). Both bases (6) and (7) will take the stairs – they will places.	Simplify – “do it.” $7^2 = 49$ $6^5 = 7776$	(reduce fraction, if necessary.)  <b>Answer</b>

Note also that any time you are dealing with constants (no variables), you can verify the result using your calculator:

```
6^-5 / 7^-2
.0063014403
Ans>Frac
49/7776
```

```
((6^(-5)))/(7^(-2))
.0063014403
Ans>Frac
49/7776
```

## Notes Section R.2 – Integer Exponents

- **EXAMPLE:** Use the product rule to simplify.  $8^0 \cdot 8^7 \cdot 8^9$  [R.2.39]

(Type exponential notation with positive exponents.)

There are two common **errors** with this:

1. Multiplying **error**  $8^0 \cdot 8^7 \cdot 8^9 = (8 \cdot 8 \cdot 8)^{0+7+9} = 512^{16}$  (INCORRECT!)
2. Multiplying **error**  $8^0 \cdot 8^7 \cdot 8^9 = 8^{(0 \cdot 7 \cdot 9)} = 8^0 = 1$  (INCORRECT!)

- **Product Rule:**  $a^m \cdot a^n = a^{m+n}$

When multiplying powers w/**same base**, **ADD** exponents.

$$8^0 \cdot 8^7 \cdot 8^9 = 8\text{-----} = 8\text{-----} \quad \textbf{Answer}$$

NOTE: because the solution must be in *exponential notation*, using the calculator isn't helpful.

- **EXAMPLE:** Multiply and simplify.  $3^5 \cdot 3^{-18}$  [R.2.35]

(Simplify your answer. Type exponential notation with positive exponents.)

$3^5 \cdot 3^{-18}$	$=$	$=$	$=$
<p>• <b>Product Rule:</b>  <math>a^m \cdot a^n = a^{m+n}</math>          When multiplying powers w/<b>same base</b>, <b>ADD</b> exponents.</p>	Can't have	exponents!	<p><b>Answer</b>          solution must be in <i>exponential notation</i>          Leave answer with exponent – don't "do it."</p>

- **EXAMPLE:** Use the product rule to simplify.  $5x^{-4} \cdot 3x^8 \cdot x^5$  [R.2.37]

(Type exponential notation with positive exponents.)

$5x^{-4} \cdot 3x^8 \cdot x^5$	$= ( \quad ) \cdot ( \quad )$	$=$
_____ factors to multiply constants separately from variables.	<p>Simplify.</p> <ul style="list-style-type: none"> <li><b>Product Rule:</b>  <math display="block">a^m \cdot a^n = a^{m+n}</math> </li> </ul> <p>When multiplying powers w/<b>same base</b>, <b>ADD</b> exponents.</p>	<p><b>Answer</b></p> <p>(No negative exponents.)</p>


## Notes Section R.2 – Integer Exponents

- EXAMPLE:** Use the quotient rule to simplify the expression.

[R.2-23]

Use positive exponents to write the answer.

$$\frac{4^{-4}}{4^8}$$

$\frac{4^{-4}}{4^8}$	$= 4$	$= 4 =$ 	$=$
<p>Same base (4).</p> <ul style="list-style-type: none"> <li><b>Quotient Rule:</b>  <math>\frac{a^m}{a^n} = a^{m-n}</math></li> </ul> <p>When dividing powers w/<b>same base</b>, <b>SUBTRACT</b> exponents.</p>	<p>Simplify.</p> $-4 - (8) = -12$	<p>Can't have negative exponents!</p> <p>Write as fraction.</p> <p>"Take the stairs" (reciprocal).</p>	<p><b>Solution</b></p> <p>Leave answer with exponent – don't "do it."</p>

### EASIER WAY? – RESET!


- EXAMPLE:** Use the ~~quotient rule~~ to simplify the expression.

[R.2-23]

Use positive exponents to write the answer.

$$\frac{4^{-4}}{4^8}$$

Rather than use the quotient rule, focus on the **negative exponent** in the \_\_\_\_\_

$\frac{4^{-4}}{4^8}$ 	$=$	$= \frac{1}{\quad}$
<p>Because of the negative exponent, "take the stairs" (reciprocal).</p> <p>Connect existing denominator <math>4^8</math> with new piece <math>4^4</math> using _____</p>	<ul style="list-style-type: none"> <li><b>Product Rule:</b>  <math>a^m \cdot a^n = a^{m+n}</math></li> </ul> <p>When multiplying powers w/<b>same base</b>, <b>ADD</b> exponents.</p>	<p><b>Answer</b></p> <p>Leave answer with exponent – don't "do it."</p>

## Notes Section R.2 – Integer Exponents

- Another way to do Division: “\_\_\_\_\_”

- EXAMPLE:** Simplify the expression. Write the answer with only positive exponents.

All variables are nonzero.

[R.2.69]

$$-\frac{24a^3b^{-2}}{18ab^{-5}}$$

To simplify this expression, work in “\_\_\_\_\_” – constants (coefficients), variable  $a$ , variable  $b$ . Then, \_\_\_\_\_(multiply) all the results together.

Constants (coefficients).	Variable $a$ .	Variable $b$ .		Merge.
$-\frac{24}{18} =$	$\frac{a^3}{a}$	1. $\left( \frac{b^{-2}}{b^{-5}} \right)$	2. $=$	Merge together the results from constants (coefficients) and variables.
Simplify the fraction of coefficients, if possible.	<b>“Face-off!”</b> 1. Do both have positive exponents?  2. Who has more, top or bottom?  3. By how much?  4. Simplifies to $a^2$ on TOP.  $\frac{a^3}{a^1} = \frac{a^2}{1} =$	<b>“Face-off!”</b> 1. Do both have positive exponents?  “take the stairs” (reciprocal) 2. Who has more, top or bottom?  3. By how much?  4. Simplifies to $b^3$ on TOP.  $\frac{b^5}{b^2} = =$		constants (coefficients): $= -\frac{4}{3}$ Variable $a$ : $= a^2$ Variable $b$ : $= b^3$  <b>Merged</b>  $=$  or  $=$  <b>Answer</b>

## Notes Section R.2 – Integer Exponents

- EXAMPLE:** Use the rules of exponents to simplify the expression. [R.2.77]

$$\left(\frac{3x^6}{5y^{-3}}\right)^2$$

(Type exponential notation with positive exponents.)

$\left(\frac{3x^6}{5y^{-3}}\right)^2$	$= \left(\frac{3x^6}{5y^{-3}}\right)^2$	$=$	$= \frac{(3^1)(x^6)(y^3)}{(5^1)}$	$=$
<p>Try to simplify INSIDE parentheses first.</p> <ul style="list-style-type: none"> <li>Fraction <math>\frac{3}{5}</math> is already simplified.</li> <li><math>x^6</math> is fine ... already has positive exponent</li> </ul>	<ul style="list-style-type: none"> <li>needs fixing ... because of the negative exponent, “take the stairs” (reciprocal).</li> </ul>	<ul style="list-style-type: none"> <li>Coefficients always have exponent of understood ____.</li> <li><b>Quotient to Power Rule</b>  <math>\left(\frac{a}{b}\right)^n = \frac{a^n}{b^n}</math>                      The exponent ( ) applies to numerator and denominator.</li> </ul>	<p>The POSITIVE exponent, 2, applies to ALL factors.</p> <ul style="list-style-type: none"> <li><b>Power Rule:</b>  <math>(a^m)^n = a^{m \cdot n}</math></li> </ul> <p>When raising a power to a power, <b>MULTIPLY</b> exponents.</p> <p>Now simplify:  <math>(3^1)^2 = 3^2 =</math>  <math>(x^6)^2 =</math>  <math>(y^3)^2 =</math>  <math>(5^1)^2 = 5^2 =</math></p>	<p><b>Answer</b></p> <p>Carefully merge together all the separate calculations.</p> <p>Be careful who goes in the numerator (TOP) and who goes in the denominator (BOTTOM).</p>

- EXAMPLE:** Simplify and write with positive exponents. [R.2.73]

$$(5x^{-3}y^3)^{-3}$$

$(5x^{-3}y^3)^{-3}$ $= (5^1x^{-3}y^3)^{-3}$	$= (5^1)(x^{-3})(y^3)$ $=$	$= \frac{(5)^{-3}x^9y^{-9}}{1}$	$=$
<ul style="list-style-type: none"> <li>Coefficients always have exponent of understood <b>1</b>.</li> <li><b>Product to Power Rule:</b>  <math>(ab)^n = a^n b^n</math>                      The exponent (– 3) applies to each factor in the parentheses.</li> </ul>	<ul style="list-style-type: none"> <li><b>Power Rule:</b>  <math>(a^m)^n = a^{m \cdot n}</math>                      When raising a power to a power, <b>MULTIPLY</b> exponents.                      Now, write as fraction.</li> </ul>	<p><math>(5)^{-3}</math> and <math>y^{-9}</math> need fixing ... because of the negative exponent, “take the stairs” (reciprocal).</p>	<p>Finally, just need to simplify <math>5^3 = 125</math> in denominator.</p> <p><b>Answer</b></p>

Sources Used:

- MyLab Math for *College Algebra with Modeling and Visualization*, 6<sup>th</sup> Edition, Rockswold, Pearson Education Inc.
- Powers of 10 chart, [https://www.eduplace.com/math/mw/background/6/01/te\\_6\\_01\\_overview.html](https://www.eduplace.com/math/mw/background/6/01/te_6_01_overview.html)