

DD2434 - Machine Learning, Advanced Course  
Assignment 1B

Tristan Perrot  
tristanp@kth.se

November 2023



## Contents

|          |                              |          |
|----------|------------------------------|----------|
| <b>1</b> | <b>CAVI for Earth quakes</b> | <b>3</b> |
| 1.1      | Question 1.1 . . . . .       | 3        |
| 1.2      | Question 1.2 . . . . .       | 3        |
| 1.3      | Question 1.3 . . . . .       | 4        |
| <b>A</b> | <b>Appendix</b>              | <b>6</b> |
| A.1      | Question 1.2 . . . . .       | 6        |

# 1 CAVI for Earth quakes

## 1.1 Question 1.1




Figure 1: Directed Graphical Model for the Earthquake problem

## 1.2 Question 1.2

Let us take the Alternative 1 in 2D. Here, we know these distributions:

- $p(Z_n|\pi) = \text{Categorical}(\pi)$
- $p(S_n|Z_n = k, \lambda_k) = \text{Poisson}(\lambda_k)$
- $p(X_n|Z_n = k, \mu_k, \tau) = \text{Normal}(\mu_k, \tau \cdot I)$
- $p(\mu_k|\nu, \rho) = \text{Normal}(\nu, \rho \cdot I)$
- $p(\lambda_k|\alpha, \beta) = \text{Gamma}(\alpha, \beta)$

Where,  $\rho$  and  $\tau$  define precision and not standard variation. Then we have:

$$\begin{aligned}
 \log p(X, S, Z, \lambda, \mu|\pi, \tau, \alpha, \beta, \nu, \rho) &= \log p(X|S, Z, \lambda, \mu, \pi, \tau, \alpha, \beta, \nu, \rho) \\
 &\quad + \log p(S, Z, \lambda, \mu|\pi, \alpha, \beta, \nu, \rho) \\
 &= \log p(X|Z, \mu, \tau) + \log p(S|Z, \lambda, \mu, \pi, \alpha, \beta, \nu, \rho) \\
 &\quad + \log p(Z, \lambda, \mu|\pi, \alpha, \beta, \nu, \rho) \\
 &= \log p(X|Z, \mu, \tau) + \log p(S|Z, \lambda) + \log p(Z|\pi) \\
 &\quad + \log p(\lambda, \mu|\alpha, \beta, \nu, \rho) \\
 \log p(X, S, Z, \lambda, \mu|\pi, \tau, \alpha, \beta, \nu, \rho) &= \log p(X|Z, \mu, \tau) + \log p(S|Z, \lambda) + \log p(Z|\pi) \\
 &\quad + \log p(\mu|\nu, \rho) + \log p(\lambda|\alpha, \beta)
 \end{aligned} \tag{1}$$

Where:

$$\begin{aligned}
 \log p(X|Z, \mu, \tau) &= \sum_{n=1}^N \sum_{k=1}^K \log p(X_n|Z_n = k, \mu_k, \tau) \\
 \log p(S|Z, \lambda) &= \sum_{n=1}^N \sum_{k=1}^K \log p(S_n|Z_n = k, \lambda_k) \\
 \log p(Z|\pi) &= \sum_{n=1}^N \log p(Z_n|\pi) \\
 \log p(\mu|\nu, \rho) &= \sum_{k=1}^K \log p(\mu_k|\nu, \rho) \\
 \log p(\lambda|\alpha, \beta) &= \sum_{k=1}^K \log p(\lambda_k|\alpha, \beta)
 \end{aligned} \tag{2}$$

### 1.3 Question 1.3

Here, the mean field approximation is not an approximation but an equality because  $Z, \mu, \lambda$  are independent. Therefore we have:

$$\begin{aligned}
 \log q^*(Z_n) &\stackrel{\pm}{=} \mathbb{E}_{\mu, \lambda} [\log p(X_n, S_n, Z_n, \lambda, \mu|\pi, \tau, \alpha, \beta, \nu, \rho)] \\
 &\stackrel{\pm}{=} \mathbb{E}_{\mu, \lambda} [\log p(X_n|Z_n, \mu, \tau) + \log p(S_n|Z_n, \lambda) + \log p(Z_n|\pi)] \\
 &= \mathbb{E}_{\mu} \left[ \sum_{k=1}^K \mathbb{1}_{\{Z_n=k\}} \left( \log \left( \frac{\tau}{2\pi} \right) - \frac{\tau}{2} ((x_n - \mu_k)^T (x_n - \mu_k)) \right) \right] \\
 &\quad + \mathbb{E}_{\lambda} \left[ \sum_{k=1}^K \mathbb{1}_{\{Z_n=k\}} (\log(\pi_k) - \lambda_k + S_n \log(\lambda_k) - \log(S_n!)) \right] \\
 &\stackrel{\pm}{=} \sum_{k=1}^K \mathbb{1}_{\{Z_n=k\}} \left( \log \left( \frac{\tau}{2\pi} \right) - \frac{\tau}{2} \mathbb{E}_{\mu} [(x_n - \mu_k)^T (x_n - \mu_k)] \right. \\
 &\quad \left. + \log(\pi_k) + \mathbb{E}_{\lambda} [-\lambda_k + S_n \log(\lambda_k)] - \log(S_n!) \right)
 \end{aligned} \tag{3}$$

Now, if we take the entire expression that is multiplied by  $\mathbb{1}_{\{Z_n=k\}}$  and we call it  $u_{n,k}$ , we have:

$$q^*(Z_n) \propto \prod_{k=1}^K u_{n,k}^{\mathbb{1}_{\{Z_n=k\}}} \tag{4}$$

And if we normalize by taking  $r_{n,k} = \frac{u_{n,k}}{\sum_{i=1}^K u_{n,i}}$  we get:

$$q^*(Z_n) = \prod_{k=1}^K r_{n,k}^{\mathbb{1}_{\{Z_n=k\}}} \tag{5}$$

Wich means that  $q^*(Z_n)$  is a categorical distribution with parameters  $r_{n,k}$ . There for we have the expectation of  $Z_n$  easily because  $\mathbb{E}[z_{n,k}] = r_{n,k}$  where  $z_{n,k} = \mathbb{1}_{\{S_n=k\}}$ . Note that  $r_{n,k}$  depends of

the expected value of  $\mu_k$ ,  $\mu_k^2$ ,  $\lambda_k$  and  $\log \lambda_k$ . We will be able to compute these expected values by finding  $q^*(\mu_k)$  and  $q^*(\lambda_k)$ .

Let us compute  $q^*(\mu_k)$ :

$$\begin{aligned}
 \log q^*(\mu_k) &\stackrel{\pm}{=} \mathbb{E}_{Z,\lambda}[\log p(X, S, Z = k, \lambda_k, \mu_k | \pi, \tau, \alpha, \beta, \nu, \rho)] \\
 &\stackrel{\pm}{=} \mathbb{E}_{Z,\lambda}[\log p(X|Z = k, \mu_k, \tau) + \log p(\mu_k | \nu, \rho)] \\
 &= \mathbb{E}_{Z,\lambda} \left[ \sum_{n=1}^N \mathbb{1}_{\{Z_n=k\}} \left( \log \left( \frac{\tau}{2\pi} \right) - \frac{\tau}{2} ((x_n - \mu_k)^T (x_n - \mu_k)) \right) \right] \\
 &\quad + \log \left( \frac{\rho}{2\pi} \right) - \frac{\rho}{2} ((\mu_k - \nu)^T (\mu_k - \nu)) \\
 &\stackrel{\pm}{=} \sum_{n=1}^N r_{n,k} \left( \log \left( \frac{\tau}{2\pi} \right) - \frac{\tau}{2} ((x_n - \mu_k)^T (x_n - \mu_k)) \right) - \frac{\rho}{2} ((\mu_k - \nu)^T (\mu_k - \nu))
 \end{aligned} \tag{6}$$

$$\begin{aligned}
 \log q^*(\lambda_k) &\stackrel{\pm}{=} \mathbb{E}_{Z,\mu}[\log p(X, S, Z = k, \lambda_k, \mu_k | \pi, \tau, \alpha, \beta, \nu, \rho)] \\
 &\stackrel{\pm}{=} \mathbb{E}_{Z,\mu}[\log p(S|Z = k, \lambda_k) + \log p(\lambda_k | \alpha, \beta)] \\
 &= \mathbb{E}_Z \left[ \sum_{n=1}^N \mathbb{1}_{\{Z_n=k\}} (-\lambda_k + S_n \log(\lambda_k) - \log(S_n!)) \right] \\
 &\quad + \log \left( \frac{\beta^\alpha}{\Gamma(\alpha)} \right) + (\alpha - 1) \log(\lambda_k) - \beta \lambda_k \\
 &\stackrel{\pm}{=} \sum_{n=1}^N r_{n,k} (-\lambda_k + S_n \log(\lambda_k)) + (\alpha - 1) \log(\lambda_k) - \beta \lambda_k \\
 &= \left( \alpha + \sum_{n=1}^N S_n r_{n,k} - 1 \right) \log(\lambda_k) - \left( \beta + \sum_{n=1}^N r_{n,k} \right) \lambda_k
 \end{aligned} \tag{7}$$

Therefore, we have  $q^*(\lambda_k) = \text{Gamma} \left( \alpha + \sum_{n=1}^N S_n r_{n,k}, \beta + \sum_{n=1}^N r_{n,k} \right)$ . And we can compute the expected value of  $\lambda_k$  and  $\log \lambda_k$  easily.

$$\begin{aligned}
 \mathbb{E}[\lambda_k] &= \frac{\alpha + \sum_{n=1}^N S_n r_{n,k}}{\beta + \sum_{n=1}^N r_{n,k}} \\
 \mathbb{E}[\log \lambda_k] &= \psi \left( \alpha + \sum_{n=1}^N S_n r_{n,k} \right) - \log \left( \beta + \sum_{n=1}^N r_{n,k} \right)
 \end{aligned} \tag{8}$$

## A Appendix

### A.1 Question 1.2

```
import matplotlib.pyplot as plt
import numpy as np
from scipy.stats import gamma, norm
from scipy.special import psi
np.random.seed(14)

def generate_data(mu, tau, N):
    # Insert your code here
    D = np.random.normal(mu, np.sqrt(1/tau), N)

    return D

MU = 1
TAU = 0.5

dataset_1 = generate_data(MU, TAU, 10)
dataset_2 = generate_data(MU, TAU, 100)
dataset_3 = generate_data(MU, TAU, 1000)

# Visualize the datasets via histograms
# Insert your code here
fig, axs = plt.subplots(1, 3, figsize=(12, 4))
axs[0].hist(dataset_1, bins=20)
axs[1].hist(dataset_2, bins=20)
axs[2].hist(dataset_3, bins=20)
plt.tight_layout()
plt.savefig('12_data.png')
plt.show()
```