DD2434 - Machine Learning, Advanced Course Assignment 1B

Tristan Perrot tristanp@kth.se

November 2023



Contents

1	CAVI for Earth quakes	3
	1.1 Question 1.1	3
	1.2 Question 1.2	3
A	Appendix	5
	A.1 Question 1.2	5

1 CAVI for Earth quakes

1.1 Question 1.1




Figure 1: Directed Graphical Model for the Earthquake problem

1.2 Question 1.2

Here, we know these distributions:

- $p(Z_n|\pi) = Categorical(\pi)$
- $p(S_n|Z_n = k, \lambda_k) = Poisson(\lambda_k)$
- $p(X_n|Z_n = k, \mu_k, \tau) = Normal(\mu_k, \tau \cdot I)$
- $p(\mu_k|\nu,\rho) = Normal(\nu,\rho\cdot I)$
- $p(\lambda_k | \alpha, \beta) = Gamma(\alpha, \beta)$

Where, ρ and τ define precision and not standard variation. Then we have:

$$\log p(X, S, Z, \lambda, \mu | \pi, \tau, \alpha, \beta, \nu, \rho) = \log p(X | S, Z, \lambda, \mu, \pi, \tau, \alpha, \beta, \nu, \rho)$$

$$+ \log p(S, Z, \lambda, \mu | \pi, \alpha, \beta, \nu, \rho)$$

$$= \log p(X | Z, \mu, \tau) + \log p(S | Z, \lambda, \mu, \pi, \alpha, \beta, \nu, \rho)$$

$$+ \log p(Z, \lambda, \mu | \pi\alpha, \beta, \nu, \rho)$$

$$= \log p(X | Z, \mu, \tau) + \log p(S | Z, \lambda) + \log p(Z | \pi)$$

$$+ \log p(\lambda, \mu | \alpha, \beta, \nu, \rho)$$

$$\log p(X, S, Z, \lambda, \mu | \pi, \tau, \alpha, \beta, \nu, \rho) = \log p(X | Z, \mu, \tau) + \log p(S | Z, \lambda) + \log p(Z | \pi)$$

$$+ \log p(\mu | \nu, \rho) + \log p(\lambda | \alpha, \beta)$$

$$(1)$$

Where:

$$\log p(X|Z, \mu, \tau) = \sum_{n=1}^{N} \sum_{k=1}^{K} \log p(X_n | Z_n = k, \mu_k, \tau)$$

$$\log p(S|Z, \lambda) = \sum_{n=1}^{N} \sum_{k=1}^{K} \log p(S_n | Z_n = k, \lambda_k)$$

$$\log p(Z|\pi) = \sum_{n=1}^{N} \log p(Z_n | \pi)$$

$$\log p(\mu | \nu, \rho) = \sum_{k=1}^{K} \log p(\mu_k | \nu, \rho)$$

$$\log p(\lambda | \alpha, \beta) = \sum_{k=1}^{K} \log p(\lambda_k | \alpha, \beta)$$
(2)

A Appendix

A.1 Question 1.2

```
import matplotlib.pyplot as plt
import numpy as np
from scipy.stats import gamma, norm
from scipy.special import psi
np.random.seed(14)
def generate_data(mu, tau, N):
    # Insert your code here
    D = np.random.normal(mu, np.sqrt(1/tau), N)
    return D
MU = 1
TAU = 0.5
dataset_1 = generate_data(MU, TAU, 10)
dataset_2 = generate_data(MU, TAU, 100)
dataset_3 = generate_data(MU, TAU, 1000)
# Visulaize the datasets via histograms
# Insert your code here
fig, axs = plt.subplots(1, 3, figsize=(12, 4))
axs[0].hist(dataset_1, bins=20)
axs[1].hist(dataset_2, bins=20)
axs[2].hist(dataset_3, bins=20)
plt.tight_layout()
plt.savefig('12_data.png')
plt.show()
```