Chapter 2: Analytic Functions

Taotao Wang, Associate Professor

College of Electronics and Information Engineering, Shenzhen University

November 1, 2026

Outline

1. Concept of Analytic Functions

2. Necessary and Sufficient Conditions for Analyticity

3. Elementary Functions

Outline

1. Concept of Analytic Functions

2. Necessary and Sufficient Conditions for Analyticity

3. Elementary Functions

1 Derivatives of Complex Functions

1) Definition of the Derivative

Definition

Let $w = f(z)$ be defined in a domain D, and let $z_0 \in D$ and $z_0 + \Delta z \in D$. If the limit

$$\lim_{\Delta z \to 0} \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z}$$

exists (and is finite), then $f(z)$ is said to be differentiable (可导) at z_0.

Definition (Cont.)

This limit is called the derivative (導数) *of* $f(z)$ *at* z_0 *and is denoted by*

$$f'(z_0) = \left.\frac{\mathrm{d}w}{\mathrm{d}z}\right|_{z=z_0} = \lim_{\Delta z \to 0} \frac{f(z_0+\Delta z) - f(z_0)}{\Delta z}. \tag{2.1.1}$$

Equivalently, for every $\varepsilon > 0$ *there is a* $\delta(\varepsilon) > 0$ *such that* $0 < |\Delta z| < \delta$ *implies*

$$\left| \frac{f(z_0+\Delta z) - f(z_0)}{\Delta z} - f'(z_0) \right| < \varepsilon.$$

As for limits in Chapter 1, $\Delta z \to 0$ *may happen along any path.*

Example 1.1

Determine whether the following functions are differentiable:

1. $f(z) = z^2$;
2. $f(z) = 2x + yi$.

Example 1.1

Determine whether the following functions are differentiable:

1. $f(z) = z^2$;
2. $f(z) = 2x + yi$.

Solution.

1. For every z in the complex plane,

$$\lim_{\Delta z \to 0} \frac{f(z+\Delta z) - f(z)}{\Delta z} = \lim_{\Delta z \to 0} \frac{(z+\Delta z)^2 - z^2}{\Delta z}$$
$$= \lim_{\Delta z \to 0} (2z + \Delta z) = 2z.$$

Hence $f(z) = z^2$ is differentiable everywhere, and $f'(z) = 2z$.

Solution (Cont.)

2 With $\Delta z = \Delta x + i\Delta y$,

$$\frac{f(z+\Delta z) - f(z)}{\Delta z} = \frac{2(x+\Delta x) + (y+\Delta y)i - (2x+yi)}{\Delta x + i\Delta y}$$
$$= \frac{2\Delta x + i\Delta y}{\Delta x + i\Delta y}.$$

If $z+\Delta z \to z$ along a line parallel to the real axis (Figure 2.1), then $\Delta y = 0$ and the quotient equals $\dfrac{2\Delta x}{\Delta x} = 2$.
If $z+\Delta z \to z$ along a line parallel to the imaginary axis, then $\Delta x = 0$ and the quotient equals $\dfrac{i\Delta y}{i\Delta y} = 1$.
The two limits differ, so $f(z) = 2x+yi$ is differentiable nowhere.

Solution (Cont.)




Figure: 2.1

2) Differentiability and Continuity
Suppose $f(z)$ is differentiable at z_0. Let

$$\rho(\Delta z) = \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z} - f'(z_0). \tag{2.1.2}$$

By the definition of the derivative,

$$\lim_{\Delta z \to 0} \rho(\Delta z) = 0.$$

From (2.1.2),

$$f(z_0 + \Delta z) = f(z_0) + f'(z_0)\Delta z + \rho(\Delta z)\Delta z, \qquad (2.1.3)$$

and therefore

$$\lim_{\Delta z \to 0} f(z_0 + \Delta z) = f(z_0).$$

Thus, differentiability at z_0 implies continuity at z_0.
The converse is false: $f(z) = 2x + yi$ is continuous everywhere but differentiable nowhere.

3) Rules of Differentiation
Complex derivatives obey the same rules as real derivatives:

a) $(c)' = 0$, where c is a complex constant;
b) $(z^n)' = nz^{n-1}$, where n is a positive integer;
c) $[f(z) \pm g(z)]' = f'(z) \pm g'(z)$;
d) $[f(z)g(z)]' = f'(z)g(z) + f(z)g'(z)$;
e) $\left[\frac{f(z)}{g(z)}\right]' = \frac{f'(z)g(z) - f(z)g'(z)}{g^2(z)}$, where $g(z) \neq 0$;
f) $\{f[g(z)]\}' = f'(w)g'(z)$, where $w = g(z)$ (chain rule);
g) $f'(z) = \frac{1}{\varphi'(w)}$, where $w = f(z)$ and $z = \varphi(w)$ are mutually inverse single-valued functions and $\varphi'(w) \neq 0$.

4) Differential

As in real analysis, suppose $w = f(z)$ is differentiable at z_0. By (2.1.3),

$$\Delta w = f(z_0 + \Delta z) - f(z_0) = f'(z_0)\Delta z + \rho(\Delta z)\Delta z,$$

where $\lim_{\Delta z \to 0} \rho(\Delta z) = 0$. Hence $\rho(\Delta z)\Delta z$ is an infinitesimal of higher order than Δz, and $f'(z_0)\Delta z$ is the linear principal part of Δw.

The linear part $f'(z_0)\Delta z$ is called the differential (微分) of $w = f(z)$ at z_0:

$$\mathrm{d}w = f'(z_0)\Delta z. \tag{2.1.4}$$

For $f(z) = z$ we have $\mathrm{d}z = \Delta z$, so (2.1.4) becomes $\mathrm{d}w = f'(z_0)\,\mathrm{d}z$, that is,

$$f'(z_0) = \left.\frac{\mathrm{d}w}{\mathrm{d}z}\right|_{z=z_0}. \tag{2.1.5}$$

Thus, a function has a differential at z_0 if and only if it has a derivative at z_0. If $f(z)$ is differentiable at every point of a domain D, it is said to be differentiable in D.

2 Analytic Functions

Definition

If $f(z)$ is differentiable at z_0 and at every point of some neighborhood of z_0, then $f(z)$ is said to be analytic (解析) *at z_0.*
If $f(z)$ is analytic at every point of a domain D, then $f(z)$ is said to be analytic in D, or an analytic function in D.
If $f(z)$ is not analytic at z_0, then z_0 is called a singular point (奇点) *of $f(z)$.*

Analyticity is a neighborhood property: differentiability at a single point is not enough.

Example 1.2

Investigate the analyticity of the following functions:

1. $f(z) = z^2$;
2. $f(z) = 2x + yi$;
3. $f(z) = |z|^2$;
4. $f(z) = 1/z$.

Example 1.2

Investigate the analyticity of the following functions:

1. $f(z) = z^2$;
2. $f(z) = 2x + yi$;
3. $f(z) = |z|^2$;
4. $f(z) = 1/z$.

Solution.

(1)(2) By Example 1.1, $f(z) = z^2$ is analytic in the whole plane, while $f(z) = 2x + yi$ is analytic nowhere.

(4) $f(z) = 1/z$ is differentiable for $z \neq 0$ with $f'(z) = -1/z^2$, so it is analytic in $\mathbb{C} \setminus \{0\}$, and $z = 0$ is a singular point.

Solution (Cont.)

(3) For $f(z) = |z|^2 = z\bar{z}$,

$$\frac{f(z_0+\Delta z)-f(z_0)}{\Delta z} = \frac{(z_0+\Delta z)(\overline{z_0}+\overline{\Delta z})-z_0\overline{z_0}}{\Delta z}$$
$$= \overline{z_0}+\overline{\Delta z}+z_0\frac{\overline{\Delta z}}{\Delta z}.$$

If $z_0 = 0$, the quotient equals $\overline{\Delta z} \to 0$, so $f'(0) = 0$.

Solution (Cont.)

If $z_0 \neq 0$, let $z_0 + \Delta z \to z_0$ along the line $y - y_0 = k(x - x_0)$. Then

$$\frac{\overline{\Delta z}}{\Delta z} = \frac{\Delta x - i\Delta y}{\Delta x + i\Delta y} = \frac{1 - ik}{1 + ik},$$

which depends on k, so the limit does not exist.
Therefore, $f(z) = |z|^2$ is differentiable only at $z = 0$ and is analytic nowhere.

Theorem 1.1

1. *If $f(z)$ and $g(z)$ are analytic in a domain D, then $f \pm g$, fg, and f/g (where $g(z) \neq 0$) are analytic in D.*
2. *Suppose $h = g(z)$ is analytic in a domain D of the z-plane, $w = f(h)$ is analytic in a domain G of the h-plane, and $g(z) \in G$ for every $z \in D$. Then the composite function $w = f[g(z)]$ is analytic in D.*

Consequently, every polynomial is analytic in the whole plane, and every rational function $P(z)/Q(z)$ is analytic except at the zeros of $Q(z)$.

Outline

1. Concept of Analytic Functions

2. Necessary and Sufficient Conditions for Analyticity

3. Elementary Functions

1 Main Theorem

Theorem 2.1

Let $f(z) = u(x, y) + iv(x, y)$ be defined in a domain D. Then $f(z)$ is differentiable at a point $z = x + iy$ of D if and only if $u(x, y)$ and $v(x, y)$ are differentiable at (x, y) and satisfy the Cauchy–Riemann equations (柯西–黎曼方程)

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \qquad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

at this point.

Proof.

1. Necessity. Suppose $f(z)$ is differentiable at $z = x + iy$. By (2.1.3), for small $|\Delta z| = |\Delta x + i\Delta y| > 0$,

$$f(z+\Delta z) - f(z) = f'(z)\Delta z + \rho(\Delta z)\Delta z,$$

where $\lim_{\Delta z \to 0} \rho(\Delta z) = 0$. Write

$$f(z+\Delta z)-f(z) = \Delta u + i\Delta v, \quad f'(z) = a+ib, \quad \rho(\Delta z) = \rho_1 + i\rho_2.$$

Proof (Cont.)

Then

$$\begin{aligned}\Delta u + i\Delta v &= (a+ib)(\Delta x + i\Delta y) + (\rho_1 + i\rho_2)(\Delta x + i\Delta y) \\ &= (a\Delta x - b\Delta y + \rho_1\Delta x - \rho_2\Delta y) \\ &\quad + i(b\Delta x + a\Delta y + \rho_2\Delta x + \rho_1\Delta y).\end{aligned}$$

Comparing real and imaginary parts,

$$\begin{aligned}\Delta u &= a\Delta x - b\Delta y + \rho_1\Delta x - \rho_2\Delta y, \\ \Delta v &= b\Delta x + a\Delta y + \rho_2\Delta x + \rho_1\Delta y.\end{aligned}$$

Proof (Cont.)

Since $\rho_1, \rho_2 \to 0$ as $(\Delta x, \Delta y) \to (0, 0)$, u and v are differentiable at (x, y), with

$$\frac{\partial u}{\partial x} = a, \quad \frac{\partial u}{\partial y} = -b, \quad \frac{\partial v}{\partial x} = b, \quad \frac{\partial v}{\partial y} = a.$$

Hence

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \qquad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}.$$

Technical Note: Equivalent Forms of the Remainder

Let

$$r = \sqrt{(\Delta x)^2 + (\Delta y)^2}, \qquad R(\Delta x, \Delta y) = o(r).$$

Then the remainder（余项）can be written as

$$R = \varepsilon_1 \Delta x + \varepsilon_2 \Delta y, \qquad \varepsilon_1, \varepsilon_2 \to 0.$$

For $r > 0$, choose the following values, and set both to 0 at $r = 0$:

$$\varepsilon_1 = \frac{R\Delta x}{r^2}, \qquad \varepsilon_2 = \frac{R\Delta y}{r^2}.$$

Then $\varepsilon_1 \Delta x + \varepsilon_2 \Delta y = R$, and

$$|\varepsilon_1|, |\varepsilon_2| \leq \frac{|R|}{r} \to 0.$$

Proof (Cont.)

2. Sufficiency. Suppose u and v are differentiable at (x, y) and satisfy the Cauchy–Riemann equations. Write

$$f(z + \Delta z) - f(z) = \Delta u + i\Delta v.$$

Proof (Cont.)

By the preceding technical note, differentiability of u and v gives

$$\Delta u = \frac{\partial u}{\partial x}\Delta x + \frac{\partial u}{\partial y}\Delta y + \varepsilon_1 \Delta x + \varepsilon_2 \Delta y,$$
$$\Delta v = \frac{\partial v}{\partial x}\Delta x + \frac{\partial v}{\partial y}\Delta y + \varepsilon_3 \Delta x + \varepsilon_4 \Delta y,$$

where $\varepsilon_k \to 0$ $(k = 1, 2, 3, 4)$ as $(\Delta x, \Delta y) \to (0, 0)$. Therefore,

$$f(z + \Delta z) - f(z) = \left(\frac{\partial u}{\partial x} + i\frac{\partial v}{\partial x}\right)\Delta x + \left(\frac{\partial u}{\partial y} + i\frac{\partial v}{\partial y}\right)\Delta y + (\varepsilon_1 + i\varepsilon_3)\Delta x + (\varepsilon_2 + i\varepsilon_4)\Delta y.$$

Proof (Cont.)

By the Cauchy–Riemann equations,

$$\frac{\partial u}{\partial y} + i\frac{\partial v}{\partial y} = -\frac{\partial v}{\partial x} + i\frac{\partial u}{\partial x} = i\left(\frac{\partial u}{\partial x} + i\frac{\partial v}{\partial x}\right).$$

Hence

$$\frac{f(z+\Delta z) - f(z)}{\Delta z} = \frac{\partial u}{\partial x} + i\frac{\partial v}{\partial x} + (\varepsilon_1 + i\varepsilon_3)\frac{\Delta x}{\Delta z} + (\varepsilon_2 + i\varepsilon_4)\frac{\Delta y}{\Delta z}.$$

Proof (Cont.)

Since $\left|\frac{\Delta x}{\Delta z}\right| \leq 1$ and $\left|\frac{\Delta y}{\Delta z}\right| \leq 1$,

$$\lim_{\Delta z \to 0} \left[(\varepsilon_1 + i\varepsilon_3)\frac{\Delta x}{\Delta z} + (\varepsilon_2 + i\varepsilon_4)\frac{\Delta y}{\Delta z} \right] = 0.$$

Therefore,

$$f'(z) = \lim_{\Delta z \to 0} \frac{f(z+\Delta z) - f(z)}{\Delta z} = \frac{\partial u}{\partial x} + i\frac{\partial v}{\partial x},$$

so $f(z)$ is differentiable at $z = x + iy$.



By the proof of Theorem 2.1, if $f(z) = u + iv$ is differentiable at $z = x + iy$, then

$$f'(z) = \frac{\partial u}{\partial x} + i\frac{\partial v}{\partial x} = \frac{\partial v}{\partial y} - i\frac{\partial u}{\partial y}. \tag{2.2.1}$$

Theorem 2.2

A function $f(z) = u(x, y) + iv(x, y)$ is analytic in a domain D if and only if $u(x, y)$ and $v(x, y)$ are differentiable in D and satisfy the Cauchy–Riemann equations in D.

How to show that a function is analytic:

1) Use the definition of the derivative and the rules of differentiation to show that $f'(z)$ exists at every point of D.
2) Show that the first-order partial derivatives of u and v exist and are continuous in D (so u and v are differentiable) and satisfy the Cauchy–Riemann equations. Then $f(z)$ is analytic in D by Theorem 2.2.

2 Examples

Example 2.1

Determine where the following functions are differentiable and where they are analytic:

1. $w = \bar{z}$;
2. $f(z) = e^x(\cos y + i \sin y)$;
3. $w = z\mathrm{Re}(z)$.

2 Examples

Example 2.1

Determine where the following functions are differentiable and where they are analytic:

1. $w = \bar{z}$;
2. $f(z) = e^x(\cos y + i \sin y)$;
3. $w = z\mathrm{Re}(z)$.

Solution.

1. $u = x$, $v = -y$: $\dfrac{\partial u}{\partial x} = 1$, $\dfrac{\partial v}{\partial y} = -1$. The Cauchy–Riemann equations never hold, so $w = \bar{z}$ is differentiable nowhere and analytic nowhere.

Solution (Cont.)

2 $u = e^x \cos y$, $v = e^x \sin y$, and the four partial derivatives

$$\frac{\partial u}{\partial x} = e^x \cos y, \quad \frac{\partial u}{\partial y} = -e^x \sin y,$$
$$\frac{\partial v}{\partial x} = e^x \sin y, \quad \frac{\partial v}{\partial y} = e^x \cos y$$

are continuous everywhere and satisfy $u_x = v_y$, $u_y = -v_x$. Therefore, $f(z)$ is analytic in the whole plane, and by (2.2.1)

$$f'(z) = u_x + iv_x = e^x(\cos y + i \sin y) = f(z).$$

This is the exponential function e^z of Section 3.

Solution (Cont.)

3 $w = z\mathrm{Re}(z) = x^2 + ixy$, so $u = x^2$, $v = xy$, and

$$\frac{\partial u}{\partial x} = 2x, \quad \frac{\partial u}{\partial y} = 0, \quad \frac{\partial v}{\partial x} = y, \quad \frac{\partial v}{\partial y} = x$$

are continuous. The Cauchy–Riemann equations $2x = x$ and $0 = -y$ hold only at $x = y = 0$.
Hence $w = z\mathrm{Re}(z)$ is differentiable only at $z = 0$ and is analytic nowhere.

Example 2.2

Prove that $\overline{z}^2$ is analytic nowhere in the complex plane.

Example 2.2

Prove that $\bar{z}^2$ is analytic nowhere in the complex plane.

Proof.

$\bar{z}^2 = x^2 - y^2 - 2ixy$, so $u = x^2 - y^2$, $v = -2xy$, and

$$\frac{\partial u}{\partial x} = 2x, \quad \frac{\partial u}{\partial y} = -2y, \quad \frac{\partial v}{\partial x} = -2y, \quad \frac{\partial v}{\partial y} = -2x.$$

The Cauchy–Riemann equations $2x = -2x$ and $-2y = 2y$ hold only at $x = y = 0$. Hence $\bar{z}^2$ is differentiable only at $z = 0$, and it is analytic nowhere. □

Example 2.3

Let $f(z) = x^2 + axy + by^2 + i(cx^2 + dxy + y^2)$. *Find the constants* a, b, c, *and* d *such that* $f(z)$ *is analytic in the whole plane.*

Example 2.3

Let $f(z) = x^2 + axy + by^2 + i(cx^2 + dxy + y^2)$. Find the constants a, b, c, and d such that $f(z)$ is analytic in the whole plane.

Solution.

$$\frac{\partial u}{\partial x} = 2x + ay, \quad \frac{\partial u}{\partial y} = ax + 2by, \quad \frac{\partial v}{\partial x} = 2cx + dy, \quad \frac{\partial v}{\partial y} = dx + 2y.$$

The Cauchy–Riemann equations $\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$ and $\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$ require, for all x and y,

$$2x + ay = dx + 2y, \qquad ax + 2by = -2cx - dy.$$

Comparing coefficients gives $a = 2$, $b = -1$, $c = -1$, $d = 2$.

Example 2.4

Prove that $f(z) = \sqrt{|xy|}$ *satisfies the Cauchy–Riemann equations at* $z = 0$ *but is not differentiable at* $z = 0$.

Example 2.4

Prove that $f(z) = \sqrt{|xy|}$ satisfies the Cauchy–Riemann equations at $z = 0$ but is not differentiable at $z = 0$.

Proof.

Here $u = \sqrt{|xy|}$ and $v = 0$. At the origin,

$$u_x(0,0) = \lim_{x \to 0} \frac{u(x,0) - u(0,0)}{x} = 0 = v_y(0,0),$$

$$u_y(0,0) = \lim_{y \to 0} \frac{u(0,y) - u(0,0)}{y} = 0 = -v_x(0,0),$$

so the Cauchy–Riemann equations hold at $z = 0$.

Proof (Cont.)

However, as $z \to 0$ along the ray $y = kx$ $(x > 0,\ k > 0)$,

$$\frac{f(z) - f(0)}{z - 0} = \frac{\sqrt{k}\,x}{x + ikx} = \frac{\sqrt{k}}{1 + ik},$$

which depends on k. Therefore, $\lim_{z \to 0} \dfrac{f(z) - f(0)}{z}$ does not exist, and $f(z)$ is not differentiable at $z = 0$.

The Cauchy–Riemann equations alone are not sufficient: u must also be differentiable, and $\sqrt{|xy|}$ is not differentiable at $(0, 0)$. □

Example 2.5

Let $f(z) = u(x, y) + iv(x, y)$ be analytic in a domain D with $v = u^2$. Find $f(z)$.

Example 2.5

Let $f(z) = u(x, y) + iv(x, y)$ be analytic in a domain D with $v = u^2$. Find $f(z)$.

Solution.

By the Cauchy–Riemann equations,

$$u_x = v_y = 2uu_y, \tag{1}$$

$$u_y = -v_x = -2uu_x. \tag{2}$$

Substituting (2) into (1) gives $u_x(1 + 4u^2) = 0$. Since $1 + 4u^2 \neq 0$, $u_x = 0$, and then (2) gives $u_y = 0$.
Hence $u = c$ is a real constant in D, and $f(z) = c + ic^2$ is constant.

Example 2.6 (Classroom practice)

Let $f(z) = my^3 + nx^2y + i(x^3 + lxy^2)$ *be analytic. Determine* l, m, *and* n.

Example 2.6 (Classroom practice)

Let $f(z) = my^3 + nx^2y + i(x^3 + lxy^2)$ *be analytic. Determine* l, m, *and* n.

Solution.

$u_x = 2nxy = v_y = 2lxy$ gives $n = l$;
$u_y = 3my^2 + nx^2 = -v_x = -3x^2 - ly^2$ gives $n = -3$ and $3m = -l$.
Hence

$$l = n = -3, \qquad m = 1.$$

Example 2.7

Suppose $f'(z) = 0$ everywhere in a domain D. Prove that $f(z)$ is constant in D.

Example 2.7

Suppose $f'(z) = 0$ everywhere in a domain D. Prove that $f(z)$ is constant in D.

Proof.

By (2.2.1),

$$f'(z) = \frac{\partial u}{\partial x} + i\frac{\partial v}{\partial x} = \frac{\partial v}{\partial y} - i\frac{\partial u}{\partial y} \equiv 0,$$

so $\frac{\partial u}{\partial x} = \frac{\partial u}{\partial y} = \frac{\partial v}{\partial x} = \frac{\partial v}{\partial y} \equiv 0$ in D. Since D is connected, u and v are constant, and hence $f(z)$ is constant in D. □

Example 2.8

Let $f(z) = u + iv$ be analytic with $f'(z) \neq 0$. Prove that the curve families $u(x, y) = c_1$ and $v(x, y) = c_2$ are orthogonal to each other, where c_1 and c_2 are constants.

Example 2.8

Let $f(z) = u + iv$ be analytic with $f'(z) \neq 0$. Prove that the curve families $u(x, y) = c_1$ and $v(x, y) = c_2$ are orthogonal to each other, where c_1 and c_2 are constants.

Proof.

By (2.2.1), $f'(z) = v_y - iu_y \neq 0$, so u_y and v_y are not both zero. First suppose $u_y \neq 0$ and $v_y \neq 0$ at an intersection point. By implicit differentiation, the slopes of the two curves are

$$k_1 = -\frac{u_x}{u_y}, \qquad k_2 = -\frac{v_x}{v_y}.$$

Proof (Cont.)

By the Cauchy–Riemann equations $u_x = v_y$ and $v_x = -u_y$,

$$k_1 \cdot k_2 = \left(-\frac{u_x}{u_y}\right) \cdot \left(-\frac{v_x}{v_y}\right) = \frac{v_y}{u_y} \cdot \frac{-u_y}{v_y} = -1,$$

so the two curves are orthogonal.
If one of u_y and v_y is zero, say $u_y = 0$ and $v_y \neq 0$, then $v_x = -u_y = 0$. Thus the tangent to $u = c_1$ is vertical and the tangent to $v = c_2$ is horizontal, and the curves are again orthogonal. □

Example 2.9

Prove that the real and imaginary parts of $f(z) = \sqrt{|\operatorname{Im} z^2|}$ *satisfy the Cauchy–Riemann equations at* $z = 0$*, but* $f(z)$ *is not differentiable at* $z = 0$*.*

Example 2.9

Prove that the real and imaginary parts of $f(z) = \sqrt{|\operatorname{Im} z^2|}$ *satisfy the Cauchy–Riemann equations at* $z = 0$*, but* $f(z)$ *is not differentiable at* $z = 0$*.*

Proof.

Since $\operatorname{Im} z^2 = 2xy$, we have $u = \sqrt{|2xy|}$ and $v = 0$. At the origin,

$$u_x(0,0) = \lim_{x \to 0} \frac{u(x,0) - u(0,0)}{x} = 0 = v_y(0,0),$$

$$u_y(0,0) = \lim_{y \to 0} \frac{u(0,y) - u(0,0)}{y} = 0 = -v_x(0,0),$$

so the Cauchy–Riemann equations hold at $z = 0$.

Proof (Cont.)

However,

$$\frac{f(\Delta z) - f(0)}{\Delta z} = \frac{\sqrt{|2\Delta x \Delta y|}}{\Delta x + i\Delta y}.$$

Along $\Delta y = \Delta x \to 0^+$, this quotient equals $\frac{\sqrt{2}}{1+i}$; along $\Delta x = 0$, $\Delta y \to 0$, it equals 0.
The two limits differ, so $f(z)$ is not differentiable at $z = 0$. □

3 Summary

A function $f(z) = u+iv$ is analytic in D if and only if $u(x, y)$ and $v(x, y)$ are differentiable in D and satisfy the Cauchy–Riemann equations

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \qquad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}.$$

In that case, $f'(z) = u_x + iv_x = v_y - iu_y$.

Example 2.10 (Question)

What should we check when using the Cauchy–Riemann equations to study the analyticity of $f(z) = u(x, y) + iv(x, y)$*?*

Solution.

1. Check whether $u(x, y)$ and $v(x, y)$ are differentiable in D (for example, whether their first-order partial derivatives are continuous);
2. Check whether they satisfy the Cauchy–Riemann equations

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \qquad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x};$$

3. Differentiability at isolated points or on a curve does not give analyticity: analyticity requires differentiability in a neighborhood.

Outline

1. Concept of Analytic Functions

2. Necessary and Sufficient Conditions for Analyticity

3. Elementary Functions

1 Exponential Function

1 Definition

For every complex number $z = x + iy$, the exponential function (指数函数) is defined by

$$\exp z = e^z = e^{x+iy} = e^x(\cos y + i \sin y).$$

- e^z is a notation for this function, not a power of the number e (powers are defined later).
- When $x = 0$, it reduces to Euler's formula $e^{iy} = \cos y + i \sin y$.
- When $y = 0$, it reduces to the real exponential function e^x.

2 Properties

1) $|e^z| = e^x$ and $\operatorname{Arg} e^z = y + 2k\pi$ $(k \in \mathbb{Z})$; in particular, $e^z \neq 0$.
2) $e^{z_1+z_2} = e^{z_1} e^{z_2}$ and $\dfrac{e^{z_1}}{e^{z_2}} = e^{z_1-z_2}$.

   However, $(e^{z_1})^{z_2} = e^{z_1 z_2}$ does not hold in general. For example, $(e^{i\pi})^{1/2} = (-1)^{1/2} = \pm i$, whereas $e^{i\pi/2} = i$.
3) e^z is analytic in the whole plane and $(e^z)' = e^z$.
4) e^z is periodic with fundamental period $2\pi i$: $e^{z+2k\pi i} = e^z$.

Example 3.1

Let $z = x + iy$. *Find*

1) $\left| e^{i-2z} \right|$;
2) $\left| e^{z^2} \right|$;
3) $\mathrm{Re}(e^{1/z})$.

Example 3.1

Let $z = x + iy$. *Find*

1) $\left|e^{i-2z}\right|$;
2) $\left|e^{z^2}\right|$;
3) $\mathrm{Re}(e^{1/z})$.

Solution.

Since $e^{x+iy} = e^x(\cos y + i \sin y)$, we have $\left|e^{x+iy}\right| = e^x$ and $\mathrm{Re}(e^{x+iy}) = e^x \cos y$.

Solution (Cont.)

1) $e^{i-2z} = e^{i-2(x+iy)} = e^{-2x+i(1-2y)}$, so $\left|e^{i-2z}\right| = e^{-2x}$.

2) $e^{z^2} = e^{(x+iy)^2} = e^{x^2-y^2+2ixy}$, so $\left|e^{z^2}\right| = e^{x^2-y^2}$.

3) $\dfrac{1}{z} = \dfrac{x-iy}{x^2+y^2}$, so $e^{1/z} = e^{\frac{x}{x^2+y^2} - i\frac{y}{x^2+y^2}}$ and

$$\operatorname{Re}\left(e^{1/z}\right) = e^{\frac{x}{x^2+y^2}} \cos \frac{y}{x^2+y^2}.$$

Example 3.2

Find the principal arguments of the following complex numbers:
(1) e^{2+i}; *(2)* e^{2-3i}; *(3)* e^{3+4i}; *(4)* e^{-3-4i};
(5) $e^{i\alpha} - e^{i\beta} \quad (0 \leq \beta < \alpha \leq 2\pi)$.

Example 3.2

Find the principal arguments of the following complex numbers:
(1) e^{2+i}; *(2)* e^{2-3i}; *(3)* e^{3+4i}; *(4)* e^{-3-4i};
(5) $e^{i\alpha} - e^{i\beta} \quad (0 \le \beta < \alpha \le 2\pi)$.

Solution.

Since $\operatorname{Arg} e^{x+iy} = y + 2k\pi \ (k \in \mathbb{Z})$, the principal argument $\arg e^{x+iy}$ is the value of $y + 2k\pi$ in $(-\pi, \pi]$.

1. $\operatorname{Arg} e^{2+i} = 1 + 2k\pi$, so $\arg e^{2+i} = 1$;
2. $\operatorname{Arg} e^{2-3i} = -3 + 2k\pi$, so $\arg e^{2-3i} = -3$;

Solution (Cont.)

(3) $\operatorname{Arg} e^{3+4i} = 4 + 2k\pi$; since $4 > \pi$, $\arg e^{3+4i} = 4 - 2\pi$;

(4) $\operatorname{Arg} e^{-3-4i} = -4+2k\pi$; since $-4 < -\pi$, $\arg e^{-3-4i} = -4+2\pi$;

(5)
$$\begin{aligned} e^{i\alpha} - e^{i\beta} &= (\cos\alpha - \cos\beta) + i(\sin\alpha - \sin\beta) \\ &= -2\sin\frac{\alpha+\beta}{2}\sin\frac{\alpha-\beta}{2} + 2i\cos\frac{\alpha+\beta}{2}\sin\frac{\alpha-\beta}{2} \\ &= 2\sin\frac{\alpha-\beta}{2}\left(-\sin\frac{\alpha+\beta}{2} + i\cos\frac{\alpha+\beta}{2}\right) \\ &= 2\sin\frac{\alpha-\beta}{2}\left(\cos\frac{\pi+\alpha+\beta}{2} + i\sin\frac{\pi+\alpha+\beta}{2}\right). \end{aligned}$$

Solution (Cont.)

Since $0 \le \beta < \alpha \le 2\pi$, we have $\sin \dfrac{\alpha - \beta}{2} > 0$, so the last expression is the trigonometric form of $e^{i\alpha} - e^{i\beta}$. Hence

$$\operatorname{Arg}(e^{i\alpha} - e^{i\beta}) = \frac{\pi + \alpha + \beta}{2} + 2k\pi,$$

and

$$\arg(e^{i\alpha} - e^{i\beta}) = \begin{cases} \dfrac{\pi + \alpha + \beta}{2}, & \alpha + \beta \le \pi, \\ \dfrac{\pi + \alpha + \beta}{2} - 2\pi, & \alpha + \beta > \pi. \end{cases}$$

Example 3.3

Find the period of $f(z) = e^{z/5}$.

Example 3.3

Find the period of $f(z) = e^{z/5}$.

Solution.

Since e^w has periods $2k\pi i$ $(k \in \mathbb{Z})$,

$$f(z) = e^{\frac{z}{5}} = e^{\frac{z}{5}+2k\pi i} = e^{\frac{z+10k\pi i}{5}} = f(z+10k\pi i).$$

Hence the periods of $f(z)$ are $10k\pi i$ $(k \neq 0)$, and its fundamental period is $10\pi i$.

2 Logarithmic Function

1. Definition
   A function $w = f(z)$ satisfying $e^w = z$ $(z \neq 0)$ is called the logarithmic function (对数函数) and is denoted by $w = \operatorname{Ln} z$.
2. Let $w = u + iv$ and $z = re^{i\theta}$. Then

$$e^w = z \iff e^u e^{iv} = re^{i\theta} \iff \begin{cases} e^u = r, \\ v = \theta + 2k\pi \ (k \in \mathbb{Z}), \end{cases}$$

so $u = \ln r$ and $v = \theta + 2k\pi$. Therefore,

$$\operatorname{Ln} z = \ln |z| + i \operatorname{Arg} z.$$

Since $\operatorname{Arg} z$ is multiple-valued, so is $\operatorname{Ln} z$.

If $\operatorname{Arg} z$ is replaced by the principal argument $\arg z$, we obtain a single-valued function, the principal value（主値）of $\operatorname{Ln} z$:

$$\ln z = \ln |z| + i \arg z.$$

All the values are

$$\operatorname{Ln} z = \ln z + 2k\pi i \quad (k \in \mathbb{Z}).$$

For each fixed k, this formula defines a single-valued function called a branch（分支）of $\operatorname{Ln} z$.
When $z = x > 0$, the principal value is $\ln z = \ln x$, the real logarithm.

Example 3.4

Find $\operatorname{Ln} 2$ *and* $\operatorname{Ln}(-1)$ *and their principal values.*

Example 3.4

Find $\operatorname{Ln} 2$ *and* $\operatorname{Ln}(-1)$ *and their principal values.*

Solution.

$\operatorname{Ln} 2 = \ln 2 + 2k\pi i \ (k \in \mathbb{Z})$, and its principal value is $\ln 2$.
$\operatorname{Ln}(-1) = \ln 1 + i\operatorname{Arg}(-1) = (2k+1)\pi i \ (k \in \mathbb{Z})$, and its principal value is πi.

Note: A negative real number has no real logarithm, but it has complex logarithms. The complex logarithm extends the real logarithm.

Example 3.5

Solve the equation $e^z - 1 - \sqrt{3}i = 0$.

Example 3.5

Solve the equation $e^z - 1 - \sqrt{3}i = 0$.

Solution.

Since $e^z = 1 + \sqrt{3}i$,

$$z = \operatorname{Ln}(1 + \sqrt{3}i) = \ln\left|1 + \sqrt{3}i\right| + i\operatorname{Arg}(1 + \sqrt{3}i)$$
$$= \ln 2 + i\left(\frac{\pi}{3} + 2k\pi\right) \quad (k \in \mathbb{Z}).$$

Example 3.6

Find (1) $\mathrm{Ln}(-2+3i)$*; (2)* $\mathrm{Ln}(3-\sqrt{3}i)$*; (3)* $\mathrm{Ln}(-3)$.

Example 3.6

Find (1) $\mathrm{Ln}(-2+3i)$*; (2)* $\mathrm{Ln}(3-\sqrt{3}i)$*; (3)* $\mathrm{Ln}(-3)$.

Solution.

(1) $-2+3i$ lies in the second quadrant, so $\arg(-2+3i) = \pi - \arctan\frac{3}{2}$ and

$$\mathrm{Ln}(-2+3i) = \frac{1}{2}\ln 13 + i\left(\pi - \arctan\frac{3}{2} + 2k\pi\right) \quad (k \in \mathbb{Z}).$$

Solution (Cont.)

(2) $\left|3-\sqrt{3}i\right| = 2\sqrt{3}$ and $\arg(3-\sqrt{3}i) = \arctan\dfrac{-\sqrt{3}}{3} = -\dfrac{\pi}{6}$, so

$$\operatorname{Ln}(3-\sqrt{3}i) = \ln(2\sqrt{3}) + i\left(2k\pi - \frac{\pi}{6}\right) \quad (k\in\mathbb{Z}).$$

(3) $|-3| = 3$ and $\arg(-3) = \pi$, so

$$\operatorname{Ln}(-3) = \ln 3 + (2k+1)\pi i \quad (k\in\mathbb{Z}).$$

3 Properties

1) $\operatorname{Ln}(z_1 z_2) = \operatorname{Ln} z_1 + \operatorname{Ln} z_2$;
2) $\operatorname{Ln} \dfrac{z_1}{z_2} = \operatorname{Ln} z_1 - \operatorname{Ln} z_2$.
   These are equalities of sets, like $\operatorname{Arg}(z_1 z_2) = \operatorname{Arg} z_1 + \operatorname{Arg} z_2$; for principal values, $\ln(z_1 z_2) \neq \ln z_1 + \ln z_2$ in general.
3) In the plane with the origin and the negative real axis removed, the principal branch and every other branch are continuous and differentiable, and

$$(\ln z)' = \frac{1}{z}, \qquad (\operatorname{Ln} z)' = \frac{1}{z}.$$

Proof of (3).

Let $z = x + iy$. For $x < 0$,

$$\lim_{y \to 0^-} \arg z = -\pi, \qquad \lim_{y \to 0^+} \arg z = \pi,$$

so $\arg z$, and hence $\ln z$, is discontinuous on the negative real axis. Elsewhere, except at the origin, $\ln z$ is continuous.
In the region $-\pi < \arg z < \pi$, $w = \ln z$ is the single-valued inverse of $z = e^w$, so

$$\frac{\mathrm{d} \ln z}{\mathrm{d} z} = \frac{1}{\frac{\mathrm{d} e^w}{\mathrm{d} w}} = \frac{1}{e^w} = \frac{1}{z}.$$



3 Powers a^b and Power Functions

1 Definition of a^b

Let $a \neq 0$ and let b be any complex number. The power (乘冪) a^b is defined by

$$a^b = e^{b \operatorname{Ln} a}.$$

Since $\operatorname{Ln} a = \ln|a| + i(\arg a + 2k\pi)$ is multiple-valued, a^b is multiple-valued in general.

Replacing $\operatorname{Ln} a$ with its principal value $\ln a$ gives the principal value of a^b:

$$e^{b \ln a} = e^{b[\ln|a| + i \arg a]}.$$

1) If b is an integer, then $e^{2kb\pi i} = 1$ and

$$a^b = e^{b[\ln|a| + i(\arg a + 2k\pi)]} = e^{b \ln a}\, e^{2kb\pi i} = e^{b \ln a},$$

so a^b is single-valued.

2) If $b = \dfrac{p}{q}$, where p and q are coprime integers and $q > 0$, then

$$\begin{aligned} a^b &= e^{\frac{p}{q}\ln|a| + i\frac{p}{q}(\arg a + 2k\pi)} \\ &= e^{\frac{p}{q}\ln|a|}\left[\cos\frac{p}{q}(\arg a + 2k\pi) + i\sin\frac{p}{q}(\arg a + 2k\pi)\right]. \end{aligned}$$

It has exactly q distinct values, obtained for $k = 0, 1, \ldots, q-1$.

3) If b is irrational or non-real, a^b has infinitely many values.

Special cases:

1) If $b = n$ is a positive integer,

$$a^n = e^{n \ln a} = \left(e^{\ln a}\right)^n = \underbrace{a \cdot a \cdots a}_{n \text{ factors}},$$

which agrees with the power defined in Chapter 1.

2) If $b = \dfrac{1}{n}$,

$$a^{\frac{1}{n}} = |a|^{\frac{1}{n}} \left[\cos \frac{\arg a + 2k\pi}{n} + i \sin \frac{\arg a + 2k\pi}{n}\right] = \sqrt[n]{a},$$

$k = 0, 1, \ldots, n-1$, which are exactly the n values of the n-th root in (1.2.10).

If $a = z$ is a complex variable, we obtain the power function (冪函数) $w = z^b$.
For a fixed exponent b, replacing $\operatorname{Ln} z$ with $\ln z$ gives the principal branch (主支)

$$w = e^{b \ln z}, \qquad -\pi < \arg z < \pi,$$

which is analytic in the cut plane $\mathbb{C} \setminus (-\infty, 0]$.
For $b = n$ and $b = \dfrac{1}{n}$, we obtain $w = z^n$ and its inverse, the n-th root function $w = z^{\frac{1}{n}} = \sqrt[n]{z}$.

Example 3.7

Find the values of $1^{\sqrt{2}}$ *and* i^i.

Example 3.7

Find the values of $1^{\sqrt{2}}$ *and* i^i.

Solution.

$$1^{\sqrt{2}} = e^{\sqrt{2}\,\mathrm{Ln}\,1} = e^{2\sqrt{2}k\pi i} = \cos(2\sqrt{2}k\pi) + i\sin(2\sqrt{2}k\pi) \quad (k \in \mathbb{Z}),$$

$$i^i = e^{i\,\mathrm{Ln}\,i} = e^{i\left(\frac{\pi}{2}+2k\pi\right)i} = e^{-\left(\frac{\pi}{2}+2k\pi\right)} \quad (k \in \mathbb{Z}).$$

All values of i^i are positive real numbers.

Example 3.8 (Classroom practice)

Find the values of $(-3)^{\sqrt{5}}$.

Example 3.8 (Classroom practice)

Find the values of $(-3)^{\sqrt{5}}$.

Solution.

$$
\begin{aligned}
(-3)^{\sqrt{5}} &= e^{\sqrt{5}\,\mathrm{Ln}(-3)} = e^{\sqrt{5}[\ln 3+(2k+1)\pi i]} \\
&= 3^{\sqrt{5}}\left[\cos\sqrt{5}(2k+1)\pi + i\sin\sqrt{5}(2k+1)\pi\right] \quad (k\in\mathbb{Z}).
\end{aligned}
$$

Example 3.9

Find the principal argument of $(1+i)^i$.

Example 3.9

Find the principal argument of $(1+i)^i$.

Solution.

$$(1+i)^i = e^{i\,\mathrm{Ln}(1+i)} = e^{i\left[\frac{1}{2}\ln 2 + i\left(\frac{\pi}{4}+2k\pi\right)\right]}$$
$$= e^{-\left(\frac{\pi}{4}+2k\pi\right)}\left[\cos\left(\frac{1}{2}\ln 2\right) + i\sin\left(\frac{1}{2}\ln 2\right)\right] \quad (k\in\mathbb{Z}).$$

All values have modulus $e^{-\left(\frac{\pi}{4}+2k\pi\right)} > 0$ and the same argument $\frac{1}{2}\ln 2 \in (-\pi,\pi]$. Therefore, the principal argument of $(1+i)^i$ is $\frac{1}{2}\ln 2$.

2 Analyticity of Power Functions

1) z^n is single-valued and analytic in the whole plane, and $(z^n)' = nz^{n-1}$.
2) $z^{\frac{1}{n}} = e^{\frac{1}{n}\operatorname{Ln} z}$ is multiple-valued with n branches. Its principal branch is $e^{\frac{1}{n}\ln z}$. Each branch is analytic in the plane with the origin and the negative real axis removed, and

$$\left(z^{\frac{1}{n}}\right)' = \left(e^{\frac{1}{n}\operatorname{Ln} z}\right)' = \frac{1}{n} z^{\frac{1}{n}-1}.$$

3) For other exponents b, $z^b = e^{b\operatorname{Ln} z}$ is multiple-valued (with infinitely many branches if b is irrational or non-real). Its principal branch is $e^{b\ln z}$. Each branch is analytic in the same cut plane, and

$$\left(z^b\right)' = bz^{b-1}.$$

4 Trigonometric and Hyperbolic Functions

1 Definition of Trigonometric Functions

For real y, $e^{iy} = \cos y + i \sin y$ and $e^{-iy} = \cos y - i \sin y$. Adding and subtracting gives

$$\cos y = \frac{e^{iy} + e^{-iy}}{2}, \qquad \sin y = \frac{e^{iy} - e^{-iy}}{2i}.$$

Replacing the real number y by a complex number z extends cosine and sine to the complex plane.

The cosine and sine functions (余弦、正弦函数) are defined by

$$\cos z = \frac{e^{iz} + e^{-iz}}{2}, \qquad \sin z = \frac{e^{iz} - e^{-iz}}{2i}.$$

$\sin z$ is odd and $\cos z$ is even:

$$\sin(-z) = -\sin z, \qquad \cos(-z) = \cos z.$$

Both are periodic with fundamental period 2π:

$$\sin(z + 2\pi) = \sin z, \qquad \cos(z + 2\pi) = \cos z.$$

Example 3.10

Find the period of $f(z) = \sin 5z$.

Example 3.10

Find the period of $f(z) = \sin 5z$.

Solution.

Since $\sin w$ has period 2π,

$$f\left(z + \frac{2\pi}{5}\right) = \sin(5z + 2\pi) = \sin 5z = f(z).$$

Therefore, the fundamental period of $f(z) = \sin 5z$ is $\dfrac{2\pi}{5}$.

Sine and cosine are analytic in the whole plane, and

$$(\sin z)' = \cos z, \qquad (\cos z)' = -\sin z.$$

Important formulas:

(1
$$\begin{cases} \cos(z_1 + z_2) = \cos z_1 \cos z_2 - \sin z_1 \sin z_2, \\ \sin(z_1 + z_2) = \sin z_1 \cos z_2 + \cos z_1 \sin z_2, \\ \sin^2 z + \cos^2 z = 1; \end{cases}$$

(2
$$\begin{cases} \cos(x + iy) = \cos x \cos iy - \sin x \sin iy, \\ \sin(x + iy) = \sin x \cos iy + \cos x \sin iy. \end{cases}$$

(3) For a purely imaginary number iy,

$$\cos iy = \frac{e^{-y} + e^{y}}{2} = \cosh y, \qquad \sin iy = \frac{e^{-y} - e^{y}}{2i} = i \sinh y.$$

Here $\cosh$ and $\sinh$ denote the hyperbolic cosine（双曲余弦) and hyperbolic sine（双曲正弦）, respectively.

(4) Hence

$$\begin{cases} \cos(x + iy) = \cos x \cosh y - i \sin x \sinh y, \\ \sin(x + iy) = \sin x \cosh y + i \cos x \sinh y. \end{cases}$$

As $y \to \infty$, $|\sin iy| \to \infty$ and $|\cos iy| \to \infty$. Unlike the real case, $\sin z$ and $\cos z$ are unbounded.

The other trigonometric functions are defined by

$$\tan z = \frac{\sin z}{\cos z}, \quad \cot z = \frac{\cos z}{\sin z}, \quad \sec z = \frac{1}{\cos z}, \quad \csc z = \frac{1}{\sin z}.$$

Their periodicity, parity, and analyticity can be studied in the same way as for $\sin z$ and $\cos z$.

All four functions are unbounded on their domains. The absolute values of $\tan z$ and $\sec z$ can become arbitrarily large as z approaches $\frac{\pi}{2} + k\pi$; the same is true of $\cot z$ and $\csc z$ as z approaches $k\pi$.

Example 3.11

Find the real and imaginary parts of $\tan z$.

Example 3.11

Find the real and imaginary parts of $\tan z$.

Solution.

Let $z = x + iy$. Then

$$\tan z = \frac{\sin(x+iy)}{\cos(x+iy)} = \frac{\sin x \cosh y + i \cos x \sinh y}{\cos x \cosh y - i \sin x \sinh y}.$$

Solution (Cont.)

Multiplying numerator and denominator by the conjugate of the denominator,

$$\begin{aligned}\tan z &= \frac{\sin x\cos x + i\sinh y\cosh y}{\cos^2 x\cosh^2 y + \sin^2 x\sinh^2 y} \\ &= \frac{\sin 2x + i\sinh 2y}{\cos 2x + \cosh 2y},\end{aligned}$$

using

$$\cos^2 x\cosh^2 y + \sin^2 x\sinh^2 y = \cos^2 x + \sinh^2 y = \frac{\cos 2x + \cosh 2y}{2}.$$

Hence

$$\mathrm{Re}(\tan z) = \frac{\sin 2x}{\cos 2x + \cosh 2y}, \qquad \mathrm{Im}(\tan z) = \frac{\sinh 2y}{\cos 2x + \cosh 2y}.$$

Example 3.12

Solve the equation $\sin z = i \sinh 1$.

Example 3.12

Solve the equation $\sin z = i \sinh 1$.

Solution.

Let $z = x + iy$. Then $\sin x \cosh y + i \cos x \sinh y = i \sinh 1$, so

$$\sin x \cosh y = 0, \qquad \cos x \sinh y = \sinh 1.$$

Since $\cosh y > 0$, $\sin x = 0$ and $x = k\pi$. Then $\sinh y = (-1)^k \sinh 1$, so $y = (-1)^k$. Therefore,

$$z = k\pi + i(-1)^k, \qquad k \in \mathbb{Z}.$$

Example 3.13

Find the values of $\cos(1+i)$ *and* $\tan(3-i)$.

Example 3.13

Find the values of $\cos(1+i)$ *and* $\tan(3-i)$.

Solution.

$$\begin{aligned}
\cos(1+i) &= \frac{e^{i(1+i)} + e^{-i(1+i)}}{2} = \frac{e^{-1+i} + e^{1-i}}{2} \\
&= \frac{1}{2}\left[e^{-1}(\cos 1 + i\sin 1) + e(\cos 1 - i\sin 1)\right] \\
&= \frac{e+e^{-1}}{2}\cos 1 - i\frac{e-e^{-1}}{2}\sin 1 \\
&= \cos 1 \cosh 1 - i \sin 1 \sinh 1.
\end{aligned}$$

Solution (Cont.)

By the formula for $\tan z$ with $x = 3$ and $y = -1$,

$$\tan(3 - i) = \frac{\sin 6 + i \sinh(-2)}{\cos 6 + \cosh(-2)} = \frac{\sin 6 - i \sinh 2}{\cos 6 + \cosh 2}.$$

Directly, $\sin(3 - i) = \sin 3 \cosh 1 - i \cos 3 \sinh 1$ and $\cos(3 - i) = \cos 3 \cosh 1 + i \sin 3 \sinh 1$; multiplying by the conjugate of the denominator gives the same result.

2 Definition of Hyperbolic Functions
The hyperbolic cosine, sine, and tangent functions（双曲函数）are defined by

$$\cosh z = \frac{e^z + e^{-z}}{2}, \qquad \sinh z = \frac{e^z - e^{-z}}{2},$$

$$\tanh z = \frac{\sinh z}{\cosh z} = \frac{e^z - e^{-z}}{e^z + e^{-z}}.$$

For real z, they agree with the real hyperbolic functions from calculus.

$\sinh z$ is odd and $\cosh z$ is even. Both are periodic with fundamental period $2\pi i$, and

$$(\sinh z)' = \cosh z, \qquad (\cosh z)' = \sinh z.$$

Moreover,

$$\cosh iy = \cos y, \qquad \sinh iy = i\sin y,$$

$$\begin{cases} \cosh(x+iy) = \cosh x\cos y + i\sinh x\sin y, \\ \sinh(x+iy) = \sinh x\cos y + i\cosh x\sin y. \end{cases}$$

Example 3.14

Solve the equation $|\tanh z| = 1$.

Example 3.14

Solve the equation $|\tanh z| = 1$.

Solution.

$\tanh z = \dfrac{e^{2z}-1}{e^{2z}+1}$, so $|\tanh z| = 1$ means $\left|e^{2z}-1\right| = \left|e^{2z}+1\right|$. Let $e^{2z} = u + iv$. Squaring gives

$$(u-1)^2 + v^2 = (u+1)^2 + v^2 \iff u = 0.$$

Since $u = \mathrm{Re}(e^{2z}) = e^{2x}\cos 2y$,

$$u = 0 \iff \cos 2y = 0 \iff y = \frac{\pi}{4} + \frac{k\pi}{2} \quad (k \in \mathbb{Z}).$$

Hence the solutions are all z with $\mathrm{Im}\, z = \dfrac{\pi}{4} + \dfrac{k\pi}{2}$ $(k \in \mathbb{Z})$: a family of horizontal lines.

5 Summary

The complex elementary functions extend the real elementary functions. They keep many basic properties, but some new features appear:

1. The exponential function is periodic, with period $2\pi i$;
2. Negative numbers have logarithms, and $\operatorname{Ln} z$ is multiple-valued;
3. $\sin z$ and $\cos z$ are unbounded;
4. The hyperbolic sine and cosine are periodic, with period $2\pi i$.

Example 3.15 (Question)

What are the similarities and differences between real and complex trigonometric functions?

Solution.

- Similarities: parity, periodicity, differentiability, the formulas for derivatives, the addition formulas, and $\sin^2 z + \cos^2 z = 1$.
- Main difference: real sine and cosine are bounded by 1, but $|\sin z| \leq 1$ and $|\cos z| \leq 1$ fail for complex z. For example, $|\cos iy| = \cosh y \to \infty$.