

Frequency Analysis: Fourier Transform (Recap)

Richard Heusdens

April 24, 2016

1

EE2S31



Frequency Decomposition

Decomposition of signals in terms of sinusoidal (or complex exponential) components

- For periodic signals, this decomposition is called a *Fourier series*
- For the class of finite energy signals, this decomposition is called the *Fourier transform*

Why important?

- Complex exponentials are eigenfunctions of linear time-invariant systems
- The response of an LTI system to a sinusoidal input signal is a sinusoid of the same frequency but of different amplitude and phase

April 24, 2016

2

EE2S31



Fourier Series

Theorem: Let $\{e_k\}_{k=1}^{\infty}$ denote a complete orthonormal system in $L^2(E)$ and let $x \in L^2(E)$. Then

$$x = \sum_{k=1}^{\infty} (x, e_k) e_k$$

Moreover, we have (Parseval's identity)

$$\|x\|^2 = \sum_{k=1}^{\infty} |(x, e_k)|^2$$

The series is called the *Fourier series* w.r.t. the system $\{e_k\}$; the coefficients (x, e_k) are called the *Fourier coefficients*.

April 24, 2016

3

EE2S31



Complex Fourier series

The *Fourier series representation* of a periodic signal $x(t)$ of period T_0 is given by

$$x(t) = \sum_{k=-\infty}^{\infty} X_k e^{jk\Omega_0 t}, \quad \Omega_0 = \frac{2\pi}{T_0}$$

with *Fourier coefficients* X_k

$$X_k = \frac{1}{T_0} \int_{t_0}^{t_0+T_0} x(t) e^{-jk\Omega_0 t} dt, \quad k \in \mathbb{Z}$$

Moreover, we have (*Parseval's identity*): $\|x\|^2 = T_0 \sum_{k=-\infty}^{\infty} |X_k|^2$

April 24, 2016

4

EE2S31



Trigonometric Fourier series

The *Fourier series representation* of a periodic signal $x(t)$ of period T_0 is given by

$$x(t) = c_0 + 2 \sum_{k=0}^{\infty} (c_k \cos(k\Omega_0 t) + d_k \sin(k\Omega_0 t)), \quad \Omega_0 = \frac{2\pi}{T_0}$$

with *Fourier coefficients* c_k and d_k

$$c_k = \frac{1}{T_0} \int_{t_0}^{t_0+T_0} x(t) \cos(k\Omega_0 t) dt, \quad k = 0, 1, 2, \dots$$
$$d_k = \frac{1}{T_0} \int_{t_0}^{t_0+T_0} x(t) \sin(k\Omega_0 t) dt, \quad k = 1, 2, \dots$$

April 24, 2016

5

EE2S31



Convergence of the Fourier Series

$$\Omega_0 = 1$$

Example: $x(t) = t, t \in [\pi, \pi]$. Since x is odd, $c_k = 0$ for all k and

$$d_k = \frac{1}{2\pi} \int_{-\pi}^{\pi} t \sin(kt) dt$$
$$= \frac{-1}{2\pi k} t \cos(kt) \Big|_{-\pi}^{\pi} + \frac{1}{2\pi k} \underbrace{\int_{-\pi}^{\pi} \cos(kt) dt}_{=0} = \frac{1}{k} (-1)^{k+1}$$

Hence, the Fourier series becomes

$$\frac{2}{1} \sin(t) - \frac{2}{2} \sin(2t) + \frac{2}{3} \sin(3t) - \frac{2}{4} \sin(4t) + \dots$$

April 24, 2016


6

EE2S31



Convergence of the Fourier Series

- Notice that $\lim_{k \rightarrow \infty} d_k = 0$ and that the decay is $\mathcal{O}(k^{-1})$
- $x(\pi) = \pi$ but substituting $t = \pi$ in the Fourier series yields 0




April 24, 2016

7

EE2S31



Convergence of the Fourier Series



April 24, 2016

8

EE2S31



Convergence of the Fourier Series

Example: $x(t) = |t|, t \in [-\pi, \pi]$. Since x is even, $d_k = 0$ for all k

$$c_k = \frac{1}{\pi} \int_0^\pi t \cos(kt) dt$$
$$\stackrel{(k \neq 0)}{=} \underbrace{\frac{1}{k\pi} t \sin(kt) \Big|_0^\pi}_{=0} - \frac{1}{k\pi} \int_0^\pi \sin(kt) dt = \begin{cases} 0, & k = 2, 4, \dots \\ \frac{-2}{\pi k^2}, & k \text{ odd} \end{cases}$$

and

$$c_0 = \frac{1}{\pi} \int_0^\pi t dt = \frac{\pi}{2}$$

April 24, 2016

9

EE2S31




Convergence of the Fourier Series

The Fourier series becomes

$$\frac{\pi}{2} - \frac{4}{\pi} \left(\frac{\cos(t)}{1^2} + \frac{\cos(3t)}{3^2} + \frac{\cos(5t)}{5^2} + \dots \right)$$

Notice that $\lim_{k \rightarrow \infty} c_k = 0$ and that the decay is of order $\mathcal{O}(k^{-2})$




April 24, 2016

10

EE2S31



Convergence of the Fourier Series



April 24, 2016

11

EE2S31



Convergence of the Fourier Series

- If x is p times differentiable and all derivatives are in $L^1(E)$, then

$$x^{(p)}(t) \quad \xleftrightarrow{\mathcal{F}} \quad (jk\Omega_0)^p X_k$$

- Applying the Riemann-Lebesgue lemma on $x^{(p)}$, we conclude that

$$\lim_{k \rightarrow \pm\infty} (k\Omega_0)^p X_k = 0$$

so that regularity of x translates to rapid decay of X_k

This explains the faster decay of the Fourier coefficients of the function $x(t) = |t|$ as compared to those of $x(t) = t$.


April 24, 2016

12

EE2S31



Convergence of the Fourier Series



April 24, 2016

13

EE2S31



Fourier Transform

Some observations:

- Fourier series to represent periodic signals
- Periodic signals have a line spectrum
- The line spacing is equal to the fundamental frequency
- What happens if the fundamental period becomes infinite (fundamental frequency zero)? \rightarrow continuous spectrum?

From Fourier Series to Fourier Integral

April 24, 2016

14

EE2S31



Fourier transform

We have

$$X(\Omega) = \int_{-\infty}^{\infty} x(t)e^{-j\Omega t} dt, \quad \Omega \in \mathbb{R}$$
$$x(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} X(\Omega)e^{j\Omega t} d\Omega, \quad t \in \mathbb{R}$$

is called the *Fourier transform* pair.

April 24, 2016


15

EE2S31



Fourier transform

Example: $x(t) = u(t) - u(t - 1)$



Solution:

$$X(\Omega) = \int_0^1 e^{-j\Omega t} dt = \frac{1}{j\Omega} (1 - e^{-j\Omega}) = \frac{\sin(\Omega/2)}{\Omega/2} e^{-j\Omega/2}$$

$$|X(\Omega)| = \left| \frac{\sin(\Omega/2)}{\Omega/2} \right|, \quad \angle X(\Omega) = \angle \frac{\sin(\Omega/2)}{\Omega/2} - \Omega/2$$

April 24, 2016


16

EE2S31



Fourier transform

Solution:



April 24, 2016

17

EE2S31



Relation to Laplace transform

If the region of convergence (ROC) of $X(s) = \mathcal{L}\{x(t)\}$ contains the j -axis, then

$$X(\Omega) = \mathcal{F}\{x(t)\} = \int_{-\infty}^{\infty} x(t)e^{-j\Omega t} dt = \mathcal{L}\{x(t)\} \Big|_{s=j\Omega}$$

This holds for causal, anti-causal, and non-causal signals

April 24, 2016

18

EE2S31



Regularity and Fourier transform

- The decay of $X(\Omega)$ depends on the worst singular behaviour of x
- If x is p times differentiable and all derivatives are in $L^1(\mathbb{R})$, then

$$\lim_{\Omega \rightarrow \pm\infty} |\Omega|^p X(\Omega) = 0$$

so that regularity of x translates to rapid decay of $X(\Omega)$

April 24, 2016

19

EE2S31



Fourier transform of periodic signal

Representing a T_0 -periodic signal $x(t)$ by its Fourier series, we have the following Fourier transform pair

$$x(t) = \sum_{k=-\infty}^{\infty} X_k e^{jk\Omega_0 t} \quad \xleftrightarrow{\mathcal{F}} \quad X(\Omega) = \sum_{k=-\infty}^{\infty} 2\pi X_k \delta(\Omega - k\Omega_0)$$

Note that x has a line spectrum!

$$\Delta\Omega_k = \Omega_0$$

Example:

$$\mathcal{F}\{\cos(\Omega_0 t)\} = \mathcal{F}\left\{\frac{1}{2}e^{j\Omega_0 t} + \frac{1}{2}e^{-j\Omega_0 t}\right\} = \pi\delta(\Omega - \Omega_0) + \pi\delta(\Omega + \Omega_0)$$

April 24, 2016

20

EE2S31



Fourier transform of distributions

The Fourier transform is well defined for distributions as well. We have

- $\mathcal{F}\{\delta(t)\} = \mathcal{L}\{\delta(t)\}\Big|_{s=j\Omega} = 1\Big|_{s=j\Omega} = 1$
- $\mathcal{F}\{\delta'(t)\} = \mathcal{L}\{\delta'(t)\}\Big|_{s=j\Omega} = s\Big|_{s=j\Omega} = j\Omega$

April 24, 2016

21

EE2S31



Time-frequency relation

The support of $X(\Omega)$ is inversely proportional to the support of $x(t)$. If the Fourier transform $X(\Omega)$ of $x(t)$ exists and $a \neq 0 \in \mathbb{R}$, then $x(at)$

- is contracted ($|a| > 1$)
- is expanded ($-1 < a < 1$)
- reflected ($a < 0$)

We have

$$x(at) \quad \xleftrightarrow{\mathcal{F}} \quad \frac{1}{|a|} X\left(\frac{\Omega}{a}\right), \quad a \neq 0$$

April 24, 2016

22

EE2S31



Discrete-Time Fourier Transform

We have

$$X(\omega) = \sum_{n=-\infty}^{\infty} x(n)e^{-j\omega n}, \quad \omega \in [0, 2\pi)$$

$$x(n) = \frac{1}{2\pi} \int_0^{2\pi} X(\omega)e^{j\omega n} d\omega, \quad n \in \mathbb{Z}$$

is called the *discrete-time Fourier transform* pair.

April 24, 2016

23

EE2S31



Relation Different Transforms

		Continuous-time signals		Discrete-time signals	
		Time-domain	Frequency-domain	Time-domain	Frequency-domain
Periodic signals Fourier series	...				
	$(T_p = 2\pi)$	$c_k = \frac{1}{T_p} \int_{T_p} x_d(t) e^{-j2\pi k F_0 t} dt$	$F_0 = \frac{1}{T_p}$	$x(n) = \frac{1}{N} \sum_{n=0}^{N-1} x(n) e^{j(2\pi/N)kn}$	$x(n) = \sum_{k=0}^{N-1} c_k e^{j(2\pi/N)kn}$
Aperiodic signals Fourier transforms	Continuous and periodic			Discrete and periodic	Discrete and periodic
	$X_d(F) = \int_{-\infty}^{\infty} x_d(t) e^{-j2\pi F t} dt$	$x_d(t) = \int_{-\infty}^{\infty} X_d(F) e^{j2\pi F t} dF$			$X(\omega) = \frac{1}{2\pi} \int_{-\infty}^{\infty} X(\omega) e^{j\omega n} d\omega$
Continuous and aperiodic		Continuous and aperiodic	Continuous and aperiodic	Discrete and aperiodic	Continuous and periodic

Figure 4.3.1 Summary of analysis and synthesis formulas.

April 24, 2016

24

EE2S31



Relation to the \mathcal{Z} -Transform

Recall that the z -transform of a sequence x was given by

$$X(z) = \sum_{n=-\infty}^{\infty} x(n)z^{-n}$$

where $r_2 < |z| < r_1$ is the region of convergence. If $X(z)$ converges for $|z| = 1$, we have on the unit circle ($z = e^{j\omega}$, $-\pi \leq \omega < \pi$):

$$X(\omega) = \sum_{n=-\infty}^{\infty} x(n)e^{-j\omega n}$$

In conclusion, if the unit circle is contained in the ROC, the Fourier transform of x is obtained by evaluating the z -transform $X(z)$ on the unit circle!

April 24, 2016

25

EE2S31



Relation to the \mathcal{Z} -Transform

The inverse z -transform, where C is taken to be the unit circle, is given by

$$\begin{aligned} x(n) &= \frac{1}{2\pi j} \oint_C X(z)z^{n-1}dz \\ &= \frac{1}{2\pi j} \oint_C X(\omega)e^{j\omega(n-1)} \underbrace{d(e^{j\omega})}_{je^{j\omega}d\omega} \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} X(\omega)e^{j\omega n}d\omega \end{aligned}$$

which is consistent with the results obtained before.

April 24, 2016

26

EE2S31



Relation to the \mathcal{Z} -Transform




Figure 4.2.9 relationship between $X(z)$ and $X(\omega)$ for the sequence in Example 4.2.4, with $A = 1$ and $L = 10$

April 24, 2016

27

EE2S31



Application: Filtering

If $x(t)$ (periodic or not) is input to a BIBO stable LTI system with impulse response $h(t)$ (and transfer function $H(s)$), then the output of the system, say $y(t)$, is given by the convolution $y(t) = (x * h)(t)$ with Fourier transform

$$Y(\Omega) = X(\Omega)H(j\Omega)$$

If $x(t)$ is T_0 -periodic, the Fourier transform of the output is given by

$$Y(\Omega) = \sum_{k=-\infty}^{\infty} 2\pi X_k H(jk\Omega_0) \delta(\Omega - k\Omega_0)$$

with X_k de Fourier series coefficients of $x(t)$ and $\Omega_0 = \frac{2\pi}{T_0}$

April 24, 2016

EE2S31



Basics of filtering





- Filtering consists in getting rid of undesirable components of a signal, and/or emphasizing desired components
- Filter design: find $H(s) = B(s)/A(s)$ satisfying certain specifications, e.g., flat frequency response up to a certain frequency Ω_p , and sufficient suppression for $|\Omega| > \Omega_s$.
- Frequency discriminating filters keep the frequency components of interest and attenuate the rest
- Filter design is a *rational approximation* problem

April 24, 2016

EE2S31



Frequency selective filters



April 24, 2016

30

EE2S31



Frequency selective filters

- Ideal filters cannot be realised:

$$H(j\Omega) = \begin{cases} c e^{-j\Omega t_0}, & |\Omega| < \Omega_c \\ 0, & \text{otherwise} \end{cases} \xrightarrow{\mathcal{F}} h(t) = \frac{\sin(\Omega_c(t - t_0))}{\pi(t - t_0)}$$

Since $h \notin L^1(\mathbb{R})$, the filter is unstable; it can produce an unbounded output for a bounded input signal.

- Ideal filters are non-causal. If we approximate h by a causal filter, the frequency response cannot be zero in *any* frequency band
- ... but we can approximate the ideal filter closely

April 24, 2016

31


EE2S31



Frequency selective filters

Design specs:

- $f_p = 10$ kHz, $f_s = 12$ kHz, stop-band attenuation 80 dB



April 24, 2016


32

EE2S31



Frequency selective filters

Example: (initial conditions zero)



$$V_i(s) = V_R(s) + V_L(s) + V_C(s)$$

April 24, 2016

33

EE2S31



Frequency selective filters

High-pass filter: voltage across the inductor

$$V_c(s) = \frac{s^2 L C V_i(s)}{s^2 L C + s R C + 1}$$

so that the transfer function is given by

$$H(s) = \frac{s^2 L C}{s^2 L C + s R C + 1}$$

and thus

$$|H(j\Omega)| = \frac{\Omega^2 L C}{\sqrt{(1 - \Omega^2 L C)^2 + (\Omega R C)^2}}$$

April 24, 2016


34

EE2S31



Frequency selective filters

High-pass filter: voltage across the inductor



April 24, 2016

35

EE2S31



Frequency response and poles/zeros

Consider a rational transfer function

$$H(s) = \frac{s - z}{s - p}$$

Frequency response:

$$H(j\Omega) = \frac{j\Omega - z}{j\Omega - p} = \underbrace{\frac{|j\Omega - z|}{|j\Omega - p|}}_{|H(j\Omega)|} e^{j(\angle(j\Omega - z) - \angle(j\Omega - p))}$$
$$e^{j\angle H(j\Omega)}$$


April 24, 2016

36

EE2S31



Frequency response and poles/zeros



$$|H(j\Omega)| = \frac{B(\Omega)}{A(\Omega)}, \quad \angle H(j\Omega) = \psi(\Omega) - \theta(\Omega)$$

April 24, 2016

37

EE2S31



Frequency response and poles/zeros

More general, for a N th-order LTI system we have

$$\begin{aligned} H(j\Omega) &= \left. \frac{b_0 + b_1 s + \cdots + b_M s^M}{a_0 + a_1 s + \cdots + a_N s^N} \right|_{s=j\Omega} \\ &= \left. \frac{b_M}{a_N} \frac{(s - z_1)(s - z_2) \cdots (s - z_M)}{(s - p_1)(s - p_2) \cdots (s - p_N)} \right|_{s=j\Omega} \\ &= \frac{b_M}{a_N} \frac{(j\Omega - z_1)(j\Omega - z_2) \cdots (j\Omega - z_M)}{(j\Omega - p_1)(j\Omega - p_2) \cdots (j\Omega - p_N)} \\ &= \frac{b_M}{a_N} \frac{\prod_{k=1}^M (j\Omega - z_k)}{\prod_{k=1}^N (j\Omega - p_k)} \end{aligned}$$

April 24, 2016

38

EE2S31



Frequency response and poles/zeros

Express the complex-valued factors in polar form as

$$j\Omega - z_k = B_k(\Omega) e^{j\psi_k(\Omega)}, \quad j\Omega - p_k = A_k(\Omega) e^{j\theta_k(\Omega)}$$

Hence, we have

$$|H(j\Omega)| = \left| \frac{b_M}{a_N} \right| \frac{\prod_{k=1}^M B_k(\Omega)}{\prod_{k=1}^N A_k(\Omega)}$$

and

$$\angle H(j\Omega) = \angle \frac{b_M}{a_N} + \sum_{k=1}^M \psi_k(\Omega) - \sum_{k=1}^N \theta_k(\Omega)$$

April 24, 2016

39

EE2S31




Frequency selective filters

Low-pass filter: voltage across the inductor

Transfer function:

$$H(s) = \frac{s^2}{s^2 + s + 1} = \frac{s^2}{(s + p_1)(s + p_2)}$$

$$p_1 = p_2^* = \frac{1}{2}(1 + j\sqrt{3})$$



April 24, 2016

40


EE2S31



Frequency response and poles/zeros

Hence

- $|H(j0)| = 0$
- $\lim_{\Omega \pm \infty} |H(j\Omega)| = 1$
- $\angle H(j0^+) = \pi$
- $\angle H(j0^-) = -\pi$
- $\lim_{\Omega \pm \infty} \angle H(j\Omega) = 0$



April 24, 2016

41

EE2S31

