

Spectral Analysis of Finite-Length Signals

Richard Heusdens

April 28, 2016

1

EE2S31



Spectral Analysis

Recall that the spectrum of the discrete-time signal x is given by

$$X(\omega) = \sum_{n=-\infty}^{\infty} x(n)e^{-j\omega n}$$

where the signal x can be recovered from its spectrum by the inverse Fourier transform

$$x(n) = \frac{1}{2\pi} \int_0^{2\pi} X(\omega)e^{j\omega n} d\omega$$

In order to compute $X(\omega)$, we need infinite many data samples:

- implementation by a numeric computer or DSP not possible

April 28, 2016

2

EE2S31



Spectral Analysis

We will approximate the Fourier transform (FT) by using the *discrete Fourier transform (DFT)*, which works with finite-length data sequences

- 1) we start by investigating the relation between the FT and the DFT when the signal to be transformed has finite support, that is, has a finite number of non-zero elements
- 2) we then study the relation between the FT of a signal with infinite support and one in which the signal is first time-windowed (thus having finite support)


April 28, 2016

3

EE2S31

TU Delft

Spectral Analysis



April 28, 2016

4

EE2S31

TU Delft

Spectral Analysis

Step 1: DFT of y :

$$\begin{aligned} Y(\omega_k) &= \sum_{n=0}^{N-1} y(n) e^{-j\omega_k n} \\ &= \sum_{n=-\infty}^{\infty} x_w(n) e^{-j\omega_k n} \\ &= X_w(\omega_k), \quad \omega_k = \frac{2\pi}{N}k, \quad k = 0, \dots, N-1 \end{aligned}$$

Hence, $Y(\omega_k)$ is composed of samples of the frequency transform $X(\omega)$, where the samples are taken at frequencies $\omega_k = \frac{2\pi}{N}k$, $k = 0, \dots, N-1$.

April 28, 2016


5

EE2S31

TU Delft

Spectral Analysis

Example:



What is the effect of padding zeros to y , thereby making it of length $L \geq N$?

April 28, 2016

6

EE2S31

TU Delft

Spectral Analysis

Let $z(n) = x_w(n)$ for $n = 0, \dots, L-1$. We then have

$$\begin{aligned} Z(\omega_k) &= \sum_{n=0}^{L-1} z(n)e^{-j\omega_k n} \\ &= \sum_{n=-\infty}^{\infty} x_w(n)e^{-j\omega_k n} \\ &= X_w(\omega_k), \quad \omega_k = \frac{2\pi}{L}k, \quad k = 0, \dots, L-1 \end{aligned}$$

Zero-padding gives us more samples of the underlying frequency-continuous spectrum!

April 28, 2016


7

EE2S31

TU Delft

Spectral Analysis

Example:



April 28, 2016

8

EE2S31

TU Delft

Spectral Analysis

Step 2: DFT of x_w :

$$\begin{aligned} X_w(\omega) &= \sum_{n=-\infty}^{\infty} x_w(n) e^{-j\omega n} \\ &= \sum_{n=-\infty}^{\infty} (x \cdot w)(n) e^{-j\omega n} \\ &= (X * W)(\omega), \quad \omega \in [0, 2\pi) \end{aligned}$$

Hence, $X_w(\omega)$ is given by the convolution of the frequency transforms X and W evaluated at the frequency ω

April 28, 2016

9



EE2S31

TU Delft

Spectral Analysis

Example:

$$x(n) = \cos\left(\frac{\pi}{4}n\right), \quad n \in \mathbb{R} \quad \Longleftrightarrow \quad X(\omega) = \pi\delta\left(\omega - \frac{\pi}{4}\right) + \pi\delta\left(\omega + \frac{\pi}{4}\right)$$



April 28, 2016



10

EE2S31

TU Delft

Spectral Analysis

$$w(n) = \begin{cases} 1, & n = 0, \dots, N-1 \\ 0, & \text{otherwise} \end{cases} \iff |W(\omega)| = \frac{\sin(N\omega/2)}{\sin(\omega/2)}$$



April 28, 2016

11

EE2S31



TU Delft

Spectral Analysis

$$x_w(n) = \begin{cases} \cos\left(\frac{\pi}{4}n\right), & n = 0, \dots, N-1 \\ 0, & \text{otherwise} \end{cases} \iff |X_w(\omega)| = |(X * W)(\omega)|$$

where

$$(W * \delta_{\omega_0})(\omega) = \int_0^{2\pi} W(\nu) \delta(\omega - \nu - \omega_0) d\nu = W(\omega - \omega_0)$$



April 28, 2016

12


EE2S31

TU Delft


Window Functions

We can use other than rectangular windows that have a better frequency decay (less spectral leakage)

rectangular



Hanning



April 28, 2016

13

EE2S31

TU Delft

Fourier Transform

We would like to design windows having rapid descent of $W(f)$. Let $w \in L^1(\mathbb{R})$:

- If w is p times differentiable and all derivatives are in $L^1(\mathbb{R})$, then

$$w^{(p)}(t) \xleftrightarrow{\mathcal{F}} (j2\pi f)^p W(f)$$

- Applying the Riemann-Lebesgue lemma on $w^{(p)}$, we conclude

$$\lim_{f \rightarrow \pm\infty} f^p W(f) = 0$$

so that regularity of w translates in rapid descent of W .


April 28, 2016

14

EE2S31

TU Delft

Window Functions



April 28, 2016

15

EE2S31

TU Delft

Window Functions




Figure 10.2.3 Shapes of several window functions.

April 28, 2016

16

EE2S31

TU Delft

Window Functions




Figure 10.2.2 Frequency response for rectangular window of lengths (a) $M = 31$, (b) $M = 61$.

April 28, 2016

17

EE2S31

TU Delft

Window Functions




Figure 10.2.4 Frequency responses of Hanning window for (a) $M = 31$ and (b) $M = 61$.

April 28, 2016

18

EE2S31

TU Delft

Window Functions




Figure 10.2.5 Frequency responses for Hamming window for (a) $M = 31$ and (b) $M = 61$.

April 28, 2016

19

EE2S31

TU Delft

Window Functions




Figure 10.2.6 Frequency responses for Blackman window for (a) $M = 31$ and (b) $M = 61$.

April 28, 2016

20

EE2S31

TU Delft

Window Functions

How can we reduce the spectral smoothing?

Example: rectangular window

$$w(n) = \begin{cases} 1, & n = 0, \dots, N-1 \\ 0, & \text{otherwise} \end{cases} \iff |W(\omega)| = \frac{\sin(N\omega/2)}{\sin(\omega/2)}$$

Hence, we have zero-crossings at $\omega_k = \frac{2\pi}{N}k$, $k = 1, \dots, N-1$

\Rightarrow transition width of the main lobe is $\frac{4\pi}{N}$ ← reduced by increasing N !

April 28, 2016

21

EE2S31

 TU Delft

Window Functions

window type	width main lobe	peak side lobe (dB)
Rectangular	$\frac{4\pi}{N}$	-13
Bartlett	$\frac{8\pi}{N}$	-25
Hanning	$\frac{8\pi}{N}$	-31
Hamming	$\frac{8\pi}{N}$	-41
Blackman	$\frac{12\pi}{N}$	-57

April 28, 2016







22

EE2S31

 TU Delft

Window Functions

Example: $x(n) = w(n) (\cos(0.25\pi n) + \cos(0.35\pi n))$



April 28, 2016

23

EE2S31

TUDelft