



HUST

ĐẠI HỌC BÁCH KHOA HÀ NỘI
HANOI UNIVERSITY OF SCIENCE AND TECHNOLOGY

ONE LOVE. ONE FUTURE.



FUNDAMENTALS OF OPTIMIZATION



ĐẠI HỌC
BÁCH KHOA HÀ NỘI
HANOI UNIVERSITY
OF SCIENCE AND TECHNOLOGY

FUNDAMENTALS OF OPTIMIZATION

Week 2: Convex Optimization

ONE LOVE. ONE FUTURE.

1. Definitions


- Convex set
- Convex combination
- Convex function
- Exercises

2. Unconstrained Optimization


3. Constrained Optimization

Convex set


- A set of points $C \subseteq \mathbb{R}^n$ is **convex** if for every pair of points $x, y \in C$, any point on the line segment between x and y is also in C
- That is, if $x, y \in C$, then $tx + (1 - t)y \in C$ for all $t \in [0,1]$



Convex Set




Not a Convex Set



Examples of convex set

- **Line Segments:** $C = \{x + t(y - x) \mid t \in [0,1]\}$ for some $x, y \in \mathbb{R}^n$




- Lines, planes, hyperplanes, etc. also define convex sets

Examples of convex set

- **Half spaces:** $C = \{x \in \mathbb{R}^n \mid w^T x + b \leq 0\}$ for some $w \in \mathbb{R}^n$ and $b \in \mathbb{R}$
- **Balls of Radius ϵ :** $C = \{x \in \mathbb{R}^n \mid \|x\|_2 \leq \epsilon\}$ for some $\epsilon \geq 0 \in \mathbb{R}$


$$\|x\|_2 = \sqrt{x^T x} = \sqrt{\sum_{i=1}^n x_i^2}$$

- This is called the Euclidean norm or Euclidean distance because $\|x - y\|_2$ is equal to the length of the line segment between the points x and y



Convex combination

- We say that $y \in \mathbb{R}^n$ is a **convex combination** of the points $x^{(1)}, \dots, x^{(k)} \in \mathbb{R}^n$ if $y = \lambda_1 x^{(1)} + \dots + \lambda_k x^{(k)}$ for some choice of $\lambda_1, \dots, \lambda_k \in [0,1]$ such that $\lambda_1 + \dots + \lambda_k = 1$
- Let C be the set of all points y that can be obtained as a convex combination of the $x^{(1)}, \dots, x^{(k)}$
 - In the special case $k = 2$, C is just a line segment
 - C is a convex set called the **convex hull** of the $x^{(1)}, \dots, x^{(k)}$




Convex function

- A **function** $f: C \rightarrow \mathbb{R}$ is **convex** if C is a convex set and

$$f(tx + (1 - t)y) \leq tf(x) + (1 - t)f(y)$$

for all $x, y \in C$ and $t \in [0,1]$

- f is called **concave** if $-f$ is convex



Exercises

1. $S = \{(x, y) \mid 0 \leq x \leq a, 0 \leq y \leq b\}$ is a convex set ?
2. *Union* of two convex sets is a convex set ?
3. $S = \{x \in R^n \mid ||x - a|| \leq r\}$ is a convex set ?
4. $S = \{(x, y, z) \mid z \geq x^2 + y^2\}$ is a convex set ?

$S = \{(x, y) \mid 0 \leq x \leq a, 0 \leq y \leq b\}$ is a convex set ?

Let (x_1, y_1) and (x_2, y_2) be two arbitrary points in S . This means:

$$0 \leq x_1 \leq a, \quad 0 \leq y_1 \leq b, \quad 0 \leq x_2 \leq a, \quad 0 \leq y_2 \leq b.$$

Consider any convex combination of these points:

$$x_\lambda = \lambda x_1 + (1 - \lambda)x_2, \quad y_\lambda = \lambda y_1 + (1 - \lambda)y_2,$$

where $\lambda \in [0, 1]$.

- Since $x_1, x_2 \in [0, a]$, the weighted sum x_λ also satisfies $0 \leq x_\lambda \leq a$.
- Similarly, since $y_1, y_2 \in [0, b]$, the weighted sum y_λ satisfies $0 \leq y_\lambda \leq b$.

Thus, every point on the line segment between (x_1, y_1) and (x_2, y_2) remains in S , proving that S is convex.

Union of two convex sets is a convex set ?

Consider two convex sets:

$$S_1 = \{(x, y) \mid x \geq 0, y = 1\}$$

$$S_2 = \{(x, y) \mid x \leq 0, y = -1\}$$

- S_1 is a horizontal line at $y = 1$, which is convex.
- S_2 is a horizontal line at $y = -1$, which is also convex.

Now, take two points:

- $A = (1, 1) \in S_1$
- $B = (-1, -1) \in S_2$

The line segment joining A and B is given by:

$$(x_\lambda, y_\lambda) = \lambda(1, 1) + (1 - \lambda)(-1, -1) = (2\lambda - 1, 2\lambda - 1), \quad \lambda \in [0, 1].$$

For values of λ strictly between 0 and 1, the intermediate points **do not belong to** S_1 or S_2 . Hence, the union $S_1 \cup S_2$ is **not convex**.

- **Nonnegative weighted sums** of convex functions are convex, i.e., if $f_1: \mathbb{R}^n \rightarrow \mathbb{R}$ and $f_2: \mathbb{R}^n \rightarrow \mathbb{R}$ are convex functions and $c_1, c_2 \geq 0$, then
$$g(x) = c_1 f_1(x) + c_2 f_2(x)$$
is a convex function.
- **Pointwise maximum** of convex functions are convex, i.e., if $f_1: \mathbb{R}^n \rightarrow \mathbb{R}$ and $f_2: \mathbb{R}^n \rightarrow \mathbb{R}$ are convex functions, then
$$g(x) = \max(f_1(x), f_2(x))$$
is a convex function.

- A differentiable function $f: \mathbb{R}^n \rightarrow \mathbb{R}$ is convex on a convex set \mathcal{C} if and only if

$$f(x) \geq f(y) + \nabla f(y)^T (x - y)$$

for all $x, y \in \mathcal{C}$

- Which of the following functions are convex?
 - $\exp(x)$
 - $\exp(-x)$
 - $\log(x)$
 - $\sin(x)$
 - x^2
 - x^8
 - $\max(x, 0)$
 - \sqrt{x}
 - $|x|$

- Which of the following functions are convex?
 - $\exp(x)$
 - $\exp(-x)$
 - $\log(x)$
 - $\sin(x)$
 - x^2
 - x^8
 - $\max(x, 0)$
 - \sqrt{x}
 - $|x|$

1. Definitions

2. Unconstrained Optimization

- Introduction to unconstrained optimization
- Descent method
- Gradient descent method
- Newton method

3. Constrained Optimization

Unconstrained convex optimization


- Unconstrained, smooth convex optimization problem:

$$\min f(x)$$


- $f: R^n \rightarrow R$ is convex and twice differentiable
- ***dom* $f = R$** : no constraint
- Assumption: the problem is solvable with $f^* = \min_x f(x)$ and $x^* = \arg\min_x f(x)$
- To find x , solve equation $\nabla f(x^*) = 0$: **not easy to solve analytically**
- Iterative scheme is preferred: compute minimizing sequence $x^{(0)}, x^{(1)}, \dots$ s.t. $f(x^{(k)}) \rightarrow f(x^*)$ as $k \rightarrow \infty$
- The algorithm stops at some point $x(k)$ when the error is within acceptable tolerance: $f(x^{(k)}) - f^* \leq \varepsilon$

Local minimizer

- x^* is a local minimizer for $f: R^n \rightarrow R$ if $f(x^*) \leq f(x)$ for $\|x^* - x\| \leq \varepsilon$ ($\varepsilon > 0$ is a constant)
- x^* is a global minimizer for $f: R^n \rightarrow R$ if $f(x^*) \leq f(x)$ for all $x \in R^n$




$f(x) = e^x$ has no minimizer




$f(x) = -x + e^{-x}$ has no minimizer

Local minimizer

- x^* is a local minimizer for $f: R^n \rightarrow R$ if $f(x^*) \leq f(x)$ for $\|x^* - x\| \leq \varepsilon$ ($\varepsilon > 0$ is a constant)
- x^* is a global minimizer for $f: R^n \rightarrow R$ if $f(x^*) \leq f(x)$ for all $x \in R^n$




$f(x) = e^x + e^{-x} - 3x^2 + x$ has two local minimizers and one global minimizer




$f(x) = e^x + e^{-x} - 3x^2$ has two global minimizers

Local minimizer

- x^* is a local minimizer for $f: R^n \rightarrow R$ if $f(x^*) \leq f(x)$ for $||x^*-x|| \leq \varepsilon$ ($\varepsilon > 0$ is a constant)
- **Theorem** (Necessary condition for local minimum) If x^* is a local minimizer for $f: R^n \rightarrow R$, then $\nabla f(x^*) = 0$ (x^* is also called *stationary point* for f)



$f(x) = e^x + e^{-x} - 3x^2 + x$ has two local minimizers and one global minimizer



$f(x) = e^x + e^{-x} - 3x^2$ has two global minimizers

Example

- $f(x, y) = x^2 + y^2 - 2xy + x$
- $\nabla f(x, y) = \begin{pmatrix} 2x - 2y + 1 \\ 2y - 2x \end{pmatrix} = 0$ has no solution

→ there is no minimizer of $f(x, y)$

Local minimizer

- **Theorem** (Necessary condition for local minimum) If x^* is a local minimizer for $f: R^n \rightarrow R$, then $\nabla f(x^*) = 0$ (x^* is also called *stationary point* for f)
- **Theorem** (Sufficient condition for a local minimum) Assume x^* is a stationary point and that $\nabla^2 f(x^*)$ is positive definite, then x^* is a local minimizer

- $\nabla^2 f(x) =$
$$\begin{pmatrix} \frac{\partial^2 f(x)}{\partial x_1 \partial x_1} & \frac{\partial^2 f(x)}{\partial x_1 \partial x_2} & \cdots & \frac{\partial^2 f(x)}{\partial x_1 \partial x_n} \\ \frac{\partial^2 f(x)}{\partial x_2 \partial x_1} & \frac{\partial^2 f(x)}{\partial x_2 \partial x_2} & \cdots & \frac{\partial^2 f(x)}{\partial x_2 \partial x_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f(x)}{\partial x_n \partial x_1} & \frac{\partial^2 f(x)}{\partial x_n \partial x_2} & \cdots & \frac{\partial^2 f(x)}{\partial x_n \partial x_n} \end{pmatrix}$$

- Matrix $A_{n \times n}$ is called positive definite if

$$A^i = \begin{pmatrix} a_{1,1} & a_{1,2} & \dots & a_{1,i} \\ a_{2,1} & a_{2,2} & \dots & a_{2,i} \\ \dots & \dots & \dots & \dots \\ a_{i,1} & \dots & a_{i,2} & \dots & a_{i,i} \end{pmatrix}, \det(A^i) > 0, i = 1, \dots, n$$

Examples

Example $f(x,y) = e^{x^2+y^2}$

$$\nabla f(x) = \begin{pmatrix} 2xe^{x^2+y^2} \\ 2ye^{x^2+y^2} \end{pmatrix} = 0 \text{ has unique solution } x^* = (0,0)$$

$$\nabla^2 f(x^*) = \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix} > 0 \rightarrow (0,0) \text{ is a minimizer of } f$$

Examples

- **Example** $f(x,y) = x^2 + y^2 - 2xy - x$

$$\nabla f(x) = \begin{pmatrix} -2x + 2y + 1 \\ -2x - 2y \end{pmatrix} = 0$$

has unique solution $x^* = (-1/4, 1/4)$

$$\nabla^2 f(x^*) = \begin{pmatrix} -2 & 2 \\ -2 & -2 \end{pmatrix} \text{ is not positive definite}$$

→ cannot conclude x^*

Descent method

Determine starting point $x^{(0)} \in \mathbb{R}^n$;

$k \leftarrow 0$;

While (stop condition not reach){

 Determine a search direction $p_k \in \mathbb{R}^n$;

 Determine a step size $\alpha_k > 0$ s.t. $f(x^{(k)} + \alpha_k p_k) < f(x^{(k)})$;

$x^{(k+1)} \leftarrow x^{(k)} + \alpha_k p_k$;

$k \leftarrow k+1$;

}

Stop condition may be

- $\|\nabla f(x^k)\| \leq \varepsilon$
- $\|x^{k+1} - x^k\| \leq \varepsilon$
- $k > K$ (maximum number of iterations)

Gradient descent method

- Gradient descent schema

$$x^{(k)} = x^{(k-1)} - \alpha_k \nabla f(x^{(k-1)})$$

```
init  $x^{(0)}$ ;  
 $k = 1$ ;  
while stop condition not reach {  
    specify constant  $\alpha_k$ ;  
     $x^{(k)} = x^{(k-1)} - \alpha_k \nabla f(x^{(k-1)})$ ;  
     $k = k + 1$ ;  
}
```

- α_k might be specified in such a way that $f(x^{(k-1)} - \alpha_k \nabla f(x^{(k-1)}))$ is minimized: $\frac{\partial f}{\partial \alpha_k} = 0$

Minimize $f(x) = x^4 - 2x^3 - 64x^2 + 2x + 63$

```
# compute the function value
def f(x):
    return x**4 - 2*x**3 - 64*x**2 + 2*x + 63

# compute the derivative value
def grad(x):
    return 4*x**3 - 6*x**2 - 128 * x + 2

# Gradient descent algorithm with given alpha and initial point
def myGD(alpha, x0):
    x = [x0]
    # loop to evaluate a series of candidate
    for it in range(100000):
        # x[k+1] = x[k] - alpha * f'(x[k])
        x_new = x[-1] - alpha*grad(x[-1])

        # check stop condition (f'(x[k+1]) <= epsilon)
        if abs(grad(x_new)) < 1e-3:
            break

        # append x[k+1] into list
        x.append(x_new)

    # return a list of evaluated candidates and the iteration at which the algorithm stops
    return (x, it)
```

Minimize $f(x) = x^4 + 3x^2 - 10x + 4$

```
def grad(x):  
    return 4*x**3+ 6*x - 10  
  
def f(x):  
    return x**4 + 3* x**2 - 10 * x + 4  
  
def myGD(alpha, x0):  
    x = [x0]  
    for it in range(1000):  
        x_new = x[-1] - alpha*grad(x[-1])  
        # if abs(grad(x_new)) < 1e-3:  
        #     break  
  
        if(abs(x[-1] - x_new) < 1e-3):  
            break  
  
        x.append(x_new)  
    return (x, it)
```

Minimize $f(x) = x^2 + 5\sin(x)$

```
def grad(x):  
    return 2*x+ 5*np.cos(x)  
  
def f(x):  
    return x**2 + 5*np.sin(x)  
  
def myGD(delta, x0):  
    x = [x0]  
    for it in range(100):  
        x_new = x[-1] - delta*grad(x[-1])  
        if abs(grad(x_new)) < 1e-3:  
            break  
        x.append(x_new)  
    return (x, it)
```

Minimize $f(x, y) = x^2 + y^2 + xy - x - y$

```
def grad(x, y):  
    return (2*x + y - 1, 2*y + x - 1)  
  
def f(x, y):  
    return x**2 + y**2 + x*y - x - y  
  
def myGD(delta, x0, y0):  
    X = [(x0, y0)]  
    for it in range(1000):  
        x_new = X[-1][0] - delta*grad(X[-1][0], X[-1][1])[0]  
        y_new = X[-1][1] - delta*grad(X[-1][0], X[-1][1])[1]  
        if abs(grad(x_new, y_new)[0]) < 1e-6 and abs(grad(x_new, y_new)[1]) < 1e-6:  
            break  
        X.append((x_new, y_new))  
    return (X, it)
```


Minimize $f(x) = x_1^2 + x_2^2 + x_3^2 - x_1x_2 - x_2x_3 + x_1 + x_3$

- α_k might be specified in such a way that $f(x^{(k-1)} - \alpha_k \nabla f(x^{(k-1)}))$ is minimized: $\frac{\partial f}{\partial \alpha_k} = 0$

```
def grad(x1, x2, x3):  
    return [2*x1 + 1 - x2, -x1 + 2*x2 - x3, -x2 + 2*x3 + 1]  
  
def f(x1, x2, x3):  
    return x1**2 + x2**2 + x3**2 - x1*x2 - x2*x3 + x1 + x3  
  
def myGD(v1, v2, v3):  
    x1 = v1  
    x2 = v2  
    x3 = v3  
    for it in range(1000):  
        print(f(x1, x2, x3))  
        [D1,D2,D3] = grad(x1,x2,x3)  
        A = 2*x1*D1 + 2*x2*D2 + 2*x3*D3 - x1*D2 - x2*D1 - x2*D3 -x3*D2 + D1 + D3  
        B = 2*D1*D1 + 2*D2*D2 + 2*D3*D3 -2*D1*D2 - 2*D2*D3  
        if B == 0:  
            break  
        alpha = A/B  
  
        x1 = x1 - alpha*D1  
        x2 = x2 - alpha*D2  
        x3 = x3 - alpha*D3  
  
        val = grad(x1, x2, x3)  
        if (val[0]**2 + val[1]**2 + val[2]**2) < 1e-6:  
            break  
  
        X.append([x1, x2, x3])  
    return (X, it)
```

Newton method

- Second-order Taylor approximation g of f at x is

$$f(x+h) \approx g(x+h) = f(x) + h \nabla f(x) + \frac{1}{2}h^2 \nabla^2 f(x)$$

- Which is a convex quadratic function of h
- $g(x+h)$ is minimized when $\frac{\partial g}{\partial h} = 0 \rightarrow h = -\nabla^2 f(x)^{-1} \nabla f(x)$

```
Generate  $x^{(0)}$ ; // starting point
 $k = 0$ ;
while stop condition not reach{
     $x^{(k+1)} \leftarrow x^{(k)} - \nabla^2 f(x^{(k)})^{-1} \nabla f(x^{(k)})$ ;
     $k = k + 1$ ;
}
```

Minimize $f(x) = x_1^2 + x_2^2 + x_3^2 - x_1x_2 - x_2x_3 + x_1 + x_3$

```
import numpy as np

def newton(f,df,Hf,x0):
    x = x0
    for i in range(10):
        iH = np.linalg.inv(Hf(x))
        D = np.array(df(x)).T #transpose matrix: convert from list to
                               #column vector

        print('df = ',D)
        y = iH.dot(D) #multiply two matrices
        if np.linalg.norm(y) == 0:
            break
        x = x - y
        print('Step ',i,': ',x,' f = ',f(x))
```

Minimize $f(x) = x_1^2 + x_2^2 + x_3^2 - x_1x_2 - x_2x_3 + x_1 + x_3$

```
def main():
    print('main start....')
    f = lambda x: x[0] ** 2 + x[1] ** 2 + x[2] ** 2 - x[0] * x[1] - x[1] *
                x[2] + x[0] + x[2] # function f to be minimized
    df = lambda x: [2 * x[0] + 1 - x[1], -x[0] + 2 * x[1] - x[2], -x[1] + 2
                * x[2] + 1] # gradient
    Hf = lambda x: [[2,-1,0],[-1,2,-1],[0,-1,2]]# Hessian
    x0 = np.array([0,0,0]).T
    newton(f,df,Hf,x0)

if __name__ == '__main__':
    main()
```

1. Definitions

2. Unconstrained Optimization

3. Constrained Optimization

- General constrained optimization problem
- Lagrange multiplier method

General constrained optimization problem

- Optimization problem in the standard form

$$\begin{array}{ll} (P) & \text{minimize } f(x) \\ \text{s.t.} & g_i(x) = 0, \quad \forall i \in \{1, 2, \dots, m\} \\ & x \in X \subseteq R^n \end{array}$$

with $x \in R^n$, and assume $D = (\cap_{i=1}^m \text{dom } g_i)$ is not empty.

- Denote f^* the optimal value of $f(x)$
- If f, g_i ($i = 1, 2, \dots, m$) are convex functions.

Lagrangian function

- Optimization problem in the standard form

$$\begin{array}{ll} (P) & \text{minimize } f(x) \\ \text{s.t.} & g_i(x) = 0, \quad \forall i \in \{1, 2, \dots, m\} \\ & x \in X \subseteq \mathbb{R}^n \end{array}$$

- Lagrangian function of the above problem is defined as follows, $L: \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}$

$$L(x, \lambda) = f(x) + \sum_{i=1}^m \lambda_i \times g_i(x)$$

Lagrangian function - properties

- Optimization problem in the standard form

$$\begin{aligned} (P) \quad & \text{minimize } f(x) \\ \text{s.t.} \quad & g_i(x) = 0, \quad \forall i \in \{1, 2, \dots, m\} \\ & x \in X \subseteq R^n \end{aligned}$$

- Lagrangian function of the above problem is defined as follows, $L: R^n \times R^m \rightarrow R$

$$L(x, \lambda) = f(x) + \sum_{i=1}^m \lambda_i \times g_i(x)$$

- **Theorem:** The optimal value of the optimization problem is the following property: $\nabla_x L(x, \lambda) = 0$

Lagrangian multiplier method

- **The method of Lagrange multipliers** is a technique in mathematics to find the local maxima or minima of a function $f(x_1, x_2, \dots, x_n)$ subject to constraints $g_i(x_1, \dots, x_n) = 0, \forall i \in \{1, \dots, m\}$.

- **Method:**

- Step 1: Solving the following system

$$\begin{aligned}\nabla f(x_1, \dots, x_n) &= \sum_{i=1}^m \lambda_i \nabla g_i(x_1, \dots, x_n) \\ g_1(x_1, \dots, x_n) &= 0 \\ &\dots \\ g_m(x_1, \dots, x_n) &= 0\end{aligned}$$

- Step 2: we get particular values of x_1, x_2, \dots, x_n , which can be plugged in $f(x_1, x_2, \dots, x_n)$ to get the extremum value if it exists.

Example 1:

- **Problem:** Find the maximum and minimum of $f(x, y) = 5x - 3y$ subject to the constraint $x^2 + y^2 = 136$.

- **Solution:**

- Region of possible solutions lies on a disk of radius $\sqrt{136}$ which is a closed and bounded region, $-\sqrt{136} \leq x, y \leq \sqrt{136}$
- The Lagrangian function $L(x, y, \lambda) = 5x - 3y - \lambda(x^2 + y^2 - 136)$
- As the result of the theorem, we have

$$\begin{aligned}5 &= 2\lambda x \\ -3 &= 2\lambda y \\ x^2 + y^2 &= 136\end{aligned}$$

- Notice that, we can't have $\lambda = 0$ since that would not satisfy the first two equations. So, since we know that $\lambda \neq 0$ we can solve the first two equations for x and y respectively. This gives,

$$x = \frac{5}{2\lambda} \qquad y = -\frac{3}{2\lambda}$$

Example 1

- **Problem:** Find the maximum and minimum of $f(x, y) = 5x - 3y$ subject to the constraint $x^2 + y^2 = 136$.

- **Solution:**

- Plugging these into the constraint gives

$$\frac{25}{4\lambda^2} + \frac{9}{4\lambda^2} = \frac{17}{2\lambda^2} = 136$$

- We can solve this for λ

$$\lambda^2 = \frac{1}{16} \Rightarrow \lambda = \pm \frac{1}{4}$$

- Now, that we know λ we can find the points that will be potential maximums and/or minimums.

- If $\lambda = -\frac{1}{4}$ we get $x = -10, y = 6$

- if $\lambda = \frac{1}{4}$ we get $x = 10, y = -6$

- So,

$$f(-10, 6) = -68$$

Minimum at $(-10, 6)$

$$f(10, -6) = 68$$

Maximum at $(10, -6)$

Example 2:

- **Problem:**

- Objective function: *maximize* $f(x, y) = xy$
- Constraint: $g(x, y) = 10x + 20y - 400 = 0$

- **Solution**

- Form the Lagrange function:

$$L(x, y, \mu) = f(x, y) - \mu(g(x, y))$$
$$L(x, y, \mu) = xy - \mu(10x + 20y - 400)$$

- Set each first order partial derivative equal to zero:

$$\frac{\partial L}{\partial x} = y - 10\mu = 0$$

$$\frac{\partial L}{\partial y} = x - 20\mu = 0$$

$$\frac{\partial L}{\partial \mu} = -(10x + 20y - 400) = 0$$

Example 2:

- **Problem:**

- Objective function: *maximize* $u(x, y) = xy$
- Constraint: $g(x, y) = 10x + 20y - 400 = 0$

- **Solution:**

- Set each first order partial derivative equal to zero:

$$\frac{\partial L}{\partial x} = y - 10\mu = 0$$

$$\frac{\partial L}{\partial y} = x - 20\mu = 0$$

$$\frac{\partial L}{\partial \mu} = -(10x + 20y - 400) = 0$$

- So,

$$10x + 20y = 400$$

$$40y = 400$$

$$y = 10$$

$$x = 2y = 20$$

Example 3:

- **Problem:**

- Objective function: *maximize* $f(x, y) = x + y$
- Constraint: $g(x, y) = x^2 + y^2 - 2 = 0$

- **Solution:**

- Form the Lagrange function:

$$\mathcal{L}(x, y, \lambda) = x + y + \lambda(x^2 + y^2 - 2)$$

- Set each first order partial derivative equal to zero:
 - $\frac{\partial L}{\partial x} = 1 + 2\lambda x = 0$
 - $\frac{\partial L}{\partial y} = 1 + 2\lambda y = 0$
 - $\frac{\partial L}{\partial \lambda} = x^2 + y^2 - 2 = 0$

Example 3

- **Problem:**

- Objective function: *maximize* $f(x, y) = x + y$
- Constraint: $g(x, y) = x^2 + y^2 - 2 = 0$

- **Solution:**

- Set each first order partial derivative equal to zero:
 - $\frac{\partial L}{\partial x} = 1 + 2\lambda x = 0$
 - $\frac{\partial L}{\partial y} = 1 + 2\lambda y = 0$
 - $\frac{\partial L}{\partial \lambda} = x^2 + y^2 - 2 = 0$
- We have , $(x, y) \in \{(1,1), (-1, -1)\}$
- So, $(x, y) = (1,1)$

A large graphic on the left side of the slide. It features a dark blue background with a circular pattern of red dots of varying sizes, creating a sense of depth and movement. The word "HUST" is centered within this graphic in a bold, white, sans-serif font.

HUST

THANK YOU !