Asparagine

Table of Contents

summary

History

Biochemical Role

Protein Synthesis

Metabolic Functions

Cancer Cell Metabolism

Dietary Sources

Biosynthesis

Health Implications

Nutritional Significance

Deficiency and Disorders

Adverse Effects

Cancer Implications

Applications

Role in Cancer Treatment

Dietary Interventions

Immunotherapy and Metabolic Fitness

Future Directions

Check https://storm.genie.stanford.edu/article/1146852 for more details

Stanford University Open Virtual Assistant Lab

The generated report can make mistakes.

Please consider checking important information.

The generated content does not represent the developer's viewpoint.

summary

Asparagine is a non-essential amino acid that plays a pivotal role in various biological processes, including protein synthesis, metabolism, and cellular communication. Initially isolated from asparagus in 1833 by French chemist Anselme Payen, asparagine has since garnered significant attention due to its biochemical importance and implications for human health. [1][2] As a key component of proteins, asparagine

contributes to structural integrity and function, and it is involved in critical metabolic pathways that support cell proliferation and differentiation.[3][4]

The amino acid is particularly notable for its association with specific health conditions, most notably Asparagine Synthetase Deficiency (ASNSD), a rare genetic disorder that manifests severe neurological symptoms and developmental delays shortly after birth. [2][5] Additionally, asparagine's role in cancer cell metabolism has sparked interest in its potential as a therapeutic target, especially for leukemias and other malignancies that exhibit heightened dependence on this amino acid. [6][7] Recent studies suggest that dietary interventions limiting asparagine intake may improve treatment outcomes for various cancers, although further research is necessary to establish effective strategies. [6][7]

Despite its essential functions, asparagine levels can also pose health risks; excess concentrations may lead to adverse effects in individuals with compromised kidney or liver function, while imbalances have been linked to immune system stress.[8][9] As research continues to unfold, the complexities of asparagine's roles in both health and disease highlight the need for a nuanced understanding of this amino acid, particularly regarding its dietary sources and implications for clinical practice.[10][11]

History

Asparagine, an amino acid significant for various biological processes, has a rich history of discovery and research. The initial identification of asparagine dates back to the early 19th century, with notable contributions from chemists such as Robiquet and Liebig. In 1833, Robiquet is credited with isolating asparagine from asparagus, a discovery that was later expanded upon by Liebig, who studied its chemical composition and properties[1][2].

Throughout the 19th and early 20th centuries, the understanding of asparagine continued to evolve, with numerous studies focusing on its biochemical roles and potential health implications. The first synthesis of asparagine was achieved in the 1870s, further solidifying its importance in both academic and medical chemistry [7].

In modern research, asparagine's role in metabolism has been critically examined, particularly in the context of certain genetic disorders such as Asparagine Synthetase Deficiency (ASNSD), which is characterized by an autosomal recessive inheritance pattern. This condition demonstrates the essential role of asparagine in human health and development, leading to ongoing investigations into treatment and management strategies[2][12].

Biochemical Role

Asparagine is a non-essential amino acid that plays a critical role in various metabolic processes within the human body. Its primary significance lies in its function as a building block for proteins and its involvement in several key biochemical pathways, including protein synthesis, N-glycosylation, nucleotide biosynthesis, and ammonium metabolism[3][4].

Protein Synthesis

Asparagine is classified as a proteinogenic amino acid with a polar side chain, which facilitates the formation of hydrogen bonds and increases the solubility of proteins in aqueous environments. It contributes to the structural integrity of proteins through the formation of asparagine-linked glycoproteins, which are essential for cellular communication, adhesion, and immune response [4]. The amino acid's content in proteins is approximately 4%, and it participates in the formation of specific structural motifs that are crucial for proper protein folding [13].

Metabolic Functions

In addition to its role in protein synthesis, asparagine is involved in several metabolic pathways. It is essential for the biosynthesis of polyamines, which are important regulators of cell proliferation and differentiation[13]. The conversion of oxaloacetate to aspartate and asparagine is crucial in metabolic pathways, particularly when the activity of citrate synthase is inhibited. This conversion can help rescue cells from glutamine-induced apoptosis, demonstrating the amino acid's protective metabolic role[6].

Cancer Cell Metabolism

Asparagine also plays a significant role in cancer metabolism, where its levels and metabolism are often altered to support the high demands of rapidly proliferating cancer cells. The interplay between asparagine and glutamine metabolism has been a focal point of research, particularly regarding how asparagine can sustain protein synthesis and cell survival when glutamine is scarce[14]. Moreover, asparagine promotes glycolysis in adipocytes, which is linked to thermogenesis and overall metabolic regulation[6].

Dietary Sources

Asparagine is an amino acid that is widely distributed in various dietary sources, both from animal and plant origins. Animal sources of asparagine include dairy products, such as cheese and whey, as well as meats like beef, poultry, and fish[1][15][16]. Eggs and seafood are also significant contributors to dietary asparagine intake[17][18].

On the plant side, asparagine can be found in numerous foods, including legumes, nuts, seeds, whole grains, and certain vegetables like asparagus and potatoes[17-][15][18][19]. Additionally, plant-based sources such as soy products and seaweed (e.g., spirulina) are noted for their asparagine content[1][20]. The consumption of these foods plays a crucial role in providing the necessary asparagine to support various biological functions, including protein synthesis and nitrogen transport[8][21].

Furthermore, dietary patterns can influence asparagine levels in the body. For instance, vegan and vegetarian diets, which exclude animal products, may lead to variations in asparagine metabolism and availability, as evidenced by increased levels of intestinal ASNase in individuals adhering to these diets[10]. Conversely,

imbalanced dietary amino acid compositions or protein restrictions have been shown to activate the ASNS gene, which is responsible for asparagine synthesis, in certain contexts[10][22].

Despite the availability of asparagine in these foods, challenges remain in accurately assessing its levels due to the limitations of current nutrient databases and common dietary assessment methods[10][22]. This underscores the need for more comprehensive research to quantify asparagine in various foods and better understand its implications for health, particularly concerning colorectal cancer (CRC) risk and dietary intake assessments[10].

Biosynthesis

Asparagine is synthesized in the body primarily through the enzymatic conversion of aspartate, utilizing either ammonia or glutamine as nitrogen sources. The key enzyme involved in this process is asparagine synthetase (AS), which is responsible for catalyzing the formation of asparagine from aspartate and either ammonia or glutamine[23][24]. The mechanism of AS involves two active sites; the N-terminal active site facilitates glutamine hydrolysis to produce glutamate and ammonia, while the C-terminal site activates the side-chain carboxylate of aspartate, resulting in the formation of an electrophilic intermediate that ultimately leads to the synthesis of asparagine[23].

Asparagine biosynthesis plays a crucial role in various metabolic pathways, including protein synthesis, N-glycosylation, nucleotide biosynthesis, and ammonium metabolism[3]. It is particularly important in maintaining the nitrogen balance within cells and in the production of amino acids essential for cellular functions. Under conditions of nutrient deprivation, such as fasting, the utilization of asparagine and its precursors allows for the conservation of energy and the support of gluconeogenesis, facilitating the production of glucose from non-carbohydrate sources[25].

In addition to its metabolic roles, asparagine is also involved in cellular stress responses. During nutrient scarcity, uncharged tRNA can activate the general control nonderepressible 2 (GCN2) pathway, leading to the phosphorylation of eukaryotic translation initiation factor 2 subunit ±e(IF2±)which subsequently inhibits global protein synthesis and helps preserve amino acid levels[26]. This integrated stress response is vital for cell survival under challenging conditions.

Furthermore, asparagine is integral to the formation of amino sugars through glycosylation processes, contributing to the synthesis of glycoproteins and glycolipids that are essential for cellular communication and structure[26].

Health Implications

Asparagine is an amino acid that plays a significant role in various physiological processes within the body, and its implications for health can be both beneficial and detrimental, particularly in individuals with specific health conditions.

Nutritional Significance

Asparagine is crucial for normal bodily function, contributing to neurotransmission, emotional regulation, and memory support. [23][27] It helps facilitate communication between nerves, which can be particularly important for individuals with nerve-related disorders. [27] The recommended daily intake of asparagine varies, with average consumption levels reported to be approximately 50 g/day for preschool children and up to 110 g/day for adult males. [28] This dietary amino acid is generally obtained from a variety of food sources, including meat, dairy, and certain vegetables.

Deficiency and Disorders

Asparagine deficiency can lead to significant health issues. Notably, it has been linked to conditions such as asparagine synthetase deficiency, a severe neurological disorder that manifests shortly after birth. Symptoms include microcephaly, severe developmental delays, intractable seizures, and muscle stiffness, ultimately leading to paralysis.[5][2][29] Individuals with this condition often experience a decline in cognitive and motor skills over time and typically do not survive past childhood.[2] Furthermore, increased prevalence of asparagine deficiencies has been observed in clinical manifestations like fatigue and immune system stress.[30]

Adverse Effects

On the other hand, excessive asparagine levels or imbalances can result in adverse effects. Individuals with kidney or liver disorders may experience a range of symptoms, from mild gastrointestinal discomfort and headaches to severe reactions such as anaphylaxis.[8][9] It has been noted that imbalances in asparagine levels can also influence biomarkers associated with liver function and overall metabolic health.[9]

Cancer Implications

Recent research has explored the relationship between asparagine and cancer, particularly colorectal cancer (CRC). Asparagine levels can be influenced by dietary intake, with red meat consumption correlating to higher serum levels of asparagine. However, the complex interplay between asparagine and other cancer-promoting compounds found in red meat complicates the understanding of its role in CRC risk.[10][11] Moreover, therapies targeting asparagine metabolism, such as the combination of metformin and asparaginase, have shown potential in impairing tumor growth in cancer cells.[6]

Applications

Role in Cancer Treatment

Asparagine has garnered attention for its potential applications in cancer therapy, particularly in the context of metabolic interventions. Research indicates that cancer cells often exhibit a dependency on asparagine for growth and proliferation, making it a target for therapeutic strategies. Notably, asparagine is critical for certain types of leukemia, as cancer cells have high demands for this amino acid that cannot be sufficiently met through endogenous synthesis alone. This sensitivity has led to the successful use of the enzyme L-asparaginase, which depletes asparagine levels in patients with acute lymphoblastic leukemia, highlighting the amino acid's importance in cancer metabolism [6][7].

Dietary Interventions

Emerging studies suggest that dietary asparagine restriction may hold promise as an adjunct treatment for various cancers, including breast, kidney, and head and neck cancers. By potentially limiting the availability of asparagine, researchers aim to hinder cancer cell growth and spread. Future clinical applications could involve implementing specific dietary modifications in conjunction with traditional cancer therapies, such as chemotherapy, to improve patient outcomes [6][7]. However, it is essential for patients to consult healthcare providers before making any dietary changes during treatment, as current evidence regarding the efficacy of food restrictions in cancer management remains inconclusive [7].

Immunotherapy and Metabolic Fitness

Recent findings have also linked asparagine to immunological responses in cancer treatment. Asparagine is believed to play a role in the activation of CD8+ T lymphocytes, which are critical for anti-tumor immunity. In this context, restricting asparagine can enhance the metabolic fitness and functionality of these immune cells, potentially leading to improved anti-tumor responses [6]. Thus, asparagine not only serves as a vital nutrient for cancer cells but also influences immune responses, indicating its dual role in cancer biology and treatment strategies.

Future Directions

The exploration of asparagine in cancer therapies is ongoing, with researchers looking to translate laboratory findings into clinical settings. Future studies will focus on identifying specific patient populations that might benefit from asparagine-targeted treatments and understanding the mechanisms by which asparagine influences both tumor growth and immune response [7]. The potential repurposing of existing therapies, such as L-asparaginase, for broader cancer treatment applications underscores the need for continued research in this area [7].

References

[1]: Asparagine - Wikipedia

[2]: Asparagine synthetase deficiency: MedlinePlus Genetics

- [3]: Possible link between breast cancer and asparagine
- [4]: Asparagine Synthetase in Cancer: Beyond Acute Lymphoblastic ...
- [5]: Asparagine an overview | ScienceDirect Topics
- [6]: What is Asparagine used for? Patsnap Synapse
- [7]: Aspartic Acid in Health and Disease MDPI
- [8]: Metabolism of asparagine in the physiological state and cancer
- [9]: Asparagine sustains cellular proliferation and c Myc expression in ...
- [10]: Disparaging Asparagine Cedars-Sinai
- [11]: Diet may influence the spread of a deadly type of breast cancer ...
- [12]: Foods rich in asparagine Botanical-online
- [13]: Fast Take: This Is No Time to Disparage Asparagus Food Insight
- [14]: Diet & the Spread of Breast Cancer Oncology Times
- [15]: Asparagine an overview | ScienceDirect Topics
- [16]: What are the side effects of Asparagine? Patsnap Synapse
- [17]: Asparagine Metabolic Pathways in Arabidopsis PubMed
- [18]: Asparagine, colorectal cancer, and the role of sex, genes, microbes ...
- [19]: A whole food, plant-based diet reduces amino acid levels in patients ...
- [20]: Harnessing Asparagine: The Amino Acid Revolutionizing Nutrac
- [21]: Asparagine synthetase Wikipedia
- [22]: Aspartate and asparagine metabolism Reactome
- [23]: Asparagine: A Key Amino Acid in Human Health
- [24]: Asparagine Getfish Seafood
- [25]: Protein and Amino Acids Recommended Dietary Allowances NCBI
- [26]: Asparagine Synthetase Deficiency
- [27]: Asparagine Synthetase Deficiency Metabolic Support UK
- [28]: [PDF] Asparagine
- [29]: High dietary protein intake (including amino acid intake) and acute ...
- [30]: Asparagine synthetase: Function, structure, and role in disease PMC