

Thionine

Table of Contents

summary

Chemical Structure

Properties

Synthesis

Applications

- Biomedical Applications

 - Peptide-Self-Assembled Monolayers (SAMs)

- Electrochemical Sensors

 - Staining and Visualization

- Environmental Monitoring

Safety and Toxicity

Related Compounds

- Thiol Derivatives

- Dye Characteristics and Classifications

- Structural Similarities with Other Compounds

Check <https://storm.genie.stanford.edu/article/1166378> for more details

Stanford University Open Virtual Assistant Lab

The generated report can make mistakes.

Please consider checking important information.

The generated content does not represent the developer's viewpoint.

summary

Thionine is a synthetic metachromatic dye widely recognized for its applications in histology, biosensing, and environmental monitoring. With the chemical formula $C_{12}H_{10}ClN_3S$, it exhibits unique staining properties that allow it to change color when bound to various tissue components, facilitating the visualization of cellular structures in biological research.^{[1][2]} Thionine's structural complexity, including functional groups such as chlorine and sulfur, contributes to its diverse functionalities, making it a notable compound in both biomedical and environmental fields.

The dye has gained prominence in the development of electrochemical biosensors, where its ability to interact with biomolecules enables the detection of disease

markers such as glucose, proteins, and DNA with high sensitivity.[\[3\]\[4\]](#) Its incorporation into peptide-self-assembled monolayers enhances biosensing capabilities, allowing for improved cell adhesion and target analyte binding, which is crucial for point-of-care diagnostics.[\[3\]](#) Additionally, thionine is utilized in histological staining, providing researchers with a reliable method for studying tissue architecture and cellular interactions under light microscopy.[\[5\]](#)

Despite its widespread use, thionine is not without controversy, particularly concerning its safety and potential toxicity. Proper handling guidelines are essential to mitigate risks associated with exposure, as inhalation can lead to respiratory irritation and skin or eye contact may cause discomfort.[\[2\]\[6\]](#) The compound's environmental implications have also been examined, as its redox properties facilitate the detection of pollutants in water quality assessments, highlighting its versatility across multiple domains.[\[4\]](#)

In summary, thionine serves as a crucial tool in both research and practical applications, with ongoing studies exploring its effectiveness and safety in various fields. Its ability to function as a metachromatic dye, combined with its role in advanced biosensing technologies, positions it as a significant compound in contemporary scientific inquiry.[\[1\]\[3\]\[4\]](#)

Chemical Structure

Thionine is a metachromatic dye characterized by the molecular formula $C_{12}H_{10}ClN_3S$ and a molar mass of 263.743 g/mol[\[1\]\[2\]](#). Its chemical structure includes multiple functional groups, which contribute to its staining properties. In particular, Thionine contains chlorine and sulfur atoms, which are integral to its functionality as a dye[\[7\]](#).

The more complex variant known as Thionine blue has the molecular formula $C_{18}H_{22}Cl_3N_3SZn$, with a molecular weight of 484.2 g/mol[\[7\]](#). This version incorporates a zinc ion, enhancing its stability and reactivity in various chemical environments[\[7\]](#).

Both forms of Thionine exhibit metachromasia, which refers to the ability of the dye to change color when bound to tissue or cellular components. This property allows for differential staining in histological applications[\[8\]\[9\]](#). Thionine's color shifts can vary from blue to green or purple depending on the specific biomolecules it interacts with, making it a versatile agent in biological staining techniques[\[5\]](#).

Properties

Thionine, a synthetic dye, is characterized by its complex molecular structure, represented by the chemical formula $C_{12}H_{10}N_3S^+$ [\[7\]](#). It possesses a molar mass of approximately 228.29 g·mol⁻¹, which indicates its size and weight within the context of chemical compounds[\[10\]](#). Thionine has been classified based on its structural and chemical properties, making it applicable in various fields, including biology and materials science.

The behavior of thionine and its interactions with surfaces can be studied through techniques such as ellipsometry and surface plasmon resonance (SPR). These methods rely on changes in the polarization state of light after reflection from surfaces that have been treated with thionine or related compounds[3]. Specifically, the SPR technique is notable for quantifying molecular interactions in real-time by detecting alterations in the refractive index of a metal thin film, typically gold (Au), that has been modified to include thionine[11].

In practical applications, thionine's binding affinity and structural properties make it a suitable candidate for use in self-assembled monolayers (SAMs) on gold surfaces, where its interactions can influence the overall properties of the surface, such as wettability and adsorption characteristics[11][12]. The ability to form SAMs with thionine can be utilized in sensor technology, offering insights into molecular recognition processes and surface chemistry.

Synthesis

The synthesis of thionine typically involves a condensation reaction between dimethylaminobenzaldehyde and thiourea, which is catalyzed by an acid. This process can be summarized in a series of steps that yield pure thionine suitable for various applications[13]. The production method is crucial as it affects the quality and functionality of the resulting compound, which is widely used in biological systems due to its metachromatic properties and potential effects on electron transfer processes[14][15].

Furthermore, thionine can be synthesized in the presence of specific reagents to enhance its properties. For instance, various modifications have been explored to improve its reactivity and compatibility with different substrates, which is particularly relevant in the context of applications like wastewater treatment[16]. The synthetic route may also involve the use of thiol functionalities or derivatives to tailor the compound for specific uses, such as in the development of self-assembled monolayers or nanocomposite materials[3][17].

Research has indicated that the optimization of thionine synthesis can lead to enhanced performance in applications such as microbial fuel cells, where it can increase microbial activity and energy output, showcasing its versatility and importance in both industrial and biomedical fields[14][18].

Applications

Biomedical Applications

Thionine has garnered significant attention in the biomedical field, particularly for its role in biosensing technologies. Its ability to interact with various biomolecules allows for the development of electrochemical biosensors that can detect disease markers with high sensitivity and specificity. These sensors can monitor biomolecules such as glucose, proteins, DNA, and cholesterol in body fluids, thereby facilitating early diagnosis and routine disease management[3][4].

Peptide-Self-Assembled Monolayers (SAMs)

One innovative application of thionine lies in its incorporation into peptide-SAMs, which are utilized for biosensing. Peptide-SAMs enable the binding of target analytes, such as enzymes, nucleic acids, and antibodies, enhancing the performance of biosensors[3]. The use of thionine in these systems allows for the manipulation of surface properties, promoting cell adhesion and biosensing capabilities, which are essential for point-of-care diagnostics[3].

Electrochemical Sensors

Thionine-based electrochemical sensors have shown promise in the early detection of biomarkers due to their rapid signal amplification capabilities. These devices are designed to provide quick, non-invasive results, which can significantly enhance patient care by allowing timely medical intervention[4]. The combination of thionine with various nanomaterials has led to advancements in sensor design, enabling higher sensitivity and selectivity in detecting biological analytes[4].

Staining and Visualization

Beyond its application in biosensing, thionine is also employed in histological staining procedures. Its affinity for specific tissue components allows for effective visualization of cells and the extracellular matrix under light microscopy. This staining capability is crucial for studying tissue architecture and cellular interactions in various biological samples[5].

Environmental Monitoring

Thionine's properties extend to environmental applications as well. It has been utilized in the development of sensors for detecting pollutants and toxic substances in environmental samples. The ability of thionine to undergo reversible redox reactions enables its use in electrochemical sensors for environmental monitoring, making it a valuable tool in assessing water quality and detecting hazardous compounds[4].

Safety and Toxicity

Thionine, like many chemical compounds, poses certain safety and toxicity concerns that need to be addressed during handling and use. General safety guidelines should be adhered to in laboratory settings to minimize risks associated with chemical exposure. It is recommended to consult the ACS Guidelines for Chemical Laboratory Safety in Secondary Schools (2016) for best practices in laboratory work, as well as the ACS Division of Chemical Education Safety Guidelines for Chemical Demonstrations when conducting experiments involving thionine[19].

Inhalation of thionine may lead to respiratory tract irritation, although its toxicological properties have not been fully investigated[2]. Therefore, appropriate safety mea-

asures such as using chemical splash goggles and resistant gloves are essential when working with thionine solutions to prevent exposure[6].

Furthermore, thionine is associated with certain biological applications, including staining in histology. When used in such contexts, it is important to consider the potential for skin and eye irritation. Any accidental exposure should be managed by rinsing the affected area thoroughly and seeking medical advice if necessary.

Related Compounds

Thionine, an organic compound characterized by its unique dye properties, is part of a larger class of compounds that share structural or functional similarities. The interactions and derivatives of thionine have been extensively studied within the realms of chemistry and biochemistry.

Thiol Derivatives

In the context of the present invention, thiols are organic compounds that are defined by the presence of a thiol (-SH) group. Thiol derivatives retain this functional group or are transformed into thio ethers or thio esters, where the -SH group is replaced with an -SR group[12]. This class of compounds can exhibit significant biological activity, making them relevant in cosmetic and therapeutic applications. Preferred thiols in relation to thionine include those derived from amino acids like cysteine, which is known for its antioxidant properties[12][3].

Dye Characteristics and Classifications

Dyes, including thionine, can be categorized based on the chemical composition of their chromophores, which are responsible for the dye's color, and the auxochromes, which enhance binding to tissue[17].

Mordant Dyes: Require metallic salts for effective staining. Thionine can function as a mordant dye, particularly in histological applications[17][5].

Basic Dyes: These are cationic dyes that bind to acidic components in tissues, such as DNA and RNA. Thionine is utilized as a basic dye due to its affinity for nucleic acids[17][8].

Acid Dyes: Composed of a colored anion that binds to basic tissue components, acid dyes like eosin can sometimes be used in conjunction with thionine for comprehensive tissue staining[17][20].

Structural Similarities with Other Compounds

Thionine shares structural and functional attributes with a variety of other dyes and compounds. For instance, ortho-quinone derivatives can lead to rapid cleavage reactions with dye molecules, which may share mechanistic pathways with thionine in specific redox processes[17]. Additionally, thionine's properties are enhanced when integrated into self-assembled monolayers (SAMs) alongside thiol functionalities,

facilitating the development of bioactive surfaces for various biomedical applications-[\[3\]\[17\]](#).

References

- [1]: [Thionine | C₁₂H₁₀ClN₃S - ChemSpider](#)
- [2]: [Thionine | CAS#:581-64-6 | Chemsr](#)
- [3]: [Thionine blue | C₁₈H₂₂Cl₃N₃SSZn | CID 24213493 - PubChem](#)
- [4]: [Thionine | C₁₂H₁₀ClN₃S | CID 65043 - PubChem](#)
- [5]: [\[PDF\] Use of Thionine dye having structurally metachromatic influ](#)
- [6]: [Histological techniques 5. GENERAL STAINING](#)
- [7]: [Thionine - Wikipedia](#)
- [8]: [Peptide-based self-assembled monolayers \(SAMs\)](#)
- [9]: [Thionin staining and imaging technique - Google Patents](#)
- [10]: [Cosmetic and dermatological sunscreen compositions containing ...](#)
- [11]: [Buy Thionine | 581-64-6 | >90%](#)
- [12]: [Interaction Of Thionine Research Articles - R Discovery](#)
- [13]: [Thionin Acetate - Biological Staining & Histology Dye - GSP Chem](#)
- [14]: [Kinetics of the oxidative degradation of thionine dye by hydrogen ...](#)
- [15]: [Bioinspired versus Enzymatic Oxidation of Some Homologous ...](#)
- [16]: [Thionine dyes can be bleached by both ET and HAT mechanism ...](#)
- [17]: [Signal amplification strategies in electrochemical biosensors via ...](#)
- [18]: [Exploring The “Two-Faced” Thionin Reaction - ChemEd X](#)
- [19]: [\[PDF\] Thionin — The Two-Faced Solution - Flinn Scientific](#)
- [20]: [Thionine | 581-64-6 - ChemicalBook](#)