

HARMONIC 1-TYPE CURVES AND WEAK BIHARMONIC CURVES IN LORENTZIAN 3-SPACE

BY

H. KOCAYIĞIT, M. ÖNDER and H.H. HACISALIHOĞLU

Abstract. In this paper, we give definitions and characterizations of harmonic 1-type and weak biharmonic curves by using the mean curvature vector field of a Frenet curve in the Lorentzian 3-space L^3 . We also study weak biharmonic curves whose mean curvature vector fields are in the kernel of normal Laplacian ∇^\perp . We give some theorems for them in L^3 . Moreover, we give some characterizations and results for a Frenet curve in the same space.

Mathematics Subject Classification 2010: 53A35, 53C50.

Key words: harmonic 1-type curve, weak biharmonic curve, geodesic, helix, degenerate helix.

1. Introduction and preliminaries

Chen and Ishikawa have classified the Biharmonic curves in pseudo-Euclidean space E^n_ν . They have obtained that every biharmonic curve lies in a 3-dimensional totally geodesic subspace. Thus, it suffices to classify biharmonic curves in a semi-Euclidean 3-space (see CHEN and ISHIKAWA [2]). KOCAYIĞIT and HACISALIHOĞLU [13] have given the characterizations of harmonic and biharmonic curves in Euclidean 3-space. Also, they have studied the characterizations of 1-type and biharmonic curves in Lorentzian 3-space L^3 and found the relationships between the curvature and the torsion of these curves (see KOCAYIĞIT and HACISALIHOĞLU [12]).

In this paper, we give the characterizations of harmonic 1-type and weak biharmonic curves in semi-Euclidean 3-space in terms of curvature and torsion. Firstly, according to normal Levi-Civita connection, we point out the

general differential equations characterizing Frenet curves (both non-null and null) in Lorentzian 3-space L^3 . Moreover, we classify the special curves from the differential equations. More precisely, we show that a circular helix is a harmonic 1-type curve in L^3 and a pseudo circle (degenerate helix) is a weak biharmonic curve in the same space. In the final section, we give some theorems, corollaries and propositions.

Let now introduce a brief summary of the Lorentzian 3-space L^3 .

Let (M, g) be a time-oriented Lorentz 3-manifold. Let $\gamma : I \rightarrow M$ be a unit speed curve on M . Namely, the velocity vector field γ' satisfies $g(\gamma', \gamma') = \varepsilon_1 = \pm 1$, where g shows the Lorentzian metric given by $g(a, b) = -a_1b_1 + a_2b_2 + a_3b_3$ for the vectors $a = (a_1, a_2, a_3)$, $b = (b_1, b_2, b_3) \in L^3$. The constant ε_1 is called the causal character of γ . Then, a unit speed curve γ is said to be spacelike or timelike if its causal character is 1 or -1, respectively. A unit speed curve γ is called geodesic if $\nabla_{\gamma'}\gamma' = 0$, where ∇ is the Levi-Civita connection of (M, g) . Moreover, the curve γ is also said to be a geodesic if $\nabla_{\gamma'}^\perp\gamma' = 0$, where ∇^\perp is the normal Levi-Civita connection of (M, g) . A unit speed curve γ is said to be a Frenet curve if $g(\gamma'', \gamma'') \neq 0$. Like Euclidean geometry, every Frenet curve γ on (M, g) admits an orthonormal Frenet frame field $\{V_1, V_2, V_3\}$ along γ such that $V_1 = \gamma'(s)$ and $\{V_1, V_2, V_3\}$ satisfies the following Frenet-Serret formula (see IZUMIYA and TAKIYAMA [8,9] and KOBAYASHI [11]):

$$(1.1) \quad \begin{bmatrix} \nabla_{\gamma'} V_1 \\ \nabla_{\gamma'} V_2 \\ \nabla_{\gamma'} V_3 \end{bmatrix} = \begin{bmatrix} 0 & \varepsilon_2 \kappa & 0 \\ -\varepsilon_1 \kappa & 0 & -\varepsilon_3 \tau \\ 0 & \varepsilon_2 \tau & 0 \end{bmatrix} \begin{bmatrix} V_1 \\ V_2 \\ V_3 \end{bmatrix}.$$

The functions $\kappa \geq 0$ and τ are called the curvature and torsion, respectively. The vector fields V_1, V_2, V_3 are called tangent vector field, principle normal vector field and binormal vector field of γ , respectively. The constants ε_2 and ε_3 are defined by

$$(1.2) \quad \varepsilon_i = g(V_i, V_i), \quad i = 2, 3,$$

and called second causal character and third causal character of γ , respectively. Note that

$$(1.3) \quad \varepsilon_3 = -\varepsilon_1 \varepsilon_2.$$

As in the case of Riemannian geometry, a Frenet curve γ is a geodesic if and only if $\kappa = 0$. A circular helix is a Frenet curve whose curvature

and torsion are constants. A Frenet curve with constant curvature and zero torsion is called a pseudo circle. Pseudo circles are regarded as degenerate helices. Helices, which are not circles, are frequently called proper helices.

Here we use the normal Laplacian.

Let $\gamma : I \rightarrow M$ be a curve parametrized by the arclength on Lorentz 3-manifold M . Denote by $\{V_1, V_2, V_3\}$ the Frenet frame of γ as before. The normal bundle $\chi^\perp(\gamma(I))$ of the curve γ is given by

$$\chi^\perp(\gamma(I)) = Sp\{V_2(s), V_3(s)\}.$$

The normal connection $\nabla_{\gamma'}^\perp$ of γ is

$$(1.4) \quad \begin{aligned} \nabla_{\gamma'}^\perp : \chi^\perp(\gamma(I)) &\rightarrow \chi^\perp(\gamma(I)) \\ \nabla_{\gamma'}^\perp \xi &= \nabla_{\gamma'} \xi - g(\nabla_{\gamma'} \xi, V_1) V_1, \quad \forall \xi \in \chi^\perp(\gamma(I)). \end{aligned}$$

$\nabla_{\gamma'}^\perp \xi$ is called normal component of $\nabla_{\gamma'} \xi$. According to (1.4), the Frenet-Serret formulas are given by

$$(1.5) \quad \nabla_{\gamma'}^\perp V_2 = -\varepsilon_3 \tau V_3, \quad \nabla_{\gamma'}^\perp V_3 = \varepsilon_2 \tau V_2.$$

Let us denote the normal Laplace-Beltrami operator by Δ^\perp of γ and mean curvature vector field along γ by H . The mean curvature vector field H of a unit speed curve γ is $H = \varepsilon_1 \nabla_{\gamma'} \gamma'$. If γ is a Frenet curve, then H is given by

$$(1.6) \quad H = -\varepsilon_3 \kappa V_2,$$

where κ is the curvature of γ .

The normal Laplacian operator of γ is defined by

$$(1.7) \quad \Delta^\perp = -\varepsilon_1 \left(\nabla_{\gamma'}^\perp \right)^2 = -\varepsilon_1 \nabla_{\gamma'}^\perp \nabla_{\gamma'}^\perp$$

(see CHEN and ISHIKAWA [3], FERRANDEZ, LUCAS and MERONO [4] and IKAWA [7]).

To close this section, we recall the notion of harmonic 1-type and weak biharmonicity for unit speed curves.

Definition 1.1. A unit speed curve $\gamma : I \rightarrow M$ on a Lorentz 3-manifold M is said to be harmonic 1-type if

$$(1.8) \quad \Delta^\perp H = \lambda H$$

holds (KILIÇ [10]).

Definition 1.2. A unit speed curve $\gamma : I \rightarrow M$ on a Lorentz 3-manifold M is said to be weak biharmonic if

$$(1.9) \quad \Delta^\perp H = 0,$$

holds (KILIÇ [10]).

Definition 1.3. A unit speed curve $\gamma : I \rightarrow M$ on a Lorentz 3-manifold M is said to be a general helix if $\frac{\kappa}{\tau}$ is constant but κ and τ are not constant (HACISALİHOĞLU and ÖZTÜRK [5,6]).

2. Harmonic 1-type non-null curves and weak biharmonic non-null curves on Lorentzian 3-manifolds

In this section, we give the characterizations of harmonic 1-type and weak biharmonic non-null curves in Lorentzian 3-space.

Theorem 2.1. *Let γ be an arclength parametrized non-null Frenet curve on Lorentzian 3-manifold (M, g) . Then, along the curve γ , $\Delta^\perp H = \lambda H$ holds if and only if*

$$(2.1) \quad \varepsilon_1 \varepsilon_2 \kappa \tau^2 - \varepsilon_1 \varepsilon_3 \kappa'' = \varepsilon_3 \lambda \kappa, \quad 2\kappa' \tau + \kappa \tau' = 0.$$

Proof. From (1.5), (1.6) and (1.7) we get

$$(2.2) \quad \Delta^\perp H = (-\varepsilon_1 \varepsilon_2 \kappa \tau^2 + \varepsilon_1 \varepsilon_3 \kappa'')V_2 - (2\kappa' \tau + \kappa \tau')V_3.$$

Then, from (2.2) and (1.8) we have

$$(2.3) \quad (-\varepsilon_1 \varepsilon_2 \kappa \tau^2 + \varepsilon_1 \varepsilon_3 \kappa'')V_2 - (2\kappa' \tau + \kappa \tau')V_3 = \varepsilon_3 \lambda \kappa V_2.$$

According to (2.3), the equations (2.1) are obtained.

Conversely, the equations (2.1) satisfy the equation (1.8). \square

Theorem 2.2. *Let γ be an arclength parametrized non-null Frenet curve on Lorentzian 3-manifold (M, g) . Then, along the curve γ , $\Delta^\perp H = \lambda H$ holds if γ is a circular helix, where*

$$(2.4) \quad \lambda = -\tau^2.$$

Proof. Since γ is a circular helix, the curvatures κ and τ are constants. Then (2.1) holds and we have $\lambda = -\tau^2$. From that $\Delta^\perp H = \lambda H$ is satisfied. \square

Corollary 2.1. *Let γ be a non-null Frenet curve on Lorentzian 3-manifold (M, g) . Then γ is harmonic 1-type if γ is a circular helix.*

Theorem 2.3. *Let γ be an arclength parametrized non-null Frenet curve on Lorentzian 3-manifold (M, g) . Then, γ is weak biharmonic curve if and only if*

$$(2.5) \quad \varepsilon_1 \varepsilon_2 \kappa \tau^2 - \varepsilon_1 \varepsilon_3 \kappa'' = 0, \quad 2\kappa' \tau + \kappa \tau' = 0.$$

Proof. Since γ is weak biharmonic, $\Delta^\perp H = 0$ holds. From Theorem 2.1, we have (2.5). Conversely, the Equations (2.5) satisfy the equation $\Delta^\perp H = 0$. Thus, γ is weak biharmonic curve. \square

Corollary 2.2. *Let γ be a non-null Frenet curve on Lorentzian 3-manifold (M, g) . Then γ is weak biharmonic if γ is a pseudo circle (degenerate helix).*

Example 2.1. Let consider the spacelike curve $\gamma(s)$ given by the parametrization

$$\gamma(s) = \left(a \cosh \frac{s}{\sqrt{a^2 + b^2}}, a \sinh \frac{s}{\sqrt{a^2 + b^2}}, \frac{bs}{\sqrt{a^2 + b^2}} \right); \quad a, b > 0.$$

Frenet vectors of $\gamma(s)$ are found as follows

$$\begin{aligned} V_1(s) &= \frac{1}{\sqrt{a^2 + b^2}} \left(a \sinh \frac{s}{\sqrt{a^2 + b^2}}, a \cosh \frac{s}{\sqrt{a^2 + b^2}}, b \right), \\ V_2(s) &= \left(\cosh \frac{s}{\sqrt{a^2 + b^2}}, \sinh \frac{s}{\sqrt{a^2 + b^2}}, 0 \right), \\ V_3(s) &= \frac{1}{\sqrt{a^2 + b^2}} \left(b \sinh \frac{s}{\sqrt{a^2 + b^2}}, b \cosh \frac{s}{\sqrt{a^2 + b^2}}, -a \right). \end{aligned}$$

Then, curvature and torsion of $\gamma(s)$ are

$$\kappa = \frac{a}{a^2 + b^2}, \quad \tau = -\frac{b}{a^2 + b^2},$$

respectively. From (1.6) the mean curvature vector field of $\gamma(s)$ is

$$H(s) = -\frac{a}{a^2 + b^2} \left(\cosh \frac{s}{\sqrt{a^2 + b^2}}, \sinh \frac{s}{\sqrt{a^2 + b^2}}, 0 \right).$$

Then by using (1.7) we have

$$\Delta^\perp H = \frac{ab^2}{(a^2 + b^2)^3} \left(\cosh \frac{s}{\sqrt{a^2 + b^2}}, \sinh \frac{s}{\sqrt{a^2 + b^2}}, 0 \right),$$

which gives us $\Delta^\perp H(s) = \frac{-b^2}{(a^2 + b^2)^2} H(s)$. By Definition 1.1 we have that $\gamma(s)$ is harmonic 1-type curve with $\lambda = \frac{-b^2}{(a^2 + b^2)^2}$. Moreover, it is easily seen that (2.1) holds.

Furthermore, since κ and τ are constants, $\gamma(s)$ is a spacelike circular helix. Then by Theorem 2.2 we have $\lambda = -\tau^2$ which can be seen from the obtained equalities of λ and τ .

Example 2.2. Let now consider timelike curve $\gamma(s)$ given by $\gamma(s) = (\sinh s, \cosh s, 0)$ which is a pseudo-circle. Then Frenet vectors of $\gamma(s)$ are

$$V_1(s) = (\cosh s, \sinh s, 0), \quad V_2(s) = (\sinh s, \cosh s, 0), \quad V_3(s) = (0, 0, 1)$$

and curvature and torsion are $\kappa = 1$, $\tau = 0$ respectively. From (1.6) the mean curvature vector field of $\gamma(s)$ is $H(s) = -(\sinh s, \cosh s, 0)$. Then by using (1.7) we have $\Delta^\perp H = 0$ which gives us that $\gamma(s)$ is weak-biharmonic timelike circle(degenerate timelike helix) and it is easily seen that (2.5) holds.

3. General characterizations for a non-null Frenet curve and some results

In this section, we give the general characterizations of a Frenet curve by means of the mean curvature vector H and Frenet vectors.

Theorem 3.1. *Let γ be a unit speed non-null Frenet curve on Lorentzian 3-manifold (M, g) . According to the normal connection, the following equation holds*

$$(3.1) \quad \Delta^\perp H + \lambda \nabla_\gamma^\perp H + \mu H = 0,$$

where

$$(3.2) \quad \lambda = \frac{2\kappa'\tau + \kappa\tau'}{\kappa\tau}, \quad \mu = \frac{\kappa\tau^2 + \varepsilon_1\kappa''}{\kappa} - \lambda \frac{\kappa'}{\kappa}.$$

Proof. By (2.2), (1.6) and Frenet formulas (1.5) we get

$$(3.3) \quad \Delta^\perp H = (-\varepsilon_1 \varepsilon_2 \kappa \tau^2 + \varepsilon_1 \varepsilon_3 \kappa'') V_2 - (2\kappa' \tau + \kappa \tau') V_3$$

and

$$(3.4) \quad \nabla_{\gamma'}^\perp H = -\varepsilon_3 \kappa' V_2 + \kappa \tau V_3.$$

Furthermore, from (1.5) and (1.6) we have

$$(3.5) \quad V_2 = -\varepsilon_3 \frac{1}{\kappa} H, \quad V_3 = \frac{1}{\tau} \left[\left(\frac{1}{\kappa} \right)' H + \frac{1}{\kappa} \nabla^\perp H \right].$$

Substituting the equations (3.5) into the (3.4) we have $\Delta^\perp H + \lambda \nabla_{\gamma'}^\perp H + \mu H = 0$ where $\lambda = \frac{2\kappa' \tau + \kappa \tau'}{\kappa \tau}$, $\mu = \frac{\kappa \tau^2 + \varepsilon_1 \kappa''}{\kappa} - \lambda \frac{\kappa'}{\kappa}$. \square

From Theorem 3.1 we have the following corollaries:

Corollary 3.1. *Let γ be an arclength parametrized non-null Frenet curve on Lorentzian 3-manifold (M, g) . If γ is a circular helix, then according to the normal connection, $\Delta^\perp H + \tau^2 H = 0$ holds.*

Proof. The proof is clear from (3.1) and (3.2). \square

Corollary 3.2. *Let γ be an arclength parametrized non-null Frenet curve on Lorentzian 3-manifold (M, g) . If γ is a general helix, then according to the normal connection, $\Delta^\perp H + \lambda \nabla_{\gamma'}^\perp H + \mu H = 0$ holds, where*

$$\lambda = 3 \frac{\kappa'}{\kappa}, \quad \mu = \frac{\kappa \tau^2 + \varepsilon_1 \kappa''}{\kappa} - 3 \left(\frac{\kappa'}{\kappa} \right)^2.$$

Proof. The proof can be easily seen from (3.1) and (3.2). \square

From Corollary 3.2 we can find the following result.

Result 3.1. Let γ be an arclength parametrized non-null Frenet curve on Lorentzian 3-manifold (M, g) . Then according to the normal connection

i) If γ is a timelike general helix, then

$$\Delta^\perp H + 3 \frac{\kappa'}{\kappa} \nabla_{\gamma'}^\perp H + \left[\frac{\kappa \tau^2 - \kappa''}{\kappa} - 3 \left(\frac{\kappa'}{\kappa} \right)^2 \right] H = 0$$

holds.

ii) If γ is a spacelike general helix, then

$$\Delta^\perp H + 3 \frac{\kappa'}{\kappa} \nabla_{\gamma'}^\perp H + \left[\frac{\kappa \tau^2 + \kappa''}{\kappa} - 3 \left(\frac{\kappa'}{\kappa} \right)^2 \right] H = 0.$$

When the curve γ is a geodesic we have Theorem 2.3 again, i.e., γ is a weak biharmonic curve.

Theorem 3.2. *Let γ be an arclength parametrized non-null Frenet curve on Lorentzian 3-manifold (M, g) . According to the normal connection, γ satisfies the following differential equation*

$$(3.6) \quad \tau \left(\nabla_{\gamma'}^\perp \right)^2 V_2 - \tau' \nabla_{\gamma'}^\perp V_2 - \varepsilon_1 \tau^3 V_2 = 0.$$

Proof. From the Frenet formulas (1.5) we have

$$(3.7) \quad \left(\nabla_{\gamma'}^\perp \right)^2 V_2 = \varepsilon_1 \tau^2 V_2 - \varepsilon_3 \tau' V_3$$

and since $V_3 = -\varepsilon_3 \frac{1}{\tau} \nabla_{\gamma'}^\perp V_2$, from (3.7) we have the equation (3.6). \square

From Theorem 3.2 we have the following corollary.

Corollary 3.3. *Let γ be an arclength parametrized non-null Frenet curve on Lorentzian 3-manifold (M, g) . If γ is a circular helix then the following equation holds $\left(\nabla_{\gamma'}^\perp \right)^2 V_2 - \varepsilon_1 \tau^2 V_2 = 0$.*

Proof. The proof is clear from (3.6). \square

Result 3.2. Let γ be an arclength parametrized non-null Frenet curve on Lorentzian 3-manifold (M, g) .

- i) If γ is a timelike circular helix then $\left(\nabla_{\gamma'}^\perp \right)^2 V_2 + \tau^2 V_2 = 0$ holds.
- ii) If γ is a spacelike circular helix then $\left(\nabla_{\gamma'}^\perp \right)^2 V_2 - \tau^2 V_2 = 0$ holds.

Corollary 3.4. *Let γ be an arclength parametrized non-null Frenet curve on Lorentzian 3-manifold (M, g) . If γ is a general helix then according to the normal connection the following equality holds $\kappa \left(\nabla_{\gamma'}^\perp \right)^2 V_2 - \kappa' \nabla_{\gamma'}^\perp V_2 - \varepsilon_1 \kappa \tau^2 V_2 = 0$.*

Proof. Since γ is a general helix, by taking $\frac{\kappa}{\tau} = \text{const.}$, i.e., $\tau' = \frac{\kappa'}{\kappa} \tau$, from (3.6) the proof follows immediately. \square

Result 3.3. Let γ be an arclength parametrized non-null Frenet curve on Lorentzian 3-manifold (M, g) . Then

i) If γ is a timelike general helix, then $\kappa \left(\nabla_{\gamma'}^\perp \right)^2 V_2 - \kappa' \nabla_{\gamma'}^\perp V_2 + \kappa \tau^2 V_2 = 0$, holds.

ii) If γ is a spacelike general helix, then $\kappa \left(\nabla_{\gamma'}^\perp \right)^2 V_2 - \kappa' \nabla_{\gamma'}^\perp V_2 - \kappa \tau^2 V_2 = 0$, holds.

Theorem 3.3. *Let γ be an arclength parametrized non-null Frenet curve on Lorentzian 3-manifold (M, g) . According to the normal connection, γ satisfies the following differential equation*

$$(3.8) \quad \tau \left(\nabla_{\gamma'}^\perp \right)^2 V_3 - \tau' \nabla_{\gamma'}^\perp V_3 - \varepsilon_1 \tau^3 V_3 = 0.$$

Proof. From the Frenet formulas (1.5) we have

$$(3.9) \quad \left(\nabla_{\gamma'}^\perp \right)^2 V_3 = \varepsilon_2 \tau' V_2 - \varepsilon_2 \varepsilon_3 \tau^2 V_3$$

and since $V_2 = \varepsilon_2 \frac{1}{\tau} \nabla_{\gamma'}^\perp V_3$, from (3.9) we have the equation (3.8). \square

From Theorem 3.3 we have the following corollary.

Corollary 3.5. *Let γ be an arclength parametrized non-null Frenet curve on Lorentzian 3-manifold (M, g) . If γ is a circular helix then the following equation holds, $\left(\nabla_{\gamma'}^\perp \right)^2 V_3 - \varepsilon_1 \tau^2 V_3 = 0$.*

Proof. By (3.8), the proof can be easily seen. \square

Result 3.4. Let γ be an arclength parametrized non-null Frenet curve on Lorentzian 3-manifold (M, g) .

i) If γ is a timelike circular helix then $\left(\nabla_{\gamma'}^\perp \right)^2 V_3 + \tau^2 V_3 = 0$ holds.

ii) If γ is a spacelike circular helix then $\left(\nabla_{\gamma'}^\perp \right)^2 V_3 - \tau^2 V_3 = 0$ holds.

Corollary 3.6. *Let γ be an arclength parametrized non-null Frenet curve on Lorentzian 3-manifold (M, g) . If γ is a general helix then according to the normal connection the following equality holds*

$$\kappa \left(\nabla_{\gamma'}^\perp \right)^2 V_3 + \kappa' \nabla_{\gamma'}^\perp V_3 - \varepsilon_1 \kappa \tau^2 V_3 = 0.$$

Proof. Since γ is a general helix, by taking $\frac{\kappa}{\tau} = \text{const.}$, i.e., $\tau' = \frac{\kappa'}{\kappa} \tau$ from (3.8) the proof is straightforward. \square

Result 3.5. Let γ be an arclength parametrized non-null Frenet curve on Lorentzian 3-manifold (M, g) . Then

i) If γ is a timelike general helix, then $\kappa \left(\nabla_{\gamma'}^\perp \right)^2 V_3 + \kappa' \nabla_{\gamma'}^\perp V_3 + \kappa \tau^2 V_3 = 0$ holds.

ii) If γ is a spacelike general helix, then $\kappa (\nabla_{\gamma'}^\perp)^2 V_3 + \kappa' \nabla_{\gamma'}^\perp V_3 - \kappa \tau^2 V_3 = 0$ holds.

Example 3.1. For the timelike curve $\gamma(s)$ given by the parametrization

$$\gamma(s) = \left(\frac{s^2}{2} + s, \frac{s^2}{2}, \frac{2\sqrt{2}}{3}s^{3/2} \right); \quad (s \neq 0),$$

the Frenet vectors are obtained as follows,

$$V_1(s) = (s+1, s, \sqrt{2s}), \quad V_2(s) = (\sqrt{2s}, \sqrt{2s}, 1), \quad V_3(s) = \left(1, 1, \frac{1}{\sqrt{2s}} \right),$$

and, curvature and torsion functions are $\kappa = \frac{\sqrt{2}}{2\sqrt{s}}$, $\tau = \frac{\sqrt{2}}{2\sqrt{s}}$, respectively. From (1.6) the mean curvature vector field of $\gamma(s)$ is

$$H(s) = \left(-\frac{\sqrt{2}}{s}, -\frac{\sqrt{2}}{s}, -\frac{1}{s^{3/2}} \right).$$

Then we have the followings

$$\Delta^\perp H = \left(\frac{2s-3}{4s^2}, \frac{2s-6}{4s^2}, \frac{\sqrt{2}(2s-3)}{8s^{5/2}} \right), \quad \nabla_{\gamma'}^\perp H = \left(\frac{1}{2s}, \frac{1}{2s}, \frac{\sqrt{2}}{4s^{3/2}} \right).$$

Thus it is obtained that (3.1) holds for $\lambda = -\frac{3}{2s}$, $\mu = \frac{\sqrt{2}(2s-3)-6\sqrt{s}}{8s^{5/2}}$. Similarly, Frenet vector V_2 satisfies equation (3.6) and equation (3.8) holds for the Frenet vector V_3 .

Moreover, we have that $\frac{\kappa}{\tau}$ is constant i.e., $\gamma(s)$ is a general timelike helix. Then it is obvious that $\gamma(s)$ satisfies the conditions of Corollary 3.2, Result 3.1 (i), Corollary 3.4, Result 3.3 (i) and Result 3.5 (i).

4. Harmonic 1-type null curves and weak biharmonic null curves on Lorentzian 3-manifolds

Let γ be a curve on Lorentzian 3-manifold (M, g) . Then, the curve γ is

called a null curve if $g(V_1, V_1) = 0$. By a Cartan frame $\{V_1, V_2, V_3\}$ of γ we mean a family of vector fields $V_1 = V_1(s)$, $V_2 = V_2(s)$, $V_3 = V_3(s)$ along the curve γ satisfying the following conditions:

$$\begin{aligned} \gamma'(s) &= V_1, \quad g(V_1, V_1) = g(V_2, V_2) = 0, \quad g(V_1, V_2) = -1, \\ g(V_1, V_3) &= g(V_2, V_3) = 0, \quad g(V_3, V_3) = 1, \end{aligned}$$

$$(4.1) \quad \begin{bmatrix} \nabla_{\gamma'} V_1 \\ \nabla_{\gamma'} V_2 \\ \nabla_{\gamma'} V_3 \end{bmatrix} = \begin{bmatrix} 0 & 0 & \kappa \\ 0 & 0 & \tau \\ \tau & \kappa & 0 \end{bmatrix} \begin{bmatrix} V_1 \\ V_2 \\ V_3 \end{bmatrix},$$

where κ and τ are called the curvature and torsion of γ , respectively (IKAWA [7]). Here V_1 and V_2 are null vectors and V_3 is a unit spacelike vector.

According to (1.4)

$$(4.2) \quad \nabla_{\gamma'}^\perp V_2 = \tau V_3, \quad \nabla_{\gamma'}^\perp V_3 = \kappa V_2.$$

Let us denote by H the mean curvature vector field of γ and by the normal Laplace-Beltrami operator Δ^\perp of γ . Then the mean curvature vector field H is defined by

$$(4.3) \quad H = \nabla_{\gamma'} V_1 = \kappa V_3$$

and the normal Laplacian operator along γ is given by

$$(4.4) \quad \Delta^\perp = -\left(\nabla_{\gamma'}^\perp\right)^2 = -\nabla_{\gamma'}^\perp \nabla_{\gamma'}^\perp$$

(see CHEN and ISHIKAWA [2,3]).

Theorem 4.1. *Let γ be a null Frenet curve on Lorentzian 3-manifold (M, g) . Then, along the curve γ , $\Delta^\perp H = \lambda H$ holds if and only if*

$$(4.5) \quad \kappa^2 \tau + \kappa'' = -\lambda \kappa, \quad \kappa' \kappa = 0.$$

Proof. From (4.2), (4.3) and (4.4) we get

$$(4.6) \quad \Delta^\perp H = -3\kappa \kappa' V_2 - (\kappa^2 \tau + \kappa'') V_3.$$

By (4.3) and (1.8), we have

$$(4.7) \quad -3\kappa \kappa' V_2 - (\kappa^2 \tau + \kappa'') V_3 = \lambda \kappa V_3.$$

According to (4.7), the equations (4.5) are obtained. □

Conversely, the equations (4.5) satisfy the equation $\Delta^\perp H = \lambda H$.

Theorem 4.2. *Let γ be a null Frenet curve on Lorentzian 3-manifold (M, g) . Then, along the curve γ , $\Delta^\perp H = \lambda H$ holds if and only if γ is a null curve with constant curvature, where $\lambda = -\kappa\tau$.*

Proof. From Theorem 4.1, we have (4.5). Since γ is a Frenet curve, $\kappa \neq 0$. Thus (4.5) shows that κ is a constant and $\lambda = -\kappa\tau$. Thus γ is a null curve with constant curvature.

Conversely, since γ is a null curve with constant curvature and $\lambda = -\kappa\tau$, $\Delta^\perp H = \lambda H$ is satisfied. \square

Corollary 4.1. *Let γ be a null Frenet curve on Lorentzian 3-manifold (M, g) . Then, γ is harmonic 1-type if and only if γ is a null curve with constant curvature.*

Theorem 4.3. *Let γ be a null Frenet curve on Lorentzian 3-manifold (M, g) . Then, along the curve γ , $\Delta^\perp H = 0$ holds if and only if*

$$(4.8) \quad \kappa^2\tau + \kappa'' = 0, \quad \kappa'\kappa = 0.$$

Proof. From Theorem 4.1 we have (4.8). Conversely, (4.8) satisfy the equation $\Delta^\perp H = 0$. \square

Theorem 4.4. *Let γ be a null Frenet curve on Lorentzian 3-manifold (M, g) . Then, along the curve γ , $\Delta^\perp H = 0$ holds if and only if γ is a pseudo null circle (degenerate null helix).*

Proof. From Theorem 4.3, we have (4.8). Since γ is a Frenet curve, $\kappa \neq 0$. Thus, the equations (4.8) show that γ is a degenerate null helix.

Conversely, since γ is a degenerate null helix, κ is a constant and $\tau = 0$. Thus the equations (4.8) satisfy the equation $\Delta^\perp H = 0$. \square

Corollary 4.2. *Let γ be a null Frenet curve on Lorentzian 3-manifold (M, g) . Then γ is a weak biharmonic if and only if γ is a pseudo null circle (degenerate null helix).*

Example 4.1. Let consider the parametrized null curve $\gamma(s)$ given by $\gamma(s) = \left(\frac{1}{2} \sinh 2s, \frac{1}{2} \cosh 2s, s\right)$, (see BALGETİR, BEKTAŞ and INOBUCHI [1]). Then, Frenet vectors of $\gamma(s)$ are obtained as follows

$$\begin{aligned} V_1(s) &= \gamma'(s) = (\cosh 2s, \sinh 2s, 1), \\ V_2(s) &= \left(\frac{1}{2} \cosh 2s, \frac{1}{2} \sinh 2s, -\frac{1}{2}\right), \\ V_3(s) &= (\sinh 2s, \cosh 2s, 0). \end{aligned}$$

The curvature and torsion of $\gamma(s)$ are $\kappa = 2$, $\tau = -1$ respectively. From (4.3) mean curvature vector field of $\gamma(s)$ is $H(s) = (2 \sinh 2s, \cosh 2s, 0)$. Then, by considering (4.4) we have $\Delta^\perp H = 2H$ which gives that $\gamma(s)$ is harmonic 1-type null curve and it is clear that $\gamma(s)$ satisfies the condition of Theorem 4.1, Theorem 4.2 and Corollary 4.1.

5. General characterizations for a null Frenet curve and some results

In this section, we give the general characterizations of a null Frenet curve by means of the mean curvature vector H and Frenet vectors according to the normal Laplacian operator Δ^\perp and the normal connection ∇^\perp .

Theorem 5.1. *Let γ be a null Frenet curve on Lorentzian 3-manifold (M, g) . According to the normal connection, the following equation holds*

$$(5.1) \quad \Delta^\perp H + \lambda \nabla_{\gamma'}^\perp H + \mu H = 0,$$

where

$$(5.2) \quad \lambda = 3 \frac{\kappa'}{\kappa}, \quad \mu = \frac{\kappa^2 \tau + \kappa''}{\kappa} - 3 \left(\frac{\kappa'}{\kappa} \right)^2.$$

Proof. By (4.6), (4.3) and Frenet formulas (4.1)-(4.2), we get

$$(5.3) \quad \Delta^\perp H = -3\kappa\kappa'V_2 - (\kappa^2\tau + \kappa'')V_3,$$

$$(5.4) \quad \nabla_{\gamma'}^\perp H = \kappa^2 V_2 + \kappa' V_3.$$

Furthermore, from (4.2) and (4.3) we have

$$(5.5) \quad V_3 = \frac{1}{\kappa} H, \quad V_2 = \frac{1}{\tau} \left[\left(\frac{1}{\kappa} \right)' H + \frac{1}{\kappa} \nabla^\perp H \right].$$

Substituting the equations (5.5) into the (5.4) we have $\Delta^\perp H + \lambda \nabla_{\gamma'}^\perp H + \mu H = 0$, where $\lambda = 3 \frac{\kappa'}{\kappa}$, $\mu = \frac{\kappa^2 \tau + \kappa''}{\kappa} - 3 \left(\frac{\kappa'}{\kappa} \right)^2$. \square

From Theorem 5.1 we have the following corollaries:

Corollary 5.1. *Let γ be a null Frenet curve on Lorentzian 3-manifold (M, g) . If γ is a circular null helix, then according to the normal connection $\Delta^\perp H + \kappa\tau H = 0$ holds.*

Proof. The proof is clear from (5.1) and (5.2). \square

Corollary 5.2. *Let γ be a null Frenet curve on Lorentzian 3-manifold (M, g) . If γ is a general null helix then according to the normal connection the following equality holds $\Delta^\perp H + \lambda \nabla_\gamma^\perp H + \mu H = 0$, where*

$$\lambda = 3 \frac{\kappa'}{\kappa}, \quad \mu = \frac{\kappa^2}{\kappa'} \tau' + \frac{\kappa''}{\kappa} - 3 \left(\frac{\kappa'}{\kappa} \right)^2.$$

Proof. By taking $\tau = \frac{\kappa}{\kappa'} \tau'$, the proof can be easily seen from (5.1) and (5.2). \square

Theorem 5.2. *Let γ be a null Frenet curve on Lorentzian 3-manifold (M, g) . According to the normal connection, γ satisfies the following differential equation*

$$(5.6) \quad \tau \left(\nabla_{\gamma'}^\perp \right)^2 V_2 - \tau' \nabla_{\gamma'}^\perp V_2 - \kappa \tau^2 V_2 = 0.$$

Proof. From the Frenet formulas (4.2) we have

$$(5.7) \quad \left(\nabla_{\gamma'}^\perp \right)^2 V_2 = \kappa \tau V_2 + \tau' V_3$$

and since $V_3 = \frac{1}{\tau} \nabla_{\gamma'}^\perp V_2$, from (5.7) we have the equation (5.6). \square

From Theorem 5.2 we have the following corollary.

Corollary 5.3. *Let γ be a null Frenet curve on Lorentzian 3-manifold (M, g) . If γ is a circular null helix then the following equation holds $(\nabla_{\gamma'}^\perp)^2 V_2 - \kappa \tau V_2 = 0$.*

Proof. The proof is clear from (5.6). \square

Corollary 5.4. *Let γ be a null Frenet curve on Lorentzian 3-manifold (M, g) . If γ is a general null helix then according to the normal connection the following equality holds $\kappa \left(\nabla_{\gamma'}^\perp \right)^2 V_2 - \kappa' \nabla_{\gamma'}^\perp V_2 - \kappa^2 \tau V_2 = 0$.*

Proof. Since γ is a general helix, by taking $\frac{\kappa}{\tau} = \text{const.}$ i.e., $\tau' = \frac{\kappa'}{\kappa} \tau$, from (5.6) the proof follows immediately. \square

Theorem 5.3. *Let γ be a null Frenet curve on Lorentzian 3-manifold (M, g) . According to the normal connection, γ satisfies the following differential equation*

$$(5.8) \quad \kappa \left(\nabla_{\gamma'}^\perp \right)^2 V_3 - \kappa' \nabla_{\gamma'}^\perp V_3 - \kappa^2 \tau V_3 = 0.$$

Proof. From the Frenet formulas (4.2) we have

$$(5.9) \quad \left(\nabla_{\gamma'}^\perp \right)^2 V_3 = \kappa' V_2 + \kappa \tau V_3$$

and since $V_2 = \frac{1}{\kappa} \nabla_{\gamma'}^\perp V_3$, from (5.9) we have the equation (5.8). \square

From Theorem 5.3 we have the following corollary.

Corollary 5.5. *Let γ be a null Frenet curve on Lorentzian 3-manifold (M, g) . If γ is a circular null helix then the following equation holds,*

$$\left(\nabla_{\gamma'}^\perp \right)^2 V_3 - \kappa \tau V_3 = 0.$$

Proof. By (5.8), the proof can be easily seen. \square

Corollary 5.6. *Let γ be a null Frenet curve on Lorentzian 3-manifold (M, g) . If γ is a general null helix then according to the normal connection the following equality holds*

$$\tau \left(\nabla_{\gamma'}^\perp \right)^2 V_3 - \tau' \nabla_{\gamma'}^\perp V_3 - \kappa \tau^2 V_3 = 0.$$

Proof. Since γ is a general helix, by taking $\frac{\kappa}{\tau} = \text{const.}$ i.e., $\tau' = \frac{\kappa'}{\kappa} \tau$ from (5.8) the proof is straightforward. \square

The null curve $\gamma(s)$ given in Example 4.1 can be taken as an example of Section 5. It is easily seen that $\gamma(s)$ satisfies all conditions given in theorems and corollaries of this section.

REFERENCES

1. BALGETIR, H.; BEKTAŞ, M.; INOBUCHI, J. – *Null Bertrand curves in Minkowski 3-space and their characterizations*, Note Mat., 23 (2004/05), 7–13.
2. CHEN, B.-Y.; ISHIKAWA, S. – *Biharmonic surfaces in pseudo-Euclidean spaces*, Mem. Fac. Sci. Kyushu Univ. Ser. A, 45 (1991), 323–347.

3. CHEN, B.-Y.; ISHIKAWA, S. – *Biharmonic pseudo-Riemannian submanifolds in pseudo-Euclidean spaces*, Kyushu J. Math., 52 (1998), 167–185.
4. FERRÁNDEZ, A.; LUCAS, P.; MEROÑO, M.A. – *Biharmonic Hopf cylinders*, Rocky Mountain J. Math., 28 (1998), 957–975.
5. HACISALİHOĞLU, H.H.; ÖZTÜRK, R. – *On the characterization of inclined curves in E^n I.*, Tensor (N.S.), 64 (2003), 157–162.
6. HACISALİHOĞLU, H.H.; ÖZTÜRK, R. – *On the characterization of inclined curves in E^n II.*, Tensor (N.S.), 64 (2003), 163–169.
7. IKAWA, T. – *On curves and submanifolds in an indefinite-Riemannian manifold*, Tsukuba J. Math., 9 (1985), 353–371.
8. IZUMIYA, S.; TAKIYAMA, A. – *A time-like surface in Minkowski 3-space which contains pseudocircles*, Proc. Edinburgh Math. Soc., 40 (1997), 127–136.
9. IZUMIYA, S.; TAKIYAMA, A. – *A time-like surface in Minkowski 3-space which contains light-like lines*, J. Geom., 64 (1999), 95–101.
10. KILIÇ, B. – *Finite type curves and surfaces*, Ph. Thesis, University of Hacettepe, 2002.
11. KOBAYASHI, O. – *Maximal surfaces in the 3-dimensional Minkowski space L^3* , Tokyo J. Math., 6 (1983), 297–309.
12. KOCAYİĞİT, H.; HACISALİHOĞLU, H.H. – *1-type and biharmonic Frenet curves in Lorentzian 3-space*, Iran. J. Sci. Technol. Trans. A Sci., 33 (2009), 159–168.
13. KOCAYİĞİT, H.; HACISALİHOĞLU, H.H. – *1-type curves and biharmonic curves in Euclidean 3-space*, Int. Electron. J. Geom., 4 (2011), 97–101.

Received: 18.VII.2011

Revised: 2.XI.2011

Accepted: 23.XI.2011

Celal Bayar University,
Faculty of Science and Arts,
Department of Mathematics,
Muradiye Campus, Manisa,
TURKEY
huseyin.kocayigit@cbu.edu.tr
mehmet.onder@cbu.edu.tr

Ankara University,
Faculty of Science,
Department of Mathematics,
Tandoğan, Ankara,
TURKEY
hacisali@science.ankara.edu.tr