

Near-consistent robust estimations of moments for unimodal distributions

Tuban Lee^{a,1}

^aInstitute of Biomathematics, Macau SAR 999078, China

This manuscript was compiled on March 22, 2023

Descriptive statistics for parametric models currently heavily rely on the accuracy of distributional assumptions. Here, based on the invariant structures of unimodal distributions, a series of sophisticated yet efficient estimators, robust to both gross errors and departures from parametric assumptions, are proposed for estimating mean and central moments with insignificant asymptotic biases for common unimodal distributions. This article also illuminates the understanding of the common nature of probability distributions and the measures of them.

orderliness | invariant | unimodal | adaptive estimation | U -statistics

The asymptotic inconsistencies between sample mean (\bar{x}) and nonparametric robust location estimators in asymmetric distributions on the real line have been noticed for more than two centuries (1), yet remain unsolved. Strictly speaking, it is unsolvable as by trimming, some information about the original distribution is removed, making it impossible to estimate the values of the removed parts without distributional assumptions. Newcomb (1886, 1912) provided the first modern approach to robust parametric estimation by developing a class of estimators that gives "less weight to the more discordant observations" (2, 3). In 1964, Huber (4) used the minimax procedure to obtain M-estimator for the contaminated normal distribution, which has played a pre-eminent role in the later development of robust statistics. However, as previously demonstrated, under growing asymmetric departures from normality, the bias of the Huber M-estimator increases rapidly. This is a common issue in parameter estimations. For example, He and Fung (1999) constructed (5) a robust M-estimator for the two-parameter Weibull distribution, from which all moments can be calculated. Nonetheless, it is inadequate for the gamma, Perato, lognormal, and the generalized Gaussian distributions (SI Dataset S1). Another old and interesting approach is arithmetically computing the parameters using one or more L -statistics as inputs, such as percentile estimators. Examples of percentile estimators for the Weibull distribution, the reader is referred to Menon (1963) (6), Dubey (1967) (7), Hassanein (1971) (8), Marks (2005) (9), and Boudt, Caliskan, and Croux (2011) (10)'s works. At the outset of the study of percentile estimators, it was known that they arithmetically utilizes the invariant structures of probability distributions (6, 11, 12). Maybe such estimators can be named as I -statistics. Formally, an estimator is classified as an I -statistic if it asymptotically satisfies $I(LE_1, \dots, LE_l) = (\theta_1, \dots, \theta_q)$ for the distribution it is consistent, where LEs are calculated with the use of L -statistics, I is defined using arithmetic operations and constants, but it may also incorporate other functions, and θ s are the population parameters it estimates. A subclass of I -statistics, arithmetic I -statistics, is defined as LEs are L -statistics, I is solely defined using arithmetic operations and constants.

Since some percentile estimators use the logarithmic function to transform all random variables before computing the L -statistics, a percentile estimator might not always be an arithmetic I -statistic (7). In this article, two subclasses of I -statistics are introduced, arithmetic I -statistics and quantile I -statistics. Examples of quantile I -statistics will be discussed later. Based on L -statistics, I -statistics are naturally robust. Compared to probability density functions (pdfs) and cumulative distribution functions (cdfs), the quantile functions of many parametric distributions are more elegant. Since the expectation of an L -statistic can be expressed as an integral of the quantile function, I -statistics are often analytically obtainable. However, the performance of the aforementioned examples is often worse than that of the robust M -statistics when the distributional assumption is violated (SI Dataset S1). Even when distributions such as the Weibull and gamma belong to the same larger family, the generalized gamma distribution, a misassumption can still result in substantial biases, rendering the approach ill-suited.

In previous research on semiparametric robust mean estimation, the binomial mean (BM_ϵ) is still inconsistent for any skewed distribution, despite having much smaller asymptotic biases than other weighted averages. All robust location estimators commonly used are symmetric due to the universality of the symmetric distributions. One can construct an asymmetric weighted average that is consistent for a semiparametric class of skewed distributions. This approach has been investigated previously, but its lack of symmetry makes it suitable only for certain applications (13). Shifting from semiparametrics to parametrics, an ideal robust location estimator would have a non-sample-dependent breakdown point (defined in Subsection ??) and be consistent for any symmetric distribution and a skewed distribution with finite second moments. This is called an invariant mean. Based on the mean-symmetric weighted

Significance Statement

Bias, variance, and contamination are the three main errors in statistics. Consistent robust estimation is unattainable without parametric assumptions. Here, based on a paradigm shift inspired by mean-median-mode inequality, Bickel-Lehmann spread, and adaptive estimation, invariant moments are proposed as a means of achieving near-consistent and robust estimations of moments, even in scenarios where moderate violations of distributional assumptions occur, while the variances are sometimes smaller than those of the sample moments.

T.L. designed research, performed research, analyzed data, and wrote the paper.

The author declares no competing interest.

¹To whom correspondence should be addressed. E-mail: tl@biomathematics.org

average-median inequality, the recombined mean is defined as

$$rm_{d,\epsilon,n} := \lim_{c \rightarrow \infty} \left(\frac{(\text{SWA}_{\epsilon,n} + c)^{d+1}}{(\text{median} + c)^d} - c \right),$$

where d is the key factor for bias correction, $\text{SWA}_{\epsilon,n}$ is $\text{BM}_{\epsilon,n}$ in the first three Subsections, but other symmetric weighted averages can also be used in practice as long as the inequalities hold. The following theorem shows the significance of this arithmetic I -statistic.

Theorem .1. *If the second moments are finite, $rm_{d \approx 0.375, \epsilon = \frac{1}{8}}$ is a consistent mean estimator for the exponential and any symmetric distributions and the Pareto distribution with quantile function $Q(p) = x_m(1-p)^{-\frac{1}{\alpha}}$, $x_m > 0$, when $\alpha \rightarrow \infty$.*

Proof. Finding d and ϵ that make $rm_{d,\epsilon}$ a consistent mean estimator is equivalent to finding the solution of $E[rm_{d,\epsilon}] = E[X]$. Rearranging the definition, $rm_{d,\epsilon} = \lim_{c \rightarrow \infty} \left(\frac{(\text{BM}_{\epsilon} + c)^{d+1}}{(\text{median} + c)^d} - c \right) = (d+1)\text{BM}_{\epsilon} - d\text{median} = \mu$. So, $d = \frac{\mu - \text{BM}_{\epsilon}}{\text{BM}_{\epsilon} - \text{median}}$. The quantile function of the exponential distribution is $Q(p) = \ln\left(\frac{1}{1-p}\right)\lambda$. $E[x] = \lambda$. $E[\text{median}] = Q\left(\frac{1}{2}\right) = \ln 2\lambda$. For the exponential distribution, the expectation of $\text{BM}_{\frac{1}{8}}$ is $E\left[\text{BM}_{\frac{1}{8}}\right] = \lambda \left(1 + \ln\left(\frac{46656}{8575\sqrt{35}}\right)\right)$. Obviously, the scale parameter λ can be canceled out, $d \approx 0.375$. The proof of the second assertion follows directly from the coincidence property. For any symmetric distribution with a finite second moment, $E[\text{BM}_{\epsilon}] = E[\text{median}] = E[X]$. Then $E[rm_{d,\epsilon}] = \lim_{c \rightarrow \infty} \left(\frac{(E[X] + c)^{d+1}}{(E[X] + c)^d} - c \right) = E[X]$. The proof for the Pareto distribution is more general. The mean of the Pareto distribution is given by $\frac{\alpha x_m}{\alpha - 1}$. The d value with two unknown percentiles p_1 and p_2 for the Pareto distribution is $d_{\text{Pareto}} = \frac{\frac{\alpha x_m}{\alpha - 1} - x_m(1-p_1)^{-\frac{1}{\alpha}}}{x_m(1-p_1)^{-\frac{1}{\alpha}} - x_m(1-p_2)^{-\frac{1}{\alpha}}}$. Since any weighted average can be expressed as an integral of the quantile function, $\lim_{\alpha \rightarrow \infty} \frac{\frac{\alpha}{\alpha-1} - (1-p_1)^{-1/\alpha}}{(1-p_1)^{-1/\alpha} - (1-p_2)^{-1/\alpha}} = -\frac{\ln(1-p_1)+1}{\ln(1-p_1)-\ln(1-p_2)}$, the d value for the Pareto distribution approaches that of the exponential distribution as $\alpha \rightarrow \infty$, regardless of the type of weighted average used. This completes the demonstration. \square

Theorem .1 implies that for the Weibull, gamma, Pareto, lognormal and generalized Gaussian distribution, $rm_{d \approx 0.375, \epsilon = \frac{1}{8}}$ is consistent for at least one particular case. The biases of $rm_{d \approx 0.375, \epsilon = \frac{1}{8}}$ for distributions with skewness between those of the exponential and symmetric distributions are tiny (SI Dataset S1). $rm_{d \approx 0.375, \epsilon = \frac{1}{8}}$ exhibits excellent performance for all these common unimodal distributions (SI Dataset S1).

Besides introducing the concept of invariant mean, the purpose of this paper is to demonstrate that, in light of previous works, the estimation of central moments can be transformed into a location estimation problem by using U -statistics, the central moment kernel distributions possess desirable properties, and a series of sophisticated yet efficient robust estimators can be constructed whose biases are typically smaller than the variances (as seen in Table ?? for $n = 5400$) for unimodal distributions.

Background and Main Results

A. Invariant mean. It has long been known that a theoretical model can be adjusted to fit the first two moments of the observed data. A continuous distribution belonging to a location-scale family takes the form $F(x) = F_0\left(\frac{x-\mu}{\lambda}\right)$, where F_0 is a "standard" distribution. Therefore, $F(x) = Q^{-1}(x) \rightarrow x = Q(p) = \lambda Q_0(p) + \mu$. Thus, any weighted average can be expressed as $\lambda \text{WA}_0(\epsilon) + \mu$, where $\text{WA}_0(\epsilon)$ is an integral of $Q_0(p)$ according to the definition of the weighted average. The simultaneous cancellation of μ and λ in $\frac{(\lambda\mu_0 + \mu) - (\lambda\text{BM}_0(\epsilon) + \mu)}{(\lambda\text{BM}_0(\epsilon) + \mu) - (\lambda\text{median}_0 + \mu)}$ assures that d is a constant. Consequently, the roles of BM_{ϵ} and median in $rm_{d,\epsilon}$ can be replaced by any weighted averages, although only symmetric weighted averages are considered in defining the invariant mean.

The performance in heavy-tailed distributions can be further improved by constructing the quantile mean as

$$qm_{d,\epsilon,n} := \hat{Q}_n \left(\left(\hat{F}_n(\text{SWA}_{\epsilon,n}) - \frac{1}{2} \right) d + \hat{F}_n(\text{SWA}_{\epsilon,n}) \right),$$

provided that $\hat{F}_n(\text{SWA}_{\epsilon,n}) \geq \frac{1}{2}$, where $\hat{F}_n(x)$ is the empirical cumulative distribution function of the sample, \hat{Q}_n is the sample quantile function. The most popular method for computing the sample quantile function was proposed by Hyndman and Fan in 1996 (14). To minimize the finite sample bias, here, $\hat{F}_n(x) := \frac{1}{n} \left(\frac{x - X_{sp}}{X_{sp+1} - X_{sp}} + sp \right)$, where $sp = \sum_{i=1}^n \mathbf{1}_{X_i \leq x}$, $\mathbf{1}_A$ is the indicator of event A . The solution of $\hat{F}_n(\text{SWA}_{\epsilon,n}) < \frac{1}{2}$ is reversing the percentile by $1 - \hat{F}_n(\text{SWA}_{\epsilon,n})$, the obtained percentile is also reversed. Without loss of generality, in the following discussion, only the case where $\hat{F}_n(\text{SWA}_{\epsilon,n}) \geq \frac{1}{2}$ is considered. Moreover, in extreme heavy-tailed distributions, the calculated percentile can exceed the breakdown point of SWA_{ϵ} , so the percentile will be modified to $1 - \epsilon$ if this occurs. The quantile mean uses the location-scale invariant in a different way as shown in the following proof.

Theorem A.1. *$qm_{d \approx 0.321, \epsilon = \frac{1}{8}}$ is a consistent mean estimator for the exponential, Pareto ($\alpha \rightarrow \infty$) and any symmetric distributions provided that the second moments are finite.*

Proof. Similarly, rearranging the definition, $d = \frac{F(\mu) - F(\text{BM}_{\epsilon})}{F(\text{BM}_{\epsilon}) - \frac{1}{2}}$. The cdf of the exponential distribution is $F(x) = 1 - e^{-\lambda^{-1}x}$, $\lambda \geq 0$, $x \geq 0$, the expectation of BM_{ϵ} can be expressed as $\lambda \text{BM}_0(\epsilon)$, so $F(\text{BM}_{\epsilon})$ is free of λ . When $\epsilon = \frac{1}{8}$, $d = \frac{-e^{-1} + e^{-\left(1 + \ln\left(\frac{46656}{8575\sqrt{35}}\right)\right)}}{\frac{1}{2} - e^{-\left(1 + \ln\left(\frac{46656}{8575\sqrt{35}}\right)\right)}} \approx 0.321$. The proof of the symmetric case is similar. Since for any symmetric distribution with a finite second moment, $F(E[\text{BM}_{\epsilon}]) = F(\mu) = \frac{1}{2}$. Then, the expectation of the quantile mean is $qm_{d,\epsilon} = F^{-1}\left(\left(F(\mu) - \frac{1}{2}\right)d + F(\mu)\right) = F^{-1}\left(0 + F(\mu)\right) = \mu$.

For the assertion related to the Pareto distribution, the cdf of it is $1 - \left(\frac{x_m}{x}\right)^{\alpha}$. So, the d value with two unknown percentile p_1 and p_2 is

$$d_{\text{Pareto}} = \frac{1 - \left(\frac{x_m}{\frac{\alpha x_m}{\alpha-1}}\right)^{\alpha} - \left(1 - \left(\frac{x_m}{x_m(1-p_1)^{-\frac{1}{\alpha}}}\right)^{\alpha}\right)}{\left(1 - \left(\frac{x_m}{x_m(1-p_1)^{-\frac{1}{\alpha}}}\right)^{\alpha}\right) - \left(1 - \left(\frac{x_m}{x_m(1-p_2)^{-\frac{1}{\alpha}}}\right)^{\alpha}\right)} = \frac{1 - \left(\frac{\alpha-1}{\alpha}\right)^{\alpha-p_1}}{p_1 - p_2}. \text{ When } \alpha \rightarrow \infty, \left(\frac{\alpha-1}{\alpha}\right)^{\alpha} = \frac{1}{e}. \text{ The } d \text{ value for the exponential distribution is identical, since } d_{\text{exp}} =$$

$$\frac{(1-e^{-1}) - \left(1-e^{-\ln\left(\frac{1}{1-p_1}\right)}\right)}{\left(1-e^{-\ln\left(\frac{1}{1-p_1}\right)}\right) - \left(1-e^{-\ln\left(\frac{1}{1-p_2}\right)}\right)} = \frac{1-\frac{1}{e}-p_1}{p_1-p_2}. \quad \text{All results are now proven.} \quad \square$$

The definitions of location and scale parameters are such that they must satisfy $F(x; \lambda, \mu) = F\left(\frac{x-\mu}{\lambda}; 1, 0\right)$. By recalling $x = \lambda Q_0(p) + \mu$, it follows that the percentile of any weighted average is free of λ and μ , which guarantees the validity of the quantile mean. The quantile mean is a quantile I -statistic. Specifically, an estimator is classified as a quantile I -statistic if LEs are percentiles of a distribution obtained by plugging L -statistics into a cumulative distribution function and I is defined with arithmetic operations, constants and quantile functions. $qm_{d \approx 0.321, \epsilon = \frac{1}{8}}$ works better in the fat-tail scenarios (SI Dataset S1). Theorem .1 and A.1 show that $rm_{d \approx 0.375, \epsilon = \frac{1}{8}}$ and $qm_{d \approx 0.321, \epsilon = \frac{1}{8}}$ are both consistent mean estimators for any symmetric distribution and a skewed distribution with finite second moments. It's obvious that the breakdown points of $rm_{d \approx 0.375, \epsilon = \frac{1}{8}}$ and $qm_{d \approx 0.321, \epsilon = \frac{1}{8}}$ are both $\frac{1}{8}$. Therefore they are all invariant means.

To study the impact of the choice of SWAs in rm and qm , it is constructive to recall that a symmetric weighted average is a linear combination of symmetric quantile averages. While using a less-biased symmetric weighted average can generally enhance performance (SI Dataset S1), there is a greater risk of violation in the semiparametric framework. However, the mean-SWA-median inequality is robust to slight fluctuations of the SQA function of the underlying distribution. Suppose the SQA function is generally decreasing in $[0, u]$, but increasing in $[u, \frac{1}{2}]$, since $1-2\epsilon$ of the symmetric quantile averages will be included in the computation of SWA_ϵ , as long as $\frac{1}{2} - u \ll 1-2\epsilon$, and other portions of the SQA function satisfy the inequality constraints that define the ν th orderliness on which the SWA_ϵ is based, the mean-SWA-median inequality will still hold. This is due to the violation being bounded (15) and therefore cannot be extreme for unimodal distributions. For instance, the SQA function is non-monotonic when the shape parameter of the Weibull distribution $\alpha > \frac{1}{1-\ln(2)} \approx 3.259$ as shown in the previous article, the violation of the third orderliness starts near this parameter as well, yet the mean-BM $_{\frac{1}{8}}$ -median inequality is still valid when $\alpha \leq 3.322$. Another key factor in determining the risk of violation is the skewness of the distribution. Previously, it was demonstrated that in a family of distributions differing by a skewness-increasing transformation in van Zwet's sense, the violation of orderliness, if it happens, often only occurs when the distribution is nearly symmetrical (16). The over-corrections in rm and qm are dependent on the SWA_ϵ -median difference, which can be a reasonable measure of skewness (17, 18), implying that the over-correction is often tiny with a moderate d . This qualitative analysis provides another perspective, in addition to the bias bounds (15), that rm and qm based on the mean-SWA-median inequality are generally safe.

B. Robust estimations of the central moments. In 1979, Bickel and Lehmann, in their final paper of the landmark series *Descriptive Statistics for Nonparametric Models* (19), generalized a class of estimators called "measures of spread," which "does not require the assumption of symmetry." From that, a popular

efficient scale estimator, the Rousseeuw-Croux scale estimator (20), was derived in 1993, but the importance of tackling the symmetry assumption has been greatly underestimated. While they had already considered one version of the trimmed standard deviation in the third paper of that series (21), in the final section of that paper (19), they explored another two possible versions, which were modified here for comparison,

$$\left[n\left(\frac{1}{2} - \epsilon\right)\right]^{-\frac{1}{2}} \left[\sum_{i=\frac{n}{2}}^{n(1-\epsilon)} [X_i - X_{n-i+1}]^2 \right]^{\frac{1}{2}}, \quad [1]$$

and

$$\left[\binom{n}{2} (1 - \epsilon - \gamma\epsilon)\right]^{-\frac{1}{2}} \left[\sum_{i=\binom{n}{2}\gamma\epsilon}^{\binom{n}{2}(1-\epsilon)} (X - X')_i^2 \right]^{\frac{1}{2}}, \quad [2]$$

where $(X - X')_1 \leq \dots \leq (X - X')_{\binom{n}{2}}$ are the order statistics of the "pseudo-sample", $X_i - X_j$, $i < j$. The paper ended with, "We do not know a fortiori which of the measures [1] or [2] is preferable and leave these interesting questions open."

Observe that the kernel of the unbiased estimation of the second central moment by using U -statistic is $\psi_2(x_1, x_2) = \frac{1}{2}(x_1 - x_2)^2$. If adding the $\frac{1}{2}$ term in [2], as $\epsilon \rightarrow 0$, the result is equivalent to the standard deviation estimated by using U -statistic (also noted by Janssen, Serfling, and Veraverbeke in 1987) (22). In fact, they also showed that, when ϵ is 0, [2] is $\sqrt{2}$ times the standard deviation.

To address their open question, the nomenclature used in this paper is introduced as follows:

Nomenclature. Given a robust estimator $\hat{\theta}$ with an adjustable breakdown point which can be infinitesimal. The name of $\hat{\theta}$ is composed of two parts: the first part denotes the type of estimator, and the second part is the name of the population parameter θ that the estimator is consistent with as $\epsilon \rightarrow 0$. The abbreviation of the estimator is formed by combining the initial letter(s) of the first part with the common abbreviation of the consistent estimator that measures the population parameter. If the estimator is symmetric, the asymptotic breakdown point, ϵ , is indicated in the subscript of the abbreviation of the estimator, except the median. For asymmetric estimators based on quantile average, the corresponding γ is also indicated after ϵ . Note that ϵ is the right breakdown point (defined in Subsection ??), while the left breakdown point should be further calculated.

In the previous article on semiparametric robust mean estimation, it was shown that the bias of a robust estimator with an adjustable breakdown point is often monotonic with respect to the breakdown point in a semiparametric distribution. Naturally, the estimator's name should correspond to the population parameter with which it is consistent as $\epsilon \rightarrow 0$. The trimmed standard deviation following this nomenclature

$$\text{is } Tsd_{\epsilon=1-\sqrt{1-\epsilon_0}, \gamma, n} := \left[TM_{\epsilon_0, \gamma} \left((\psi_2(X_{N_1}, X_{N_2}))_{N=1}^{\binom{n}{2}} \right) \right]^{-\frac{1}{2}},$$

where $TM_{\epsilon_0, \gamma}(Y)$ denotes the ϵ_0, γ -trimmed mean with the sequence $(\psi_2(X_{N_1}, X_{N_2}))_{N=1}^{\binom{n}{2}}$ as an input, the proof of the breakdown point is given in Subsection ??. Removing the square root yields the trimmed variance ($Tvar_{\epsilon, \gamma, n}$). It is now very clear that this definition, essentially the same as

[2], should be preferable. Not only because it is essentially a trimmed U -statistic for the standard deviation but also because the γ -orderliness of the second central moment kernel distribution is ensured by the next exciting theorem.

Theorem B.1. *The second central moment kernel distribution generated from any unimodal distribution is γ -ordered.*

Proof. The monotonic increasing of the pairwise difference distribution was first implied in its unimodality proof done by Hodges and Lehmann in 1954 (23). Whereas they used induction to get the result in Theorem ??, Dharmadhikari and Jogdeo in 1982 (24) provided a modern proof of the unimodality using Khintchine's representation (25). Assuming absolute continuity, Purkayastha (26) introduced a much simpler proof in 1998. Transforming the pairwise difference distribution by squaring and multiplying by $\frac{1}{2}$ does not change the monotonicity, making the pdf become monotonically decreasing with mode at zero. In the previous article, it was proven that a right skewed distribution with a monotonic decreasing pdf is always γ -ordered, which gives the desired result. \square

Previously, it was shown that any symmetric distribution with a finite second moment is ν th ordered, indicating that orderliness does not require unimodality, e.g., a symmetric bimodal distribution is also ordered. An analysis of the Weibull distribution showed that unimodality does not guarantee orderliness. Theorem B.1 reveals another profound relationship between unimodality and orderliness, which is sufficient for trimming inequality.

In 1928, Fisher constructed k -statistics as unbiased estimators of cumulants (27). Halmos (1946) proved that the functional θ admits an unbiased estimator if and only if it is a regular statistical functional of degree k and showed a relation of symmetry, unbiasedness and minimum variance (28). In 1948, Hoeffding generalized U -statistics (29) which enable the derivation of a minimum-variance unbiased estimator from each unbiased estimator of an estimable parameter. Hoeffman (1997) (30) obtained an unbiased estimator of the k th central moment by using U -statistics and demonstrated that it is the minimum variance unbiased estimator for distributions with finite moments (31, 32). In 1976, Saleh generalized the Hodges-Lehmann (H-L) estimator (33) to the trimmed H-L mean (34). In 1984, Serfling pointed out the speciality of Hodges-Lehmann estimator, which is neither a simple L -statistic nor a U -statistic, and considered the generalized L -statistics and U -statistic structure (35). Also in 1984, Janssen and Serfling and Veraverbeke (36) showed that the Bickel-Lehmann spread also belongs to the same class. It gradually became clear that the Hodges-Lehmann estimator, trimmed H-L mean and trimmed standard deviation are all trimmed U -statistics (37–39).

Extending the trimmed U -statistic to weighted U -statistic, i.e., replacing the trimmed mean with weighted average. The weighted k th central moment ($k \leq n$) is defined as,

$$Wkm_{\epsilon=1-(1-\epsilon_0)\frac{1}{k}, \gamma, n} := WA_{\epsilon_0, \gamma, n} \left((\psi_k(X_{N_1}, \dots, X_{N_k}))_{N=1}^n \right),$$

where X_{N_1}, \dots, X_{N_k} are the n choose k elements from X , $\psi_k(x_1, \dots, x_k) = \sum_{j=0}^{k-2} (-1)^j \left(\frac{1}{k-j} \right) \sum (x_{i_1}^{k-j} \dots x_{i_{j+1}}) + (-1)^{k-1} (k-1) x_1 \dots x_k$, the second summation is over $i_1, \dots, i_{j+1} = 1$ to k with $i_1 < \dots < i_{j+1}$ (30). Despite the

complexity, the structure of the k th central moment kernel distributions can be elucidated by decomposing.

Theorem B.2. *For each pair $(Q(p_i), Q(p_j))$ of the original distribution such that $Q(p_i) < Q(p_j)$, let $x_1 = Q(p_i)$ and $x_k = Q(p_j)$, $\Delta = Q(p_i) - Q(p_j)$, the k th central moment kernel distribution, $k > 2$, can be seen as a mixture distribution and each of the components has the support $(-\left(\frac{k}{3+(-1)^k}\right)^{-1}(-\Delta)^k, \frac{1}{k}(-\Delta)^k)$.*

Proof. Without loss of generality, generating the distribution of the k -tuple $(Q(p_{i_1}), \dots, Q(p_{i_k}))$ under continuity, $k > 2$, $i_1 < \dots < i_k$, $p_{i_1} < \dots < p_{i_k}$, the corresponding probability density is $f_{X, \dots, X}(Q(p_{i_1}), \dots, Q(p_{i_k})) = k! f(Q(p_{i_1})) \dots f(Q(p_{i_k}))$. Transforming the distribution of the k -tuple by the function $\psi_k(x_1, \dots, x_k)$, denoting $\bar{\Delta} = \psi_k(Q(p_{i_1}), \dots, Q(p_{i_k}))$. The probability $f_{\Xi_k}(\bar{\Delta}) = \sum_{\bar{\Delta}=\psi_k(Q(p_{i_1}), \dots, Q(p_{i_k}))} f_{X, \dots, X}(Q(p_{i_1}), \dots, Q(p_{i_k}))$ is the summation of the probabilities of all k -tuples such that $\bar{\Delta}$ is equal to $\psi_k(Q(p_{i_1}), \dots, Q(p_{i_k}))$. The following Ξ_k is equivalent.

Ξ_k : Every pair with a difference equal to $\Delta = Q(p_{i_1}) - Q(p_{i_k})$ can generate a pseudodistribution (but the integral is not equal to 1, so "pseudo") such that x_2, \dots, x_{k-1} exhaust all combinations under the inequality constraints, i.e., $Q(p_{i_1}) = x_1 < x_2 < \dots < x_{k-1} < x_k = Q(p_{i_k})$. The combination of all the pseudodistributions with the same Δ is ξ_Δ . The combination of ξ_Δ , i.e., from $\Delta = 0$ to $Q(0) - Q(1)$, is Ξ_k .

The support of ξ_Δ is the extrema of ψ_k subject to the inequality constraints. Using the Lagrange multiplier, one can easily determine the only critical point at $x_1 = \dots = x_k = 0$, where $\psi_k = 0$. Other candidates are within the boundaries, i.e., $\psi_k(x_1 = x_1, x_2 = x_k, \dots, x_k = x_k)$, \dots , $\psi_k(x_1 = x_1, \dots, x_i = x_1, x_{i+1} = x_k, \dots, x_k = x_k)$, \dots , $\psi_k(x_1 = x_1, \dots, x_{k-1} = x_1, x_k = x_k)$. $\psi_k(x_1 = x_1, \dots, x_i = x_1, x_{i+1} = x_k, \dots, x_k = x_k)$ can be divided into k groups. If $\frac{k+1-i}{2} \leq j \leq \frac{k-1}{2}$, from $j+1$ st to $k-j$ th group, the g th group has $i \binom{i-1}{g-j-1} \binom{k-i}{j}$ terms having the form $(-1)^{g+1} \frac{1}{k-g+1} x_1^{k-j} x_k^j$, from $k-j+1$ th to $i+j$ th group, the g th group has $i \binom{i-1}{g-j-1} \binom{k-i}{j} + (k-i) \binom{k-i-1}{j-k+g-1} \binom{i}{k-j}$ terms having the form $(-1)^{g+1} \frac{1}{k-g+1} x_1^{k-j} x_k^j$. If $j < \frac{k+1-i}{2}$, from $j+1$ st to $i+j$ th group, the g th group has $i \binom{i-1}{g-j-1} \binom{k-i}{j}$ terms having the form $(-1)^{g+1} \frac{1}{k-g+1} x_1^{k-j} x_k^j$. If $j \geq \frac{k}{2}$, from $k-j+1$ st to j th group, the g th group has $(k-i) \binom{k-i-1}{j-k+g-1} \binom{i}{k-j}$ terms having the form $(-1)^{g+1} \frac{1}{k-g+1} x_1^{k-j} x_k^j$, from $j+1$ th to $j+i$ th group, $i+j < k$, the g th group has $i \binom{i-1}{g-j-1} \binom{k-i}{j} + (k-i) \binom{k-i-1}{j-k+g-1} \binom{i}{k-j}$ terms having the form $(-1)^{g+1} \frac{1}{k-g+1} x_1^{k-j} x_k^j$. The final k th group is the term $(-1)^{k-1} (k-1) x_1^{k-i} x_k^i$. So, if $i+j = k$, $j \geq \frac{k}{2}$, $i \leq \frac{k}{2}$, the summed coefficient of $x_1^{k-i} x_k^i$ is $(-1)^{k-1} (k-1) + \sum_{g=i+1}^{k-1} (-1)^{g+1} \frac{1}{k-g+1} (k-i) \binom{k-i-1}{g-i-1} + \sum_{g=k-i+1}^{k-1} (-1)^{g+1} \frac{1}{k-g+1} i \binom{i-1}{g-k+i-1} = (-1)^{k-1} (k-1) + (-1)^{k+1} + (k-i) (-1)^k + (-1)^k (i-1) = (-1)^{k+1}$. The summation identities are $\sum_{g=i+1}^{k-1} (-1)^{g+1} \frac{1}{k-g+1} (k-i) \binom{k-i-1}{g-i-1} = (k-i) \int_0^1 \sum_{g=i+1}^{k-1} (-1)^{g+1} \binom{k-i-1}{g-i-1} t^{k-g} dt = (k-i) \int_0^1 ((-1)^i (t-1)^{k-i-1} - (-1)^{k+1}) dt$

370 $(k-i) \binom{(-1)^k}{i-k} + (-1)^k = (-1)^{k+1} + (k-i)(-1)^k$
 371 and $\sum_{g=k-i+1}^{k-1} (-1)^{g+1} \frac{1}{k-g+1} i \binom{i-1}{g-k+i-1} =$
 372 $\int_0^1 \sum_{g=k-i+1}^{k-1} (-1)^{g+1} i \binom{i-1}{g-k+i-1} t^{k-g} dt =$
 373 $\int_0^1 (i(-1)^{k-i} (t-1)^{i-1} - i(-1)^{k+1}) dt = (-1)^k (i-1).$
 374 If $j < \frac{k+1-i}{2}$, $i > k-1$, if $i = k$, $\psi_k = 0$, if $\frac{k+1-i}{2} \leq j \leq \frac{k-1}{2}$,
 375 $\frac{k+1}{2} \leq i \leq k-1$, the summed coefficient of $x_1^i x_k^{k-i}$ is
 376 $(-1)^{k-1} (k-1) + \sum_{g=k-i+1}^{k-1} (-1)^{g+1} \frac{1}{k-g+1} i \binom{i-1}{g-k+i-1} +$
 377 $\sum_{g=i+1}^{k-1} (-1)^{g+1} \frac{1}{k-g+1} (k-i) \binom{k-i-1}{g-i-1}$, the same as above. If
 378 $i+j < k$, since $\binom{i}{k-j} = 0$, the related terms can be ignored, so,
 379 using the binomial theorem and beta function, the summed co-
 380 efficient of $x_1^{k-j} x_k^j$ is $\sum_{g=j+1}^{i+j} (-1)^{g+1} \frac{1}{k-g+1} i \binom{i-1}{g-j-1} \binom{k-i}{j} =$
 381 $i \binom{k-i}{j} \int_0^1 \sum_{g=j+1}^{i+j} (-1)^{g+1} \binom{i-1}{g-j-1} t^{k-g} dt =$
 382 $\binom{k-i}{j} i \int_0^1 \left((-1)^j t^{k-j-1} \left(\frac{t}{1-t} \right)^{1-i} \right) dt =$
 383 $\binom{k-i}{j} i \frac{(-1)^{j+i+1} \Gamma(i) \Gamma(k-j-i+1)}{\Gamma(k-j+1)} = \frac{(-1)^{j+i+1} i! (k-j-i)! (k-i)!}{(k-j)! j! (k-j-i)!} =$
 384 $(-1)^{j+i+1} \frac{i! (k-i)!}{k!} \frac{k!}{(k-j)! j!} = \binom{k}{i}^{-1} (-1)^{1+i} \binom{k}{j} (-1)^j.$
 385 The coefficient of $x_1^i x_k^{k-i}$ in $\binom{k}{i}^{-1} (-1)^{1+i} (x_1 - x_k)^k$
 386 is $\binom{k}{i}^{-1} (-1)^{1+i} \binom{k}{i} (-1)^{k-i} = (-1)^{k+1}$, same as the
 387 summed coefficient if $i+j = k$. If $i+j < k$,
 388 the coefficient of $x_1^{k-j} x_k^j$ is $\binom{k}{i}^{-1} (-1)^{1+i} \binom{k}{j} (-1)^j$,
 389 same as the corresponding summed coefficient. There-
 390 fore, $\psi_k(x_1 = x_1, \dots, x_i = x_1, x_{i+1} = x_k, \dots, x_k = x_k) =$
 391 $\binom{k}{i}^{-1} (-1)^{1+i} (x_1 - x_k)^k$, the maximum and minimum of ψ_k
 392 follow directly from the properties of the binomial coeffi-
 393 cient. \square

394 ξ_Δ is closely related to $f_\Xi(\Delta)$, which is the pairwise differ-
 395 ence distribution, since the probability density of ξ_Δ can be ex-
 396 pressed as $f_{\Xi_k}(\bar{\Delta}|\Delta) = \sum_{\bar{\Delta} = -(\frac{k}{3+(-1)^k})}^{\frac{1}{k}(-\Delta)^k} f_{\Xi_k}(\bar{\Delta}|\Delta) =$
 397 $f_\Xi(\Delta) = \int_0^\infty 2f(t)f(t-\Delta)dt$. The support of the original
 398 distribution is assumed to be $[0, \infty)$ for simplicity. Recall that
 399 $f_\Xi(\Delta)$ is monotonic increasing with a mode at the origin if the
 400 original distribution is unimodal. Thus, in general, ignoring
 401 the shape of ξ_Δ , Ξ_k is monotonic left and right around zero.
 402 In fact, the median of Ξ_k also exhibits a strong tendency to be
 403 close to zero, as it can be cast as a weighted mean of the medi-
 404 ans of ξ_Δ . When Δ is small, all values of ξ_Δ are close to zero,
 405 resulting in the median of ξ_Δ being close to zero as well. When
 406 Δ is large, the median of ξ_Δ depends on its skewness, but the
 407 corresponding weight is much smaller, so even if ξ_Δ is highly
 408 skewed, the median of Ξ_k will only be slightly shifted from
 409 zero. Denote the median of Ξ_k as m_{Ξ_k} , for the five parametric
 410 distributions here, $|m_{\Xi_k}|$ s are all $\leq 0.1\sigma$ for Ξ_3 and Ξ_4 (SI
 411 Dataset S1). Assuming $m_{\Xi_k} = 0$, for the even ordinal central
 412 moment kernel distribution, the average probability density on
 413 the left side of zero is greater than that on the right side, since
 414 $\frac{\frac{1}{2}}{\binom{k}{2}^{-1} (Q(0)-Q(1))^k} > \frac{\frac{1}{2}}{\frac{1}{k} (Q(0)-Q(1))^k}$. This means that, on aver-
 415 age, the inequality $f(Q(\epsilon)) \geq f(Q(1-\epsilon))$ holds. For the odd
 416 ordinal distribution, the discussion is more challenging since
 417 it is generally symmetric. Just consider Ξ_3 , let $x_1 = Q(p_i)$
 418 and $x_3 = Q(p_j)$, changing the value of x_2 from $Q(p_i)$ to
 419 $Q(p_j)$ will monotonically change the value of $\psi_3(x_1, x_2, x_3)$,
 420 since $\frac{\partial \psi_3(x_1, x_2, x_3)}{\partial x_2} = -\frac{x_2}{2} - x_1 x_2 + 2x_1 x_3 + x_2^2 - x_2 x_3 - \frac{x_2^2}{2}$,
 421 $-\frac{3}{4}(x_1 - x_3)^2 \leq \frac{\partial \psi_3(x_1, x_2, x_3)}{\partial x_2} \leq -\frac{1}{2}(x_1 - x_3)^2 \leq 0$. If the
 422 original distribution is right-skewed, ξ_Δ will be left-skewed,

423 so, for Ξ_3 , the average probability density of the right side of
 424 zero will be greater than that of the left side, which means,
 425 on average, the inequality $f(Q(\epsilon)) \leq f(Q(1-\epsilon))$ holds (the
 426 same result can be inferred from the definition of central mo-
 427 ments, where the positivity of the odd order central moment
 428 is directly related to the left-skewness of the corresponding
 429 kernel distribution). In all, the monotonicity of the pairwise
 430 difference distribution guides the general shape of the k th
 431 central moment kernel distribution, $k > 2$, forcing it to be
 432 unimodal-like with the mode and median close to zero, then,
 433 the inequality $f(Q(\epsilon)) \leq f(Q(1-\epsilon))$ or $f(Q(\epsilon)) \geq f(Q(1-\epsilon))$
 434 holds in general. If a distribution is ordered and all of its cen-
 435 tral moment kernel distributions are also ordered, it is called
 436 completely ordered. Although strict complete orderliness is
 437 difficult to prove, even if the inequality may be violated in
 438 a small range, as discussed in Subsection A, the mean-SWA-
 439 median inequality remains valid, in most cases, for the central
 440 moment kernel distribution.

441 Another crucial property of the central moment kernel dis-
 442 tribution, location invariant, is introduced in the next theorem.
 443 The proof is provided in the SI Text.

Theorem B.3. $\psi_k(x_1 = \lambda x_1 + \mu, \dots, x_k = \lambda x_k + \mu) =$
 444 $\lambda^k \psi_k(x_1, \dots, x_k).$ 445

A direct result of Theorem B.3 is that, Wkm after stan-
 dardization is invariant to location and scale. So, the weighted
 standardized k th moment is defined to be

$$Wskm_{\epsilon, \gamma, n} := \frac{Wkm_{\epsilon, \gamma, n}}{Wvar_{\epsilon, \gamma, n}^{\frac{k}{2}}}.$$

Consider two continuous distributions belonging to the
 same location-scale family, their corresponding k th central
 moment kernel distributions only differ in scaling. So d is
 invariant, as shown in Subsection A. The recombined k th
 central moment, based on rm , is defined by,

$$rkm_{d, \epsilon=1-(1-\epsilon_0)^{\frac{1}{k}}, n} := (d+1)SWkm_{\epsilon, n} - dmkmn,$$

where $SWkm_{\epsilon, n}$ is using the binomial k th central moment
 ($Bkm_{\epsilon_0, n}$) here, $mkmn$ is the median k th central moment.
 Since $SWkm_{\epsilon, n}$ is an L -statistic, the resulting $rkm_{d, \epsilon, n}$ is an
 arithmetic I -statistic. Similarly, the quantile will not change
 after scaling. The quantile k th central moment is thus defined as

$$qkm_{d, \epsilon, n} := \hat{Q}_n \left(\left(pSWkm_{\epsilon, n} - \frac{1}{2} \right) d + pSWkm_{\epsilon, n} \right),$$

where $pSWkm_{\epsilon, n} = \hat{F}_{\psi, n}(SWkm_{\epsilon, n})$, $\hat{F}_{\psi, n}$ is the empirical
 cumulative distribution function of the corresponding central
 moment kernel distribution. $qkm_{d, \epsilon, n}$ is a quantile I -statistic.

For standardized moments, quantile skewness and quan-
 tile kurtosis are defined to be $qskew_{d, \epsilon, n} := \frac{qtm_{d, \epsilon, n}}{qsd_{d, \epsilon, n}^{\frac{3}{2}}}$ and
 449 $qkurt_{d, \epsilon, n} := \frac{qfm_{d, \epsilon, n}}{qsd_{d, \epsilon, n}^4}$. Quantile standard deviation ($qsd_{d, \epsilon, n}$),
 450 recombined standard deviation ($rsd_{d, \epsilon, n}$), quantile third cen-
 451 tral moment ($qtm_{d, \epsilon, n}$), quantile fourth central moment
 452 ($qfm_{d, \epsilon, n}$), recombined third central moment ($rtm_{d, \epsilon, n}$), re-
 453 combined fourth central moment ($rftm_{d, \epsilon, n}$), recombined skew-
 454 ness ($rskew_{d, \epsilon, n}$), and recombined kurtosis ($rkurt_{d, \epsilon, n}$) are all
 455 defined similarly as above and not repeated here. The trans-
 456 formation to a location problem can also empower related
 457
 458

statistical tests. From the better performance of the quantile mean in heavy-tailed distributions, quantile central moments are generally better than recombined central moments regarding asymptotic bias.

To avoid confusion, it should be noted that the robust location estimations of the kernel distributions discussed in this paper differ from the approach taken by Joly and Lugosi (2016) (40), which is computing the median of all U -statistics from different disjoint blocks. Compared to bootstrap median U -statistics, this approach can produce two additional kinds of finite sample bias, one arises from the limited numbers of blocks, another is due to the size of the U -statistics (consider the mean of all U -statistics from different disjoint blocks, it is definitely not identical to the original U -statistic, except when the kernel is the Hodges-Lehmann kernel). Laforgue, Clemencon, and Bertail (2019)'s median of randomized U -statistics (41) is more sophisticated and can overcome the limitation of the number of blocks, but the second kind of bias remains unsolved.

C. Congruent distribution. In the realm of nonparametric statistics, the precise values of robust estimators are of secondary importance. What is of primary importance is their relative differences or orders. Based on this principle, in the absence of contamination, as the parameters of the distribution vary, all reasonable nonparametric location estimates should asymptotically change in the same direction. Otherwise if the results obtained based on the trimmed mean are completely different from those based on the median, a contradiction arises. However, such contradictions are possible, as in the case of the Weibull distribution, $E[m] = \lambda \sqrt[\gamma]{\ln(2)}$, $E[\mu] = \lambda \Gamma(1 + \frac{1}{\alpha})$, then, when $\alpha = 1$, $E[m] = \lambda \ln(2) \approx 0.693\lambda$, $E[\mu] = \lambda$, but when $\alpha = \frac{1}{2}$, $E[m] = \lambda \ln^2(2) \approx 0.480\lambda$, $E[\mu] = 2\lambda$, the mean increases, but the median decreases. To study the conditions that avoid such scenarios by classifying distributions through the signs of derivatives, let the quantile average function of a parametric distribution be denoted as $QA(\epsilon, \gamma, \alpha_1, \dots, \alpha_i, \dots, \alpha_k)$, where α_i represent the parameters of the distribution, then, a distribution is γ -congruent if and only if the sign of $\frac{\partial QA}{\partial \alpha_i}$ remains the same for all $0 \leq \epsilon \leq \frac{1}{1+\gamma}$. If this partial derivative is equal to zero or undefined, it can be considered both positive and negative, and thus does not impact the analysis. Asymptotically, any weighted average can be expressed as an integral of the quantile average function. Since the sign does not change after integration, the sign of $\frac{\partial QA}{\partial \alpha_i}$ remains the same for all $0 \leq \epsilon \leq \frac{1}{1+\gamma}$ implies that all γ -weighted averages change in the same direction as the parameters change, as long as they are not undefined. A distribution is completely γ -congruent if and only if it is γ -congruent and all its central moment kernel distributions are also γ -congruent. Setting $\gamma = 1$ constitutes the definitions of congruence and complete congruence. Chebyshev's inequality implies that, for any probability distribution with finite moments, even if some weighted averages change in a direction different from that of the sample mean, the deviations are bounded. Furthermore, distributions with infinite moments can be γ -congruent, since the definition is based on the quantile average, not the sample mean.

The following theorems show the conditions that a distribution is congruent or γ -congruent.

Theorem C.1. *A symmetric distribution with a finite second*

moment is always congruent.

Proof. For any symmetric distribution with a finite second moment, all symmetric quantile averages coincide. The conclusion follows immediately. \square

Theorem C.2. *A positive define location-scale distribution with a finite second moment is always γ -congruent.*

Proof. As shown in discussions in Subsection A, for a location-scale distribution, any weighted average can be expressed as $\lambda WA_0(\epsilon) + \mu$, where $WA_0(\epsilon)$ is an integral of $Q_0(p)$ according to the definition of the weighted average. Therefore, the derivatives with respect to the parameters λ or μ are always positive. By application of the definition, the desired outcome is obtained. \square

Theorem C.3. *The second central moment kernel distribution derived from a continuous location-scale unimodal distribution with a finite second moment is always γ -congruent.*

Proof. Theorem B.3 shows that the corresponding central moment kernel distribution is also a location-scale family distribution. Theorem B.1 shows that it is positively defined. Implementing Theorem C.2 yields the desired result. \square

For the Pareto distribution, $\frac{\partial Q(p, \alpha)}{\partial \alpha} = \frac{x_m(1-p)^{-1/\alpha} \ln(1-p)}{\alpha^2}$. Since $\ln(1-p) < 0$ for all $0 < p < 1$, $(1-p)^{-1/\alpha} > 0$ for all $0 < p < 1$ and $\alpha > 0$, so $\frac{\partial Q(p, \alpha)}{\partial \alpha} < 0$, and therefore $\frac{\partial QA(\epsilon, \gamma, \alpha)}{\partial \alpha} < 0$, the Pareto distribution is γ -congruent. The derivative for the lognormal distribution is $\frac{\partial SQA(\epsilon, \sigma)}{\partial \sigma} = \frac{-\operatorname{erfc}^{-1}(2\epsilon)e^{\mu - \sqrt{2}\sigma \operatorname{erfc}^{-1}(2\epsilon)} - \operatorname{erfc}^{-1}(2-2\epsilon)e^{\mu - \sqrt{2}\sigma \operatorname{erfc}^{-1}(2-2\epsilon)}}{\sqrt{2}}$. Since the inverse complementary error function is positive when the input is smaller than 1, and negative when the input is larger than 1, $\operatorname{erfc}^{-1}(2\epsilon) = -\operatorname{erfc}^{-1}(2-2\epsilon)$, $e^{\mu - \sqrt{2}\sigma \operatorname{erfc}^{-1}(2-2\epsilon)} > e^{\mu - \sqrt{2}\sigma \operatorname{erfc}^{-1}(2\epsilon)}$, $\frac{\partial SQA(\epsilon, \sigma)}{\partial \sigma} > 0$, the lognormal distribution is congruent. Theorem C.1 implies that the generalized Gaussian distribution is congruent. For the Weibull distribution, when α changes from 1 to $\frac{1}{2}$, the average probability density on the left side of the median increases, since $\frac{1}{\lambda \ln(2)} < \frac{1}{\lambda \ln^2(2)}$, but the mean increases, indicating that the distribution is more heavy-tailed, the probability density of large values will also increase. The main reason for non-congruence of a right-skewed smooth partial bounded probability distribution lies in the simultaneous increase of probability densities on two opposite sides: one approaching the bound and the other approaching infinity. Note that the gamma distribution does not have this issue, it looks to be congruent.

Although some common parametric distributions are not congruent, Theorem C.2 establishes that γ -congruence always holds for a positive define location-scale family distribution and thus for the second central moment kernel distribution generated from a continuous location-scale unimodal distribution as shown in Theorem C.3. Theorem B.2 demonstrates that all their central moment kernel distributions are unimodal-like with mode and median close to zero, as long as they are unimodal distributions. Assuming finite moments and constant $Q(0) - Q(1)$, increasing the mean of the kernel distribution will result in a more heavy-tailed distribution, i.e., the probability density of the values close to $\frac{1}{k}(-\Delta)^k$ increases. While the total probability density on either side of zero remains

unchanged as the median is generally close to zero and much less impacted by increasing the mean, the probability density of the values close to zero decreases. This transformation will increase nearly all symmetric weighted averages, in the general sense. Therefore, except for the median, which is assumed to be zero, nearly all symmetric weighted averages for all central moment kernel distributions derived from unimodal distributions should change in the same direction when the parameters change. Therefore, they are valid measures for nonparametric descriptive statistics.

D. A shape-scale distribution as the consistent distribution.

Up to this point, in this article, the consistent robust estimation has been limited to a location-scale distribution, with the location parameter often being omitted for simplicity. To construct probability distributions can be made to fit the observed skewness and kurtosis arbitrarily well, in 1894, Pearson (42) introduced a family of continuous probability distributions that are now often characterized by the square of the skewness and the kurtosis. If the skewness and the kurtosis are interrelated by a shape parameter, a distribution specified by a shape parameter (denoted as α) and a scale parameter (denoted as λ) is often referred to as a shape-scale distribution. Weibull, gamma, Pareto, lognormal, and generalized Gaussian distributions (when μ is a constant) are all shape-scale unimodal distributions. Moreover, if α or skewness or kurtosis is a constant, the shape-scale distribution is reduced to a location-scale distribution. The above discussion shows that, due to the invariant property, if a location-scale distribution is chosen as the consistent distribution, the type of invariant moments and their related weighted moments are given, there should exist a unique k -tuple (d_{im}, \dots, d_{ikm}) calibrated by the distribution and the corresponding kernel distributions generated from this distribution. For a right skewed shape-scale distribution, let $D(|skewness|, kurtosis, k, etype, dtype, n) = d_{ikm}$ denote these relations, where the first input is the absolute value of the skewness, the second input is the kurtosis, the third is the order of the central moment (if $k = 1$, the mean), the fourth is the type of estimator, the fifth is the type of consistent distribution, and the sixth input is the sample size. For simplicity, the last three inputs will be omitted in the following discussion. Hold in awareness that due to the invariant property of scale, specifying d values for a shape-scale distribution only requires either skewness or kurtosis, while the other may be also omitted. Since many common shape-scale distributions are always right skewed (if not, only the right skewed or left skewed part is used for calibration, while the other part is omitted), the absolute value of the skewness should be identical to the skewness for them and it can also handle the left skew scenario well.

For recombined moments, the object of using a shape-scale distribution as the consistent distribution is to find solutions for the system of equations

$$\begin{cases} rm(SWA, median, D(|rskew|, rkurt, 1)) = \mu \\ rvar(SWvar, mvar, D(|rskew|, rkurt, 2)) = \mu_2 \\ rtm(SWtm, mtm, D(|rskew|, rkurt, 3)) = \mu_3 \\ rfm(SWfm, mfm, D(|rskew|, rkurt, 4)) = \mu_4 \\ rskew = \frac{\mu_3}{\mu_2} \\ rkurt = \frac{\mu_4}{\mu_2^2} \end{cases},$$

where μ_2 , μ_3 and μ_4 are the population second,

third and fourth central moments. $|rskew|$ and $rkurt$ should be the invariant points of the functions $\varsigma(|rskew|) = \left| \frac{rtm(SWtm, mtm, D(|rskew|, 3))}{rvar(SWvar, mvar, D(|rskew|, 2))^{3/2}} \right|$ and $\varkappa(rkurt) = \frac{rfm(SWfm, mfm, D(rkurt, 4))}{rvar(SWvar, mvar, D(rkurt, 2))^2}$. Clearly, this is an overdetermined nonlinear system of equations, given that the skewness and kurtosis are interrelated for a shape-scale distribution. Since an overdetermined system constructed with random coefficients is almost always inconsistent, it is natural to optimize them separately using the fixed-point iteration (see Algorithm 1, only $rkurt$ is provided, others are the same).

Algorithm 1 $rkurt$ for a shape-scale distribution

Input: D ; $SWvar$; $SWfm$; $mvar$; mfm ; $maxit$; δ

Output: $rkurt_{i-1}$

```

1:  $i = 0$ 
2:  $rkurt_i \leftarrow \varkappa(kurtosis_{max})$   $\triangleright$  Using the maximum kurtosis
   available in  $D$  as an initial guess.
3: repeat
4:    $i = i + 1$ 
    $rkurt_{i-1} \leftarrow rkurt_i$ 
5:    $rkurt_i \leftarrow \varkappa(rkurt_{i-1})$ 
6: until  $i > maxit$  or  $|rkurt_i - rkurt_{i-1}| < \delta$   $\triangleright maxit$  is
   the maximum number of iterations,  $\delta$  is a small positive
   number.
```

The following theorem shows the validity of Algorithm 1.

Theorem D.1. Assuming $mkms$ are all equal to zero, $|rskew|$ and $rkurt$, defined as the largest attracting fix points of the functions $\varsigma(|rskew|)$ and $\varkappa(rkurt)$, are consistent estimators of $\tilde{\mu}_3$ and $\tilde{\mu}_4$ for a shape-scale distribution whose central moment kernel distributions are all congruent, as long as they are within the domain of D , where $\tilde{\mu}_3$ and $\tilde{\mu}_4$ are the population skewness and kurtosis.

Proof. Without loss of generality, only $rkurt$ is considered here, while the logic for $|rskew|$ is the same. Also, according to the property of invariance, the second central moments of the sample and consistent distribution are all assumed to be 1. From the definition of D , $\frac{\varkappa(rkurt_D)}{rkurt_D} = \frac{fm_D - SWfm_D}{SWfm_D - mfm_D} \frac{(SWfm - mfm) + SWfm}{(var_D - SWvar_D)(SWvar - mvar) + SWvar}$, where the subscript D indicates that the estimates are from the central moment kernel distributions generated from the consistent distribution used to calibrate the d values, while other estimates are from the sample.

Then, assuming the $mkms$ are all equal to zero, $\frac{\varkappa(rkurt_D)}{rkurt_D} = \frac{fm_D - SWfm_D}{SWfm_D} \frac{(SWfm) + SWfm}{(SWvar_D)^2} = \frac{(fm_D - SWfm_D + 1)(SWfm)}{fm_D (SWvar_D)^2} = \frac{SWfm SWvar_D^2}{SWfm_D SWvar_D^2} = \frac{SWkurt}{SWkurt_D}$. Since $SWfm_D$ are from the same kernel distribution as $fm_D = rkurt_D var_D^2$, according to the congruence, an increase in fm_D will also result in an increase in $SWfm_D$. Combining with Theorem B.3, $SWkurt$ is a measure of kurtosis that is invariant to location and scale, so $\lim_{rkurt_D \rightarrow \infty} \frac{\varkappa(rkurt_D)}{rkurt_D} < 1$. As a result, if there is at least one fix point, let the largest one be

666 fix_{max} , then it is attracting since $|\frac{\partial(\kappa(rkurt_D))}{\partial(rkurt_D)}| < 1$ for all
 667 $rkurt_D \in [fix_{max}, kurtosis_{max}]$.

668 Asymptotically, consider any $SWkurt_D > SWkurt$,
 669 $\frac{\kappa(rkurt_D)}{rkurt_D} < 1$, the same logic applies, a consistent estimator
 670 must be the last attracting fix point, fix_{max} is the consistent
 671 estimator. \square

672 As a result of Theorem D.1, assuming continuity, $mkms$
 673 are all equal to zero, and congruence of the central moment
 674 kernel distributions, Algorithm 1 converges surely provided
 675 that a fix point exists within the domain of D . At this stage,
 676 D can only be approximated through a Monte Carlo study.
 677 Continuity can be ensured by using linear interpolation. One
 678 common encountered problem is that the domain of D de-
 679 pends on both the consistent distribution and the Monte Carlo
 680 study, so the iteration may halt at the boundary if the fix
 681 point is not within the domain. However, by setting a proper
 682 maximum number of iterations, the algorithm can return the
 683 optimal boundary value. For quantile moments, the logic is
 684 similar, if the percentiles do not exceed the breakdown point.
 685 If this is the case, consistent estimation is impossible, and
 686 the algorithm will stop due to the maximum number of iter-
 687 ations. The fix point iteration is, in principle, similar to the
 688 iterative reweighing in M-estimator, but an advantage of this
 689 algorithm is that the optimization is solely related to the d
 690 value function and is independent of the sample size (except
 691 for the quantile moments, which require re-computation of the
 692 quantile function, but this operation has a time complexity of
 693 $O(1)$ for a sorted sample). Since $|rskew|$ can specify d_{rm} after
 694 optimization, this algorithm enables the robust estimations of
 695 all four moments to reach a near-consistent level for common
 696 unimodal distributions (Table ??, SI Dataset S1), just using
 697 the Weibull distribution as the consistent distribution.

698 **E. Variance.** As one of the fundamental theorems in statistics,
 699 the central limit theorem declares that the standard deviation
 700 of the limiting form of the sampling distribution of the sample
 701 mean is $\frac{\sigma}{\sqrt{n}}$. The principle, asymptotic normality, was later
 702 applied to the sampling distributions of robust location esti-
 703 mators (2, 36, 43–50) and it was found that the efficiencies
 704 of the robust location estimators can differ significantly from
 705 that of the arithmetic mean. Daniell (1920) stated (43) that
 706 comparing the efficiencies of various kinds of estimators is
 707 useless unless they all tend to coincide asymptotically. Bickel
 708 and Lehmann, also in the landmark series (49, 50), argued
 709 that meaningful comparisons can be made by studying the
 710 standardized variances, asymptotic variances, and sharp lower
 711 bounds of these estimators. Here, the scaled standard error
 712 (SSE) is proposed to estimate the variances of all estimators, in-
 713 cluding recombined/quantile moments, on a scale comparable
 714 to that of the sample mean.

715 **Definition E.1** (Scaled standard error). Let $\mathcal{M}_{s_i s_j} \in \mathbb{R}^{i \times j}$
 716 denote the sample-by-statistics matrix, i.e., the first column
 717 is the main statistics of interest, $\widehat{\theta}_m$, the second to the j th
 718 column are $j - 1$ statistics required to scale, $\widehat{\theta}_{r_1}, \widehat{\theta}_{r_2}, \dots$,
 719 $\widehat{\theta}_{r_{j-1}}$. Then, the scaling factor $\mathcal{S} = \left[1, \frac{\widehat{\theta}_{r_1}}{\widehat{\theta}_m}, \frac{\widehat{\theta}_{r_2}}{\widehat{\theta}_m}, \dots, \frac{\widehat{\theta}_{r_{j-1}}}{\widehat{\theta}_m}\right]^T$
 720 is a $j \times 1$ matrix, which $\widehat{\theta}$ is the mean of the column. The
 721 normalized matrix is $\mathcal{M}_{s_i s_j}^N = \mathcal{M}_{s_i s_j} \mathcal{S}$. The SSEs are the
 722 unbiased standard deviations of the corresponding columns.

Setting the U -statistic moments as the main statistics of
 interest, the SSEs of all robust estimators proposed here are
 often between those of the median moments and those of the
 unbiased sample moments (SI Dataset S1). This is because
 similar monotonic relations between robustness and variance
 are also very common, e.g., Bickel and Lehmann (50) proved
 that a lower bound for the efficiency of TM_ϵ to sample mean
 is $(1 - 2\epsilon)^2$ and this monotonic bound holds true for any dis-
 tribution. However, the direction of monotonicity differs for
 distributions with different kurtosis. Lehmann and Scheffé
 (1950, 1955) (51, 52) in their two early papers provided a
 way to construct a uniformly minimum-variance unbiased
 estimator (UMVUE). From that, the sample mean and unbi-
 ased sample second moment can be proven as the UMVUEs
 for the population mean and population second moment for
 the Gaussian distribution. While their performance for sub-
 Gaussian distributions is generally satisfied, they performs
 poorly when the distribution has a heavy tail and completely
 fails for distributions with infinite second moments. For sub-
 Gaussian distributions, the variance of a robust estimator is
 generally monotonic increasing as its robustness increases, but
 for heavy-tailed distributions, the relation is reversed. The gen-
 eral monotonic relation between the concentration bound of an
 estimator and the kurtosis of the underlying distribution was
 also implied by Devroye, Lerasle, Lugosi, and Oliveira (2016)
 (53). As a result, for distributions with different kurtosis, the
 variance-optimal choice can be very different.

Lai, Robbins, and Yu (1983) proposed an estimator that
 adaptively chooses the mean or median in a symmetric distri-
 bution and showed that the results were typically as good as
 the better of the sample mean and median regarding variance
 (54). It can be interpreted as an attempt to use the variance
 version of the mean-SWA-median inequality. While they used
 bootstrap standard error as the criterion, another approach
 can be dated back to Laplace (1812) (55) is using a linear
 combination of median and mean and the weight is deduced
 to achieve minimum variance; examples for symmetric distri-
 butions see Samuel-Cahn (1994), Chan and He (1994), and
 Damilano and Puig (2004)'s papers (56–58).

Scaled standard error allows for direct comparison of vari-
 ances for different location estimators in asymmetric distribu-
 tions. In this study, twelve possible combinations were created
 using two invariant means and related symmetric weighted av-
 erages ($BM_{\frac{1}{8}}$, $SQM_{\frac{1}{8}}$, $BM_{\nu=2, \epsilon=\frac{1}{8}}$, $WM_{\frac{1}{8}}$, $BWM_{\frac{1}{8}}$, and $TM_{\frac{1}{8}}$
 used here). Each combination has a SSE for a single-parameter
 distribution, which can be inferred through a Monte Carlo
 study. Then, the combination with the smallest SSE is chosen
 (if the percentiles of quantile moments exceed the breakdown
 point, this combination will be excluded). Similar to Subsec-
 tion D, let $I(|skewness|, kurtosis, k, dtype, n) = ikm_{WA}$ de-
 note these relations. Then, since $\lim_{rkurt \rightarrow \infty} \frac{I(rkurt, 4)}{I(rkurt, 2)^2 rkurt} <$
 1, the same fix point iteration algorithm can be used to choose
 the variance-optimum combination. The only difference is
 that unlike D , I is defined to be discontinuous but linear
 interpolation can also ensure continuity. This approach yields
 results that are often nearly optimal (SI Datasets S1).

Data Availability. Data for Table ?? are given in SI Dataset S1.
 All codes have been deposited in [GitHub](#).

ACKNOWLEDGMENTS. I gratefully acknowledge the construc-
 tive comments made by the editor which substantially improved

- 784 1. CF Gauss, *Theoria combinationis observationum erroribus minimis obnoxiae*. (Henricus
785 Dieterich), (1823).
- 786 2. S Newcomb, A generalized theory of the combination of observations so as to obtain the best
787 result. *Am. journal Math.* **8**, 343–366 (1886).
- 788 3. S Newcomb, Researches on the motion of the moon. part ii, the mean motion of the moon and
789 other astronomical elements derived from observations of eclipses and occultations extending
790 from the period of the babylonians until ad 1908. *United States. Naut. Alm. Off. Astron. paper*;
791 v. **9** **9**, 1 (1912).
- 792 4. PJ Huber, Robust estimation of a location parameter. *Ann. Math. Stat.* **35**, 73–101 (1964).
- 793 5. X He, WK Fung, Method of medians for lifetime data with weibull models. *Stat. medicine* **18**,
794 1993–2009 (1999).
- 795 6. M Menon, Estimation of the shape and scale parameters of the weibull distribution. *Techno-*
796 *metrics* **5**, 175–182 (1963).
- 797 7. SD Dubey, Some percentile estimators for weibull parameters. *Technometrics* **9**, 119–129
798 (1967).
- 799 8. KM Hassanein, Percentile estimators for the parameters of the weibull distribution. *Biometrika*
800 **58**, 673–676 (1971).
- 801 9. NB Marks, Estimation of weibull parameters from common percentiles. *J. applied Stat.* **32**,
802 17–24 (2005).
- 803 10. K Boudt, D Caliskan, C Croux, Robust explicit estimators of weibull parameters. *Metrika* **73**,
804 187–209 (2011).
- 805 11. SD Dubey, *Contributions to statistical theory of life testing and reliability*. (Michigan State
806 University of Agriculture and Applied Science. Department of statistics), (1960).
- 807 12. LJ Bain, CE Antle, Estimation of parameters in the weibull distribution. *Technometrics* **9**,
808 621–627 (1967).
- 809 13. RV Hogg, Adaptive robust procedures: A partial review and some suggestions for future
810 applications and theory. *J. Am. Stat. Assoc.* **69**, 909–923 (1974).
- 811 14. RJ Hyndman, Y Fan, Sample quantiles in statistical packages. *The Am. Stat.* **50**, 361–365
812 (1996).
- 813 15. C Bernard, R Kazzi, S Vanduffel, Range value-at-risk bounds for unimodal distributions under
814 partial information. *Insur. Math. Econ.* **94**, 9–24 (2020).
- 815 16. WR van Zwet, *Convex Transformations of Random Variables: Nebst Stellingen*. (1964).
- 816 17. AL Bowley, *Elements of statistics*. (King) No. 8, (1926).
- 817 18. RA Groeneveld, G Meeden, Measuring skewness and kurtosis. *J. Royal Stat. Soc. Ser. D*
818 *(The Stat.)* **33**, 391–399 (1984).
- 819 19. PJ Bickel, EL Lehmann, Descriptive statistics for nonparametric models iv. spread in *Selected*
820 *Works of EL Lehmann*. (Springer), pp. 519–526 (2012).
- 821 20. PJ Rousseeuw, C Croux, Alternatives to the median absolute deviation. *J. Am. Stat. associa-*
822 *tion* **88**, 1273–1283 (1993).
- 823 21. PJ Bickel, EL Lehmann, Descriptive statistics for nonparametric models. iii. dispersion in
824 *Selected works of EL Lehmann*. (Springer), pp. 499–518 (2012).
- 825 22. P Janssen, R Serfling, N Veraverbeke, Asymptotic normality of u-statistics based on trimmed
826 samples. *J. statistical planning inference* **16**, 63–74 (1987).
- 827 23. J Hodges, E Lehmann, Matching in paired comparisons. *The Annals Math. Stat.* **25**, 787–791
828 (1954).
- 829 24. S Dharmadhikari, K Jogdeo, Unimodal laws and related in *A Festschrift For Erich L. Lehmann*.
830 (CRC Press), p. 131 (1982).
- 831 25. AY Khintchine, On unimodal distributions. *Izv. Nauchno-Isled. Inst. Mat. Mech.* **2**, 1–7 (1938).
- 832 26. S Purkayastha, Simple proofs of two results on convolutions of unimodal distributions. *Stat. &*
833 *probability letters* **39**, 97–100 (1998).
- 834 27. RA Fisher, Moments and product moments of sampling distributions. *Proc. Lond. Math. Soc.*
835 **2**, 199–238 (1930).
- 836 28. PR Halmos, The theory of unbiased estimation. *The Annals Math. Stat.* **17**, 34–43 (1946).
- 837 29. W Hoeffding, A class of statistics with asymptotically normal distribution. *The Annals Math.*
838 *Stat.* **19**, 293–325 (1948).
- 839 30. PM Heffernan, Unbiased estimation of central moments by using u-statistics. *J. Royal Stat.*
840 *Soc. Ser. B (Statistical Methodol.)* **59**, 861–863 (1997).
- 841 31. D Fraser, Completeness of order statistics. *Can. J. Math.* **6**, 42–45 (1954).
- 842 32. AJ Lee, *U-statistics: Theory and Practice*. (Routledge), (2019).
- 843 33. J Hodges Jr, E Lehmann, Estimates of location based on rank tests. *The Annals Math. Stat.*
844 **34**, 598–611 (1963).
- 845 34. A Ehsanes Saleh, Hodges-lehmann estimate of the location parameter in censored samples.
846 *Annals Inst. Stat. Math.* **28**, 235–247 (1976).
- 847 35. RJ Serfling, Generalized l-, m-, and r-statistics. *The Annals Stat.* **12**, 76–86 (1984).
- 848 36. P Janssen, R Serfling, N Veraverbeke, Asymptotic normality for a general class of statistical
849 functions and applications to measures of spread. *The Annals Stat.* **12**, 1369–1379 (1984).
- 850 37. MG Akritas, Empirical processes associated with v-statistics and a class of estimators under
851 random censoring. *The Annals Stat.* **14**, 619–637 (1986).
- 852 38. I Gijbels, P Janssen, N Veraverbeke, Weak and strong representations for trimmed u-statistics.
853 *Probab. theory related fields* **77**, 179–194 (1988).
- 854 39. J Choudhury, R Serfling, Generalized order statistics, bahadur representations, and sequential
855 nonparametric fixed-width confidence intervals. *J. Stat. Plan. Inference* **19**, 269–282 (1988).
- 856 40. E Joly, G Lugosi, Robust estimation of u-statistics. *Stoch. Process. their Appl.* **126**, 3760–3773
857 (2016).
- 858 41. P Laforge, S Cléménçon, P Bertail, On medians of (randomized) pairwise means in *Interna-*
859 *tional Conference on Machine Learning*. (PMLR), pp. 1272–1281 (2019).
- 860 42. K Pearson, Contributions to the mathematical theory of evolution. *Philos. Transactions Royal*
861 *Soc. London. A* **185**, 71–110 (1894).
- 862 43. P Daniell, Observations weighted according to order. *Am. J. Math.* **42**, 222–236 (1920).
- 863 44. F Mosteller, On some useful "inefficient" statistics. *The Annals Math. Stat.* **17**, 377–408 (1946).
- 864 45. CR Rao, *Advanced statistical methods in biometric research*. (Wiley), (1952).
- 865 46. PJ Bickel, , et al., Some contributions to the theory of order statistics in *Proc. Fifth Berkeley*

- Sympos. Math. Statist. and Probability*. Vol. 1, pp. 575–591 (1967). 866
47. H Chernoff, JL Gastwirth, MV Johns, Asymptotic distribution of linear combinations of functions
867 of order statistics with applications to estimation. *The Annals Math. Stat.* **38**, 52–72 (1967). 868
48. L LeCam, On the assumptions used to prove asymptotic normality of maximum likelihood
869 estimates. *The Annals Math. Stat.* **41**, 802–828 (1970). 870
49. P Bickel, E Lehmann, Descriptive statistics for nonparametric models i. introduction in *Selected*
871 *Works of EL Lehmann*. (Springer), pp. 465–471 (2012). 872
50. PJ Bickel, EL Lehmann, Descriptive statistics for nonparametric models ii. location in *selected*
873 *works of EL Lehmann*. (Springer), pp. 473–497 (2012). 874
51. EL Lehmann, H Scheffé, Completeness, similar regions, and unbiased estimation-part i in
875 *Selected works of EL Lehmann*. (Springer), pp. 233–268 (2011). 876
52. EL Lehmann, H Scheffé, *Completeness, similar regions, and unbiased estimation—part II*.
877 (Springer), (2012). 878
53. L Devroye, M Lerasle, G Lugosi, RI Oliveira, Sub-gaussian mean estimators. *The Annals Stat.*
879 **44**, 2695–2725 (2016). 880
54. T Lai, H Robbins, K Yu, Adaptive choice of mean or median in estimating the center of a
881 symmetric distribution. *Proc. Natl. Acad. Sci.* **80**, 5803–5806 (1983). 882
55. PS Laplace, *Theorie analytique des probabilités*. (1812). 883
56. E Samuel-Cahn, Combining unbiased estimators. *The Am. Stat.* **48**, 34 (1994). 884
57. Y Chan, X He, A simple and competitive estimator of location. *Stat. & Probab. Lett.* **19**,
885 137–142 (1994). 886
58. G Damilano, P Puig, Efficiency of a linear combination of the median and the sample mean:
887 The double truncated normal distribution. *Scand. J. Stat.* **31**, 629–637 (2004). 888