Near-consistent robust estimations of moments for unimodal distributions

Tuban Lee

This manuscript was compiled on June 10, 2023

orderliness | invariant | unimodal | adaptive estimation | U-statistics

A. Congruent distribution. In the realm of nonparametric statistics, the relative differences, or orders, of robust estimators are of primary importance. A key implication of this principle is that when there is a shift in the parameters of the underlying distribution, all nonparametric estimates should asymptotically change in the same direction, if they are estimating the same attribute of the distribution. If, on the other hand, the mean suggests an increase in the location of the distribution while the median indicates a decrease, a contradiction arises. It is worth noting that such contradiction is not possible for any LL-statistics in a location-scale 11 distribution, as explained in the previous article on semipara-12 metric robust mean. However, it is possible to construct 13 counterexamples to the aforementioned implication in a shape-14 scale distribution. In the case of the Weibull distribution, 15 its quantile function is $Q_{Wei}(p) = \lambda(-\ln(1-p))^{1/\alpha}$, where 16 $0 \le p \le 1, \ \alpha > 0, \ \lambda > 0, \ \lambda$ is a scale parameter, α is a shape parameter, ln is the natural logarithm function. Then, $m = \lambda \sqrt[\alpha]{\ln(2)}, \ \mu = \lambda \Gamma \left(1 + \frac{1}{\alpha}\right), \text{ where } \Gamma \text{ is the gamma func-}$ 19 tion. When $\alpha = 1$, $m = \lambda \ln(2) \approx 0.693\lambda$, $\mu = \lambda$, when $\alpha = \frac{1}{2}$, $m = \lambda \ln^2(2) \approx 0.480\lambda$, $\mu = 2\lambda$, the mean increases as α 21 changes from 1 to $\frac{1}{2}$, but the median decreases. Previously, the fundamental role of quantile average and its relation to 23 nearly all common nonparametric robust location estimates were demonstrated by using the method of classifying distributions through the signs of derivatives. To avoid such scenarios, this method can also be used.

Data Availability. Data for Table ?? are given in SI Dataset S1.
All codes are attached.

ACKNOWLEDGMENTS. I gratefully acknowledge the constructive comments made by the editor which substantially improved the clarity and quality of this paper.