

# Robust estimations of moments for unimodal distributions

Tuban Lee

This manuscript was compiled on June 11, 2023

**A. Invariant Moments.** All popular robust location estimators, such as the symmetric trimmed mean, symmetric Winsorized mean, Hodges-Lehmann estimator, Huber  $M$ -estimator, and median of means, are symmetric. As shown in RSSM I, a  $\gamma$ -weighted Hodges-Lehmann mean ( $\text{WHLM}_{k,\epsilon,\gamma}$ ) can achieve consistency for the population mean in any  $\gamma$ -symmetric distribution with a finite mean. However, it falls considerably short of consistently handling other parametric distributions that are not  $\gamma$ -symmetric. Shifting from semiparametrics to parametrics, consider a robust estimator with a non-sample-dependent breakdown point (defined in Subsection ??) which is consistent simultaneously for both a semiparametric distribution and a parametric distribution that does not belong to that semiparametric distribution, it is named with the prefix ‘invariant’ followed by the name of the population parameter it is consistent with. Here, the recombined  $I$ -statistic is defined as

$$\text{RI}_{d,\mathbf{k}_1,\mathbf{k}_2,k_1,k_2,\epsilon_1,\epsilon_2,\gamma_1,\gamma_2,n,LU_1,LU_2} := \lim_{c \rightarrow \infty} \left( \frac{(LU_1 \mathbf{k}_1, k_1, \epsilon_1, \gamma_1, n + c)^{d+1}}{(LU_2 \mathbf{k}_2, k_2, \epsilon_2, \gamma_2, n + c)^d} - c \right),$$

where  $d$  is the key factor for bias correction,  $LU_{\mathbf{k},k,\epsilon,\gamma,n}$  is the  $LU$ -statistic,  $\mathbf{k}$  is the degree of the  $U$ -statistic,  $k$  is the degree of the  $LL$ -statistic,  $\epsilon$  is the upper asymptotic breakdown point of the  $LU$ -statistic. It is assumed in this series that in the subscript of an estimator, if  $\mathbf{k}$ ,  $k$  and  $\gamma$  are omitted,  $\mathbf{k} = 1$ ,  $k = 1$ ,  $\gamma = 1$  are assumed, if just one  $\gamma$  is indicated,  $\gamma_1 = \gamma_2$ , if  $n$  is omitted, only the asymptotic behavior is considered, in the absence of subscripts, no assumptions are made. The subsequent theorem shows the significance of a recombined  $I$ -statistic.

**Theorem A.1.** Define the recombined mean as  $rm_{d,k_1,k_2,\epsilon_1,\epsilon_2,\gamma_1,\gamma_2,n,WL_1,WL_2} := \text{RI}_{d,\mathbf{k}_1=1,\mathbf{k}_2=1,k_1,k_2,\epsilon_1,\epsilon_2,\gamma_1,\gamma_2,n,LU_1=WL_1,LU_2=WL_2}$ . Assuming finite means,  $rm_{d,k_1,k_2,\epsilon_1,\epsilon_2,\gamma_1,\gamma_2,n,WL_1,WL_2} = \frac{\mu - WL_1 k_1, \epsilon_1, \gamma_1}{WL_1 k_1, \epsilon_1, \gamma_1 - WL_2 k_2, \epsilon_2, \gamma_2}$ ,  $k_1, k_2, \epsilon_1, \epsilon_2, \gamma_1, \gamma_2, WL_1, WL_2$  is a consistent mean estimator for a location-scale distribution, where  $\mu$ ,  $WL_1 k_1, \epsilon_1, \gamma_1$ , and  $WL_2 k_2, \epsilon_2, \gamma_2$  are different location parameters from that location-scale distribution. If  $\gamma_1 = \gamma_2$ ,  $WL = \text{WHLM}$ ,  $rm$  is also consistent for any  $\gamma$ -symmetric distributions.

*Proof.* Finding  $d$  that make  $rm_{d,k_1,k_2,\epsilon_1,\epsilon_2,\gamma_1,\gamma_2,WL_1,WL_2}$  a consistent mean estimator is equivalent to finding the solution of  $rm_{d,k_1,k_2,\epsilon_1,\epsilon_2,\gamma_1,\gamma_2,WL_1,WL_2} = \mu$ . First consider the location-scale distribution. Since  $rm_{d,k_1,k_2,\epsilon_1,\epsilon_2,\gamma_1,\gamma_2,WL_1,WL_2} = \frac{\mu - WL_1 k_1, \epsilon_1, \gamma_1}{WL_1 k_1, \epsilon_1, \gamma_1 - WL_2 k_2, \epsilon_2, \gamma_2}$ ,  $k_1, k_2, \epsilon_1, \epsilon_2, \gamma_1, \gamma_2, WL_1, WL_2$  is a consistent mean estimator for a location-scale distribution, where  $\mu$ ,  $WL_1 k_1, \epsilon_1, \gamma_1$ , and  $WL_2 k_2, \epsilon_2, \gamma_2$  are different location parameters from that location-scale distribution. If  $\gamma_1 = \gamma_2$ ,  $WL = \text{WHLM}$ ,  $rm$  is also consistent for any  $\gamma$ -symmetric distributions.

$dWL_2 k_2, \epsilon_2, \gamma_2 = \mu$ . So,  $d = \frac{\mu - WL_1 k_1, \epsilon_1, \gamma_1}{WL_1 k_1, \epsilon_1, \gamma_1 - WL_2 k_2, \epsilon_2, \gamma_2}$ . In RSSM I, it was established that any  $WL(k, \epsilon, \gamma)$  can be expressed as  $\lambda WL_0(k, \epsilon, \gamma) + \mu$  for a location-scale distribution parameterized by a location parameter  $\mu$  and a scale parameter  $\lambda$ , where  $WL_0(k, \epsilon, \gamma)$  is a function of  $Q_0(p)$ , the quantile function of a standard distribution without any shifts or scaling, according to the definition of the weighted  $L$ -statistic. The simultaneous cancellation of  $\mu$  and  $\lambda$  in  $\frac{(\lambda\mu_0 + \mu) - (\lambda WL_0(k_1, \epsilon_1, \gamma_1) + \mu)}{(\lambda WL_0(k_1, \epsilon_1, \gamma_1) + \mu) - (\lambda WL_0(k_2, \epsilon_2, \gamma_2) + \mu)}$  assures that the  $d$  in  $rm$  is always a constant for a location-scale distribution. The proof of the second assertion follows directly from the coincidence property. According to Theorem 18 in RSSM I, for any  $\gamma$ -symmetric distribution with a finite mean,  $\text{WHLM}_{1k_1, \epsilon_1, \gamma} = \text{WHLM}_{2k_2, \epsilon_2, \gamma} = \mu$ . Then  $rm_{d,k_1,k_2,\epsilon_1,\epsilon_2,\gamma,WL_1,WL_2} = \lim_{c \rightarrow \infty} \left( \frac{(\mu + c)^{d+1}}{(\mu + c)^d} - c \right) = \mu$ . This completes the demonstration.  $\square$

For example, the Pareto distribution has a quantile function  $Q_{Par}(p) = x_m(1-p)^{-\frac{1}{\alpha}}$ , where  $x_m$  is the minimum possible value that a random variable following the Pareto distribution can take, serving a scale parameter,  $\alpha$  is a shape parameter. The mean of the Pareto distribution is given by  $\frac{\alpha x_m}{\alpha - 1}$ . As  $WL(k, \epsilon, \gamma)$  can be expressed as a function of  $Q(p)$ , one can set the two  $WL_{k,\epsilon,\gamma}$ s in the  $d$  value as two arbitrary quantiles  $Q_{Par}(p_1)$  and  $Q_{Par}(p_2)$ . For the Pareto distribution,  $d_{Per} = \frac{\mu_{Per} - Q_{Par}(p_1)}{Q_{Par}(p_1) - Q_{Par}(p_2)} = \frac{\frac{\alpha x_m}{\alpha - 1} - x_m(1-p_1)^{-\frac{1}{\alpha}}}{x_m(1-p_1)^{-\frac{1}{\alpha}} - x_m(1-p_2)^{-\frac{1}{\alpha}}} \cdot x_m$  can be canceled out. Intriguingly, the quantile function of exponential distribution is  $Q_{exp}(p) = \ln\left(\frac{1}{1-p}\right)\lambda$ ,  $\lambda \geq 0$ .  $\mu_{exp} = \lambda$ . Then,  $d_{exp} = \frac{\mu_{exp} - Q_{exp}(p_1)}{Q_{exp}(p_1) - Q_{exp}(p_2)} = \frac{\lambda - \ln\left(\frac{1}{1-p_1}\right)\lambda}{\ln\left(\frac{1}{1-p_1}\right)\lambda - \ln\left(\frac{1}{1-p_2}\right)\lambda} = -\frac{\ln(1-p_1)+1}{\ln(1-p_1)-\ln(1-p_2)}$ . Since  $\lim_{\alpha \rightarrow \infty} \frac{\frac{\alpha}{\alpha-1} - (1-p_1)^{-1/\alpha}}{(1-p_1)^{-1/\alpha} - (1-p_2)^{-1/\alpha}} = -\frac{\ln(1-p_1)+1}{\ln(1-p_1)-\ln(1-p_2)}$ , the  $d$  value for the Pareto distribution approaches that of the exponential distribution, as  $\alpha \rightarrow \infty$ , regardless of the type of weighted  $L$ -statistic used. That means, for the Weibull, gamma,

## Significance Statement

Bias, variance, and contamination are the three main errors in statistics. Consistent robust estimation is unattainable without parametric assumptions. In this article, invariant moments are proposed as a means of achieving near-consistent and robust estimations of moments, even in scenarios where moderate violations of distributional assumptions occur, while the variances are sometimes smaller than those of the sample moments.

T.L. designed research, performed research, analyzed data, and wrote the paper.

The author declares no competing interest.

<sup>1</sup>To whom correspondence should be addressed. E-mail: tl@biomathematics.org

58 Pareto, lognormal and generalized Gaussian distribution,  
59  $rm_{d=\frac{\mu - \text{WHLM}_{1k_1, \epsilon_1, \gamma}}{\text{WHLM}_{1k_1, \epsilon_1, \gamma} - \text{WHLM}_{2k_2, \epsilon_2, \gamma}}, k_1, k_2, \epsilon_1, \epsilon_2, \gamma, \text{WHLM}_1, \text{WHLM}_2}$   
60 is consistent for at least one particular case, where  
61  $\mu$ ,  $\text{WHLM}_{1k_1, \epsilon_1, \gamma}$ , and  $\text{WHLM}_{2k_2, \epsilon_2, \gamma}$  are differ-  
62 ent location parameters from an exponential dis-  
63 tribution. Let  $\text{WHLM}_{1k_1, \epsilon_1, \gamma} = \text{BM}_{\nu=3, \epsilon=\frac{1}{24}}$ ,  
64  $\text{WHLM}_{2k_2, \epsilon_2, \gamma} = m$ , then  $\mu = \lambda$ ,  $m = Q\left(\frac{1}{2}\right) = \ln 2\lambda$ ,  
65  $\text{BM}_{\nu=3, \epsilon=\frac{1}{24}} = \lambda \left(1 + \ln \left(\frac{26068394603446272 \sqrt[6]{\frac{\pi}{247}} \sqrt[3]{11}}{391^{5/6} 101898752449325 \sqrt{5}}\right)\right)$ ,  
66 the detailed formula is given in the SI Text. So,  $d =$   
67  $\frac{\mu - \text{BM}_{\nu=3, \epsilon=\frac{1}{24}}}{\text{BM}_{\nu=3, \epsilon=\frac{1}{24}} - m} = \frac{\lambda - \lambda \left(1 + \ln \left(\frac{26068394603446272 \sqrt[6]{\frac{\pi}{247}} \sqrt[3]{11}}{391^{5/6} 101898752449325 \sqrt{5}}\right)\right)}{\lambda \left(1 + \ln \left(\frac{26068394603446272 \sqrt[6]{\frac{\pi}{247}} \sqrt[3]{11}}{391^{5/6} 101898752449325 \sqrt{5}}\right)\right) - \ln 2\lambda} =$   
68  $-\frac{\ln \left(\frac{26068394603446272 \sqrt[6]{\frac{\pi}{247}} \sqrt[3]{11}}{391^{5/6} 101898752449325 \sqrt{5}}\right)}{1 - \ln(2) + \ln \left(\frac{26068394603446272 \sqrt[6]{\frac{\pi}{247}} \sqrt[3]{11}}{391^{5/6} 101898752449325 \sqrt{5}}\right)} \approx 0.103$ . The biases of  
69  $rm_{d \approx 0.103, \nu=3, \epsilon_1=\frac{1}{24}, \epsilon_2=\frac{1}{2}, \text{BM}, m}$  for distributions with skewness  
70 between those of the exponential and symmetric distributions  
71 are tiny (SI Dataset S1).  $rm_{d \approx 0.103, \nu=3, \epsilon_1=\frac{1}{24}, \epsilon_2=\frac{1}{2}, \text{BM}, m}$   
72 exhibits excellent performance for all these common unimodal  
73 distributions (SI Dataset S1).  
74 The recombined mean is an recombined  $I$ -statistic.