

Semiparametric robust mean estimations based on the orderliness of quantile averages

Tuban Lee

This manuscript was compiled on May 28, 2023

As one of the most fundamental problems in statistics, the robust location estimation has many prominent solutions, such as the symmetric trimmed mean, symmetric Winsorized mean, Hodges–Lehmann estimator, Huber M -estimator, and median of means. Recent studies suggest that their maximum biases concerning the mean can be quite different in asymmetric distributions, but the underlying mechanisms and average performance remain largely unclear. This study establishes several forms of orderliness among quantile averages, similar to the mean-median-mode inequality, within a wide range of semiparametric distributions, particularly highlighting the unique role of γ -symmetric distributions. From this, a sequence of advanced robust mean estimators emerges, which also explains why the Winsorized mean and median of means typically have smaller biases compared to the trimmed mean. Building on the U -orderliness, the superiority of the median Hodges–Lehmann mean is discussed.

semiparametric | mean-median-mode inequality | asymptotic | unimodal | Hodges–Lehmann estimator

In 1823, Gauss (1) proved that for any unimodal distribution, $|m - \mu| \leq \sqrt{\frac{3}{4}}\omega$ and $\sigma \leq \omega \leq 2\sigma$, where μ is the population mean, m is the population median, ω is the root mean square deviation from the mode, and σ is the population standard deviation. This pioneering work revealed that despite potential bias in robust mean estimates, the deviation remains bounded in units of a scale parameter under certain assumptions. Bernard, Kazzi, and Vanduffel (2020) (2) further derived asymptotic bias bounds of any quantile for unimodal distributions with finite second moments, by reducing this optimization problem to a parametric one, which can be solved analytically. They showed that m has the smallest maximum distance to μ among all symmetric quantile averages (SQA _{ϵ}). Daniell, in 1920, (3) analyzed a class of estimators, linear combinations of order statistics, and identified that ϵ -symmetric trimmed mean (STM _{ϵ}) belongs to this class. Another popular choice, the ϵ -symmetric Winsorized mean (SWM _{ϵ}), named after Winsor and introduced by Tukey (4) and Dixon (5) in 1960, is also an L -estimator. Bieniek (2016) derived exact bias upper bounds of the Winsorized mean based on Danielak and Rychlik's work (2003) on the trimmed mean for any distribution with a finite second moment and confirmed that the former is smaller than the latter (6, 7). In 1963, Hodges and Lehmann (8) proposed a class of nonparametric location estimators based on rank tests and, from the Wilcoxon signed-rank statistic (9), deduced the median of pairwise means as a robust location estimator for a symmetric population. Both L -statistics and R -statistics achieve robustness essentially by removing a certain proportion of extreme values. In 1964, Huber (10) generalized maximum likelihood estimation to the minimization of the sum of a specific loss function, which measures the residuals between the data points and the model's parameters. Some L -estimators are also M -estimators, e.g., the sample mean is an M -estimator with a squared error loss

function, the sample median is an M -estimator with an absolute error loss function (10). The Huber M -estimator is obtained by applying the Huber loss function that combines elements of both squared error and absolute error to achieve robustness against gross errors and high efficiency for contaminated Gaussian distributions (10). Sun, Zhou, and Fan (2020) examined the concentration bounds of Huber M -estimator (11). Mathieu (2022) (12) further derived the concentration bounds of M -estimators and demonstrated that, by selecting the tuning parameter which depends on the variance, Huber M -estimator can also be a sub-Gaussian estimator. The concept of the median of means (MoM _{$k, b = \frac{n}{k}, n$}), which was first introduced by Nemirovsky and Yudin (1983) in their work on stochastic optimization (13). Given its good performance even for distributions with infinite second moments, MoM has received increasing attention over the past decade (14–17). Devroye, Lerasle, Lugosi, and Oliveira (2016) showed that MoM _{$k, b = \frac{n}{k}, n$} nears the optimum of sub-Gaussian mean estimation with regards to concentration bounds when the distribution has a heavy tail (15). Laforge, Clemencon, and Bertail (2019) proposed the median of randomized means (MoRM _{k, b, n}) (16), wherein, rather than partitioning, an arbitrary number, b , of blocks are built independently from the sample, and showed that MoRM _{k, b, n} has a better non-asymptotic sub-Gaussian property compared to MoM _{$k, b = \frac{n}{k}, n$} . In fact, asymptotically, the Hodges–Lehmann (H-L) estimator is equivalent to MoM _{$k=2, b = \frac{n}{k}$} and MoRM _{$k=2, b$} , and they can be seen as the pairwise mean distribution is approximated by the sampling without replacement and bootstrap, respectively. When $k \ll n$, the difference between sampling with replacement and without replacement is negligible. For the asymptotic validity, readers are referred to the foundational works of Efron (1979) (18), Bickel and Freedman (1981, 1984) (19, 20), and Helmers, Janssen, and Veraverbeke (1990) (21).

Significance Statement

In 1964, van Zwet introduced the convex transformation order for comparing the skewness of two distributions. This paradigm shift played a fundamental role in defining robust measures of distributions, from spread to kurtosis. Here, instead of examining the stochastic ordering between two distributions, the orderliness of quantile averages within a distribution is investigated. By classifying distributions through the signs of derivatives, a series of sophisticated robust mean estimators is deduced. Nearly all common nonparametric robust location estimators are found to be special cases thereof.

T.L. designed research, performed research, analyzed data, and wrote the paper.

The author declares no competing interest.

¹To whom correspondence should be addressed. E-mail: tl@biomathematics.org

Here, the ϵ, b -stratified mean is defined as

$$\text{SM}_{\epsilon, b, n} := \frac{b}{n} \left(\sum_{j=1}^{\lfloor \frac{b-1}{2b\epsilon} \rfloor} \sum_{i_j = \frac{(2bj-b-1)n\epsilon}{b-1}}^{\frac{(2bj-b+1)n\epsilon}{b-1}} X_{i_j} \right),$$

where $X_1 \leq \dots \leq X_n$ denote the order statistics of a sample of n independent and identically distributed random variables X_1, \dots, X_n . $b \in \mathbb{N}$, $b \geq 3$. The definition was further refined to guarantee the continuity of the breakdown point by incorporating an additional block in the center when $\lfloor \frac{b-1}{2b\epsilon} \rfloor \bmod 2 = 0$, or by adjusting the central block when $\lfloor \frac{b-1}{2b\epsilon} \rfloor \bmod 2 = 1$ (SI Text). If the subscript n is omitted, only the asymptotic behavior is considered. If b is omitted, $b = 3$ is assumed. $\text{SM}_{\epsilon, b=3}$ is equivalent to STM_{ϵ} , when $\epsilon > \frac{1}{6}$. When $\frac{b-1}{2\epsilon} \in \mathbb{N}$, $b \bmod 2 = 1$, the basic idea of the stratified mean is to distribute the data into $\frac{b-1}{2\epsilon}$ equal-sized non-overlapping blocks according to their order, then further sequentially group these blocks into b equal-sized strata and compute the mean of the middle stratum, which is the median of means of each stratum. In situations where $i \bmod 1 \neq 0$, a potential solution is to generate multiple smaller samples that satisfy the equality by sampling without replacement, and subsequently calculate the mean of all estimations. The details of determining the sample size and sampling times are provided in the SI Text. Although the principle resembles that of the median of means, $\text{SM}_{\epsilon, b, n}$ is different from $\text{MoM}_{k=\frac{n}{b}, b, n}$ as it does not include the random shift. Additionally, the stratified mean differs from the mean of the sample obtained through stratified sampling methods, introduced by Neyman (1934) (22) or ranked set sampling (23), introduced by McIntyre in 1952, as these sampling methods aim to obtain more representative samples or improve the efficiency of sample estimates, but the sample means based on them are not robust. When $b \bmod 2 = 1$, the stratified mean can be regarded as replacing the other equal-sized strata with the middle stratum, which, in principle, is analogous to the Winsorized mean that replaces extreme values with less extreme percentiles. Furthermore, while the bounds confirm that the Winsorized mean and median of means outperform the trimmed mean (6, 7, 15) in worst-case performance, the complexity of bound analysis makes it difficult to achieve a complete and intuitive understanding of these results. Also, a clear explanation for the average performance of them remains elusive. The aim of this paper is to define a series of semi-parametric models using the signs of derivatives, reveal their elegant interrelations and connections to parametric models, and show that by exploiting these models, a set of sophisticated mean estimators can be deduced, which exhibit strong robustness to departures from assumptions.

Quantile average and weighted average

The symmetric trimmed mean, symmetric Winsorized mean, and stratified mean are all L -estimators. More specifically, they are symmetric weighted averages, which are defined as

$$\text{SWA}_{\epsilon, n} := \frac{\sum_{i=1}^{\lceil \frac{n}{2} \rceil} \frac{X_i + X_{n-i+1}}{2} w_i}{\sum_{i=1}^{\lceil \frac{n}{2} \rceil} w_i},$$

where w_i s are the weights applied to the symmetric quantile averages according to the definition of the corresponding L -estimators. For example, for the ϵ -symmetric trimmed mean,

$w_i = \begin{cases} 0, & i < n\epsilon \\ 1, & i \geq n\epsilon \end{cases}$, provided that $n\epsilon \in \mathbb{N}$. The mean and median are indeed two special cases of the symmetric trimmed mean.

To extend the symmetric quantile average to the asymmetric case, two definitions for the ϵ, γ -quantile average ($\text{QA}(\epsilon, \gamma, n)$) are proposed. The first definition is:

$$\frac{1}{2}(\hat{Q}_n(\gamma\epsilon) + \hat{Q}_n(1-\epsilon)), \quad [1]$$

and the second definition is:

$$\frac{1}{2}(\hat{Q}_n(\epsilon) + \hat{Q}_n(1-\gamma\epsilon)), \quad [2]$$

where $\hat{Q}_n(p)$ is the empirical quantile function; γ is used to adjust the degree of asymmetry, $\gamma \geq 0$; and $0 \leq \epsilon \leq \frac{1}{1+\gamma}$. For trimming from both sides, [1] and [2] are essentially equivalent. The first definition along with $\gamma \geq 0$ and $0 \leq \epsilon \leq \frac{1}{1+\gamma}$ are assumed in the remainder of this article unless otherwise specified, since many common asymmetric distributions are right-skewed, and [1] allows trimming only from the right side by setting $\gamma = 0$.

Analogously, the weighted average can be defined as

$$\text{WA}_{\epsilon, \gamma, n} := \frac{\int_{\epsilon_0=0}^{\frac{1}{1+\gamma}} \text{QA}(\epsilon_0, \gamma, n) w_{\epsilon_0}}{\int_{\epsilon_0=0}^{\frac{1}{1+\gamma}} w_{\epsilon_0}}.$$

For any weighted average, if γ is omitted, $\gamma = 1$ is assumed. The ϵ, γ -trimmed mean ($\text{TM}_{\epsilon, \gamma, n}$) is a weighted average with a left trim size of $\gamma\epsilon n$ and a right trim size of ϵn , where $w_{\epsilon_0} = \begin{cases} 0, & \epsilon_0 < \epsilon \\ 1, & \epsilon_0 \geq \epsilon \end{cases}$. Using this definition, the TM computation remains the same, regardless of whether $\gamma\epsilon n \notin \mathbb{N}$ or $\epsilon n \notin \mathbb{N}$, since this definition is based on the empirical quantile function. However, in this article, considering the computational cost in practice, the non-asymptotic definitions of various types of weighted averages are essentially based on order statistics, in most cases. Unless stated otherwise, the solution to their decimal issue is the same as that in SM.

Classifying distributions by the signs of derivatives

Let \mathcal{P}_R denote the set of all continuous distribution over \mathbb{R} . Without loss of generality, the discussion of all the classes outlined below is restricted to the intersection with this non-parametric class of distributions. Besides fully and smoothly parameterizing by a Euclidean parameter or just assuming regularity conditions, there are many ways to classify distributions. In 1956, Stein initiated the problem of estimating parameters in the presence of an infinite dimensional nuisance shape parameter (24). A notable example discussed in his groundbreaking work was the estimation of the center of symmetry for an unknown symmetric distribution. In 1993, Bickel, Klaassen, Ritov, and Wellner published an influential semi-parametrics textbook (25) which systematically categorized many common models into three classes: parametric, non-parametric, and semiparametric. Yet, there is another old and commonly encountered class of distributions that receives little attention in semiparametric literature: the unimodal distribution. It is a very unique semiparametric model because its definition is based on the signs of derivatives, i.e.,

($f'(x) > 0$ for $x \leq M$) \wedge ($f'(x) < 0$ for $x \geq M$), where $f(x)$ is the probability density function (pdf) of a random variable X , M is the mode. Let \mathcal{P}_U denote the set of all unimodal distributions. There was a widespread misbelief that the median of an arbitrary unimodal distribution always lies between its mean and mode until Runnenburg (1978) and van Zwet (1979) (26, 27) endeavored to determine sufficient conditions for the inequality to hold, thereby implying the possibility of its violation. The class of distributions that satisfy the mean-median-mode inequality constitutes a subclass of \mathcal{P}_U . By analogy, a right-skewed distribution is called γ -ordered, if and only if

$$\forall 0 \leq \epsilon_1 \leq \epsilon_2 \leq \frac{1}{1+\gamma}, QA_{\epsilon_1, \gamma} \geq QA_{\epsilon_2, \gamma}.$$

The necessary and sufficient condition below hints at the relation between the mean-median-mode inequality and the γ -orderliness.

Theorem .1. *A distribution is γ -ordered if and only if its pdf satisfies the inequality $f(Q(\gamma\epsilon)) \geq f(Q(1-\epsilon))$ for all $0 \leq \epsilon \leq \frac{1}{1+\gamma}$ or $f(Q(\gamma\epsilon)) \leq f(Q(1-\epsilon))$ for all $0 \leq \epsilon \leq \frac{1}{1+\gamma}$.*

Proof. Without loss of generality, consider the case of right-skewed distribution. From the definition of γ -orderliness, it is deduced that $\frac{Q(\gamma\epsilon-\delta)+Q(1-\epsilon+\delta)}{2} \geq \frac{Q(\gamma\epsilon)+Q(1-\epsilon)}{2} \Leftrightarrow Q(\gamma\epsilon-\delta) - Q(\gamma\epsilon) \geq Q(1-\epsilon) - Q(1-\epsilon+\delta) \Leftrightarrow Q'(1-\epsilon) \geq Q'(\gamma\epsilon)$, where δ is an infinitesimal positive quantity. Observing that the quantile function is the inverse function of the cumulative distribution function (cdf), $Q'(1-\epsilon) \geq Q'(\gamma\epsilon) \Leftrightarrow F'(Q(\gamma\epsilon)) \geq F'(Q(1-\epsilon))$, thereby completing the proof, given that the derivative of cdf is pdf. \square

According to Theorem .1, if a probability distribution is right-skewed and monotonic, it will always be γ -ordered. For a right-skewed unimodal distribution, if $Q(\gamma\epsilon) > M$, the inequality $f(Q(\gamma\epsilon)) \geq f(Q(1-\epsilon))$ holds. The principle is extendable to unimodal-like distributions. Suppose there is a right-skewed multimodal distribution following the mean- γ -median-first mode inequality with several smaller modes on the right side, with the first mode, M_1 , having the greatest probability density, and the γ -median, $Q(\frac{\gamma}{1+\gamma})$, falling within the first dominant mode (i.e., if $x > Q(\frac{\gamma}{1+\gamma})$, $f(Q(\frac{\gamma}{1+\gamma})) \geq f(x)$), then if $Q(\gamma\epsilon) > M_1$, the inequality $f(Q(\gamma\epsilon)) \geq f(Q(1-\epsilon))$ also holds. In other words, while a distribution following the mean- γ -median-mode inequality may not be strictly γ -ordered, the inequality that defines γ -orderliness remains valid for most quantile averages. The mean- γ -median-mode inequality can also indicate possible bounds for γ in practice, e.g., for any distributions, when $\gamma \rightarrow \infty$, the γ -median will be greater than the mean and the mode, when $\gamma \rightarrow 0$, the γ -median will be smaller than the mean and the mode.

Consider the sign of the derivative of the quantile average with respect to the breakdown point; the above definition of γ -orderliness can also be expressed as

$$\forall 0 \leq \epsilon \leq \frac{1}{1+\gamma}, \frac{\partial QA_{\epsilon, \gamma}}{\partial \epsilon} \leq 0.$$

The left-skewed case can be obtained by reversing the inequality $\frac{\partial QA_{\epsilon, \gamma}}{\partial \epsilon} \leq 0$ to $\frac{\partial QA_{\epsilon, \gamma}}{\partial \epsilon} \geq 0$ and employing the second definition of QA, as given in [2]. For simplicity, it will be

omitted in the following discussion. If $\gamma = 1$, the γ -ordered distribution is referred to as ordered.

Furthermore, many common right-skewed distributions are partial bounded, indicating a convex behavior of the QA function when $\epsilon \rightarrow 0$. By further assuming convexity, the second γ -orderliness can be defined for a right-skewed distribution as follows,

$$\forall 0 \leq \epsilon \leq \frac{1}{1+\gamma}, \frac{\partial^2 QA_{\epsilon, \gamma}}{\partial \epsilon^2} \geq 0 \wedge \frac{\partial QA_{\epsilon, \gamma}}{\partial \epsilon} \leq 0.$$

Analogously, the ν th γ -orderliness of a right-skewed distribution can be defined as $(-1)^\nu \frac{\partial^\nu QA_{\epsilon, \gamma}}{\partial \epsilon^\nu} \geq 0 \wedge \dots \wedge -\frac{\partial QA_{\epsilon, \gamma}}{\partial \epsilon} \geq 0$. If $\gamma = 1$, the ν th γ -orderliness is referred as ν th orderliness. Let \mathcal{P}_O denote the set of all distributions that are ordered and \mathcal{P}_{O_ν} and $\mathcal{P}_{\gamma O_\nu}$ represent the sets of all distributions that are ν th ordered and ν th γ -ordered, respectively. When the shape parameter of the Weibull distribution, α , is smaller than 3.258, it can be shown that the Weibull distribution belong to $\mathcal{P}_U \cap \mathcal{P}_O \cap \mathcal{P}_{O_2} \cap \mathcal{P}_{O_3}$ (SI Text). At $\alpha \approx 3.602$, the Weibull distribution is symmetric, and as $\alpha \rightarrow \infty$, the skewness of the Weibull distribution reaches 1. Therefore, the parameters that prevent it from being included in the set correspond to cases when it is near-symmetric, as shown in the SI Text. Nevertheless, computing the derivatives of the QA function is often intricate and, at times, challenging. The following theorems establish the relationship between \mathcal{P}_O , \mathcal{P}_{O_ν} , and $\mathcal{P}_{\gamma O_\nu}$, and a wide range of other semi-parametric distributions. They can be used to quickly identify some parametric distributions in \mathcal{P}_O , \mathcal{P}_{O_ν} , and $\mathcal{P}_{\gamma O_\nu}$.

Theorem .2. *For any random variable X whose probability distribution function belongs to a location-scale family, the distribution is ν th γ -ordered if and only if the family of probability distributions is ν th γ -ordered.*

Proof. Let Q_0 denote the quantile function of the standard distribution without any shifts or scaling. After a location-scale transformation, the quantile function is $Q(p) = \lambda Q_0(p) + \mu$, where λ is the scale parameter and μ is the location parameter. According to the definition of the ν th γ -orderliness, the signs of derivatives of the QA function are invariant after this transformation. As the location-scale transformation is reversible, the proof is complete. \square

Theorem .2 demonstrates that in the analytical proof of the ν th γ -orderliness of a parametric distribution, both the location and scale parameters can be regarded as constants. It is also instrumental in proving other theorems.

Theorem .3. *Define a γ -symmetric distribution as one for which the quantile function satisfies $Q(\gamma\epsilon) = 2Q(\frac{\gamma}{1+\gamma}) - Q(1-\epsilon)$, for all $0 \leq \epsilon \leq \frac{1}{1+\gamma}$. Any γ -symmetric distribution is ν th γ -ordered.*

Proof. The equality implies that $\frac{\partial Q(\gamma\epsilon)}{\partial \epsilon} = \gamma Q'(\gamma\epsilon) = \frac{\partial (-Q(1-\epsilon))}{\partial \epsilon} = Q'(1-\epsilon)$. From the definition of QA, the QA function of the γ -symmetric distribution is a horizontal line, since $\frac{\partial QA_{\epsilon, \gamma}}{\partial \epsilon} = \gamma Q'(\gamma\epsilon) - Q'(1-\epsilon) = 0$. So, the ν th order derivative of QA is always zero. \square

Theorem .4. *A symmetric distribution is a special case of a γ -symmetric distribution when $\gamma = 1$.*

Proof. Without loss of generality, assuming continuity. A symmetric distribution is a probability distribution such that for all x , $f(x) = f(2m - x)$. Its cdf satisfies $F(x) = 1 - F(2m - x)$. Let $x = Q(p)$, then, $F(Q(p)) = p = 1 - F(2m - Q(p))$ and $F(Q(1 - p)) = 1 - p \Leftrightarrow p = 1 - F(Q(1 - p))$. Therefore, $F(2m - Q(p)) = F(Q(1 - p))$. Since the cdf is monotonic, $2m - Q(p) = Q(1 - p)$. \square

As a consequence of Theorem .3, and due to the property that the generalized Gaussian distribution is symmetric around the median, it is found to be ν th ordered.

Theorem .5. Any right-skewed distribution whose quantile function Q satisfies $Q^{(\nu)}(p) \geq 0 \wedge \dots Q^{(i)}(p) \geq 0 \dots \wedge Q^{(2)}(p) \geq 0$, $i \bmod 2 = 0$, is ν th γ -ordered, provided that $0 \leq \gamma \leq 1$.

Proof. Since $(-1)^i \frac{\partial^i Q_{\epsilon, \gamma}}{\partial \epsilon^i} = \frac{1}{2}((- \gamma)^i Q^i(\gamma \epsilon) + Q^i(1 - \epsilon))$ and $1 \leq i \leq \nu$, when $i \bmod 2 = 0$, $(-1)^i \frac{\partial^i Q_{\epsilon, \gamma}}{\partial \epsilon^i} \geq 0$ for all $\gamma \geq 0$. When $i \bmod 2 = 1$, if further assuming $0 \leq \gamma \leq 1$, $(-1)^i \frac{\partial^i Q_{\epsilon, \gamma}}{\partial \epsilon^i} \geq 0$, since $Q^{(i+1)}(p) \geq 0$. \square

It is now straightforward to show that the Pareto distribution follows the ν th γ -orderliness, provided that $0 \leq \gamma \leq 1$, since the quantile function of the Pareto distribution is $Q(p) = x_m(1 - p)^{-\frac{1}{\alpha}}$, where $x_m > 0$, $\alpha > 0$, and so $Q^{(\nu)}(p) \geq 0$ for all $\nu \in \mathbb{N}$ according to the chain rule.

Theorem .6. A right-skewed distribution with a monotonic decreasing pdf is second γ -ordered.

Proof. A monotonic decreasing pdf implies $f'(x) = F^{(2)}(x) \leq 0$. Since $Q'(p) \geq 0$, let $x = Q(F(x))$, then by differentiating both sides of the equation twice, one can obtain $0 = Q^{(2)}(F(x))(F'(x))^2 + Q'(F(x))F^{(2)}(x) \Leftrightarrow Q^{(2)}(F(x)) = -\frac{Q'(F(x))F^{(2)}(x)}{(F'(x))^2} \geq 0$. The desired result is derived from Theorem .1 and .5. \square

Theorem .6 provides valuable insights into the relation between modality and orderliness. The conventional definition states that a distribution with a monotonic pdf is still considered unimodal. However, within its supported interval, the mode number is zero. The number of modes and their magnitudes within a distribution are closely related to the possibility of orderliness being valid, although counterexamples can always be constructed for non-monotonic distributions. It can be easily established that the gamma distribution is second γ -ordered, when $\alpha \leq 1$ as the pdf of the gamma distribution is $f(x) = \frac{\lambda^{-\alpha} x^{\alpha-1} e^{-\frac{x}{\lambda}}}{\Gamma(\alpha)}$, where $x \geq 0$, $\lambda > 0$, $\alpha > 0$, Γ is the gamma function, it is a product of two monotonic decreasing functions under constraints. For $\alpha > 1$, an analytical analysis becomes challenging. Numerical results show that the orderliness is valid until $\alpha > 140$, the second orderliness is valid until $\alpha > 78$, and the third orderliness is valid until $\alpha > 55$ (SI Text). It is instructive to consider that when $\alpha \rightarrow \infty$ the gamma distribution converges to a Gaussian distribution with mean $\mu = \alpha\lambda$ and variance $\sigma = \alpha\lambda^2$. The skewness of the gamma distribution, $\frac{\alpha+2}{\sqrt{\alpha(\alpha+1)}}$, is monotonic with respect to α , since $\frac{\partial \tilde{\mu}_3(\alpha)}{\partial \alpha} = \frac{-3\alpha-2}{2(\alpha(\alpha+1))^{3/2}} < 0$. When $\alpha = 55$, $\tilde{\mu}_3(\alpha) = 1.027$. Therefore, similar to the Weibull distribution, the parameters that let the distribution not be included in $\mathcal{P}_U \cap \mathcal{P}_O \cap \mathcal{P}_{O_2} \cap \mathcal{P}_{O_3}$ also correspond to cases when it is near-symmetric.

Theorem .7. Consider a γ -symmetric random variable X . Let it be transformed using a function $\phi(x)$ such that $\phi^{(2)}(x) \geq 0$ over the interval supported, the resulting convex transformed distribution is γ -ordered. Moreover, if the quantile function of X satisfies $Q^{(2)}(p) \leq 0$, the convex transformed distribution is second γ -ordered.

Proof. Let $\phi_{QA}(\epsilon) = \frac{1}{2}(\phi(Q(\gamma\epsilon)) + \phi(Q(1 - \epsilon)))$. Then, $\frac{\partial \phi_{QA}}{\partial \epsilon} = \frac{1}{2}(\gamma\phi'(Q(\gamma\epsilon))Q'(\gamma\epsilon) - \phi'(Q(1 - \epsilon))Q'(1 - \epsilon)) = \frac{1}{2}\gamma Q'(\gamma\epsilon)(\phi'(Q(\gamma\epsilon)) - \phi'(Q(1 - \epsilon))) \leq 0$, since for a γ -symmetric distribution, $Q(\frac{1}{1+\gamma}) - Q(\gamma\epsilon) = Q(1 - \epsilon) - Q(\frac{1}{1+\gamma})$, differentiating both sides, $-\gamma Q'(\gamma\epsilon) = -Q'(1 - \epsilon)$, where $Q'(p) \geq 0$, $\phi^{(2)}(x) \geq 0$. If further differentiating the equality, $\gamma^2 Q^{(2)}(\gamma\epsilon) = -Q^{(2)}(1 - \epsilon)$. Since $\frac{\partial^{(2)} \phi_{QA}}{\partial \epsilon^{(2)}} = \frac{1}{2}(\gamma^2 \phi^2(Q(\gamma\epsilon))(Q'(\gamma\epsilon))^2 + \phi^2(Q(1 - \epsilon))(Q'(1 - \epsilon))^2) + \frac{1}{2}(\gamma^2 \phi'(Q(\gamma\epsilon))(Q^2(\gamma\epsilon)) + \phi'(Q(1 - \epsilon))(Q^2(1 - \epsilon))) = \frac{1}{2}((\phi^{(2)}(Q(\gamma\epsilon)) + \phi^{(2)}(Q(1 - \epsilon)))(\gamma^2 Q'(\gamma\epsilon))^2) + \frac{1}{2}((\phi'(Q(\gamma\epsilon)) - \phi'(Q(1 - \epsilon)))\gamma^2 Q^{(2)}(\gamma\epsilon))$. If $Q^{(2)}(p) \leq 0$, $\frac{\partial^{(2)} \phi_{QA}}{\partial \epsilon^{(2)}} \geq 0$. \square

The mean-median-mode inequality for distributions of the powers and roots of the variates of a given distribution was investigated by Henry Rietz in 1927 (28), but the most straightforward solution is the exponential transformation since the derivatives are invariably positive. An application of Theorem .7 is that the lognormal distribution is ordered as it is exponentially transformed from the Gaussian distribution. The quantile function of the Gaussian distribution meets the condition $Q^{(2)}(p) = -2\sqrt{2\pi}\sigma e^{2\text{erfc}^{-1}(2p)^2} \text{erfc}^{-1}(2p) \leq 0$, where σ is the standard deviation of the Gaussian distribution and erfc denotes the complementary error function. Thus, the lognormal distribution is second ordered. Numerical results suggest that it is also third ordered, although analytically proving this result is challenging.

Theorem .7 also reveals a relation between convex transformation and orderliness, since ϕ is the non-decreasing convex function in van Zwet's trailblazing work *Convex transformations of random variables* (29) if adding an additional constraint that $\phi'(x) \geq 0$. Consider a near-symmetric distribution S , such that SQA_ϵ as a function of ϵ fluctuates from 0 to $\frac{1}{2}$, with $\mu = m$. By definition, S is not ordered. Let s be the pdf of S . Applying the transformation $\phi(x)$ to S decreases $s(Q_S(\epsilon))$, and the decrease rate, due to the order, is much smaller for $s(Q_S(1 - \epsilon))$. As a consequence, as the second derivative of $\phi(x)$ increases, eventually, after a point, $s(Q_S(\epsilon))$ becomes greater than $s(Q_S(1 - \epsilon))$ even if it was not previously. Thus, the SQA_ϵ function becomes monotonically decreasing, and S becomes ordered. Accordingly, in a family of distributions that differ by a skewness-increasing transformation in van Zwet's sense, violations of orderliness typically occur only when the distribution is near-symmetric.

Pearson proposed using the mean-median difference $\mu - m$ as a measure of skewness after standardization in 1895 (30). Bowley (1926) proposed a measure of skewness based on the SQA -median difference $\text{SQA}_\epsilon - m$ (31). Groeneveld and Meeden (1984) (32) generalized these measures of skewness based on van Zwet's convex transformation (29) while exploring their properties. A distribution is called monotonically right-skewed if and only if $\forall 0 \leq \epsilon_1 \leq \epsilon_2 \leq \frac{1}{2}$, $\text{SQA}_{\epsilon_1} - m \geq \text{SQA}_{\epsilon_2} - m$. Since m is a constant, the monotonic skewness is equivalent

to the orderliness. For a nonordered distribution, the signs of $\text{SQA}_\epsilon - m$ with different breakdown points might be different, implying that some skewness measures indicate left-skewed distribution, while others suggest right-skewed distribution. Although it seems reasonable that such a distribution is likely to be generally near-symmetric, however, counterexamples can be constructed. For example, consider the Weibull distribution, when $\alpha > \frac{1}{1-\ln(2)}$, it is near-symmetric and nonordered, the non-monotonicity of the SQA function arises when ϵ is close to $\frac{1}{2}$. Replacing the third quartile with one from a right-skewed heavy-tailed distribution leads to a right-skewed, heavy-tailed, and nonordered distribution. Therefore, the validity of robust measures of skewness based on the SQA-median difference is closely related to the orderliness of the distribution.

Remarkably, in 2018, Li, Shao, Wang, Yang (33) proved the bias bound of any quantile for arbitrary continuous distributions with finite second moments. Here, let $\mathcal{P}_{\mu,\sigma}$ denotes the set of continuous distributions whose mean is μ and standard deviation is σ , the bias upper bound of the quantile average is given in the following theorem.

Theorem .8. *The bias upper bound of the quantile average for any continuous distribution whose mean is zero and standard deviation is one is*

$$\sup_{P \in \mathcal{P}_{\mu=0, \sigma=1}} \text{QA}(\epsilon, \gamma) = \frac{1}{2} \left(\sqrt{\frac{\gamma\epsilon}{1-\gamma\epsilon}} + \sqrt{\frac{1-\epsilon}{\epsilon}} \right), \quad [3]$$

where $0 \leq \epsilon \leq \frac{1}{1+\gamma}$.

Proof. Since $\sup_{P \in \mathcal{P}_{\mu=0, \sigma=1}} \frac{1}{2}(Q(\gamma\epsilon) + Q(1-\epsilon)) \leq \frac{1}{2}(\sup_{P \in \mathcal{P}_{\mu=0, \sigma=1}} Q(\gamma\epsilon) + \sup_{P \in \mathcal{P}_{\mu=0, \sigma=1}} Q(1-\epsilon))$, the assertion follows directly from the Lemma 2.6 in (33). \square

In 2020, Bernard et al. (2) further refined these bounds for unimodal distributions and derived the bias bound of the symmetric quantile average. Here, the bias upper bound of the quantile average, $0 \leq \gamma < 5$, for $P \in \mathcal{P}_U \cap \mathcal{P}_{\mu=0, \sigma=1}$ is given as

$$\sup_{P \in \mathcal{P}_U \cap \mathcal{P}_{\mu=0, \sigma=1}} \text{QA}(\epsilon, \gamma) = \begin{cases} \frac{1}{2} \left(\sqrt{\frac{4}{9\epsilon} - 1} + \sqrt{\frac{3\gamma\epsilon}{4-3\gamma\epsilon}} \right) & 0 \leq \epsilon \leq \frac{1}{6} \\ \frac{1}{2} \left(\sqrt{\frac{3(1-\epsilon)}{4-3(1-\epsilon)}} + \sqrt{\frac{3\gamma\epsilon}{4-3\gamma\epsilon}} \right) & \frac{1}{6} < \epsilon \leq \frac{1}{1+\gamma} \end{cases}$$

The proof based on the bias bounds of any quantile (2) and the $\gamma \geq 5$ case are given in the SI Text. Subsequent theorems reveal the safeguarding role these bounds play in defining estimators based on ν th γ -orderliness. The proof of Theorem .9 is provided in the SI Text.

Theorem .9. $\sup_{P \in \mathcal{P}_{\mu=0, \sigma=1}} \text{QA}(\epsilon, \gamma)$ is monotonic decreasing with respect to ϵ over the interval $[0, \frac{1}{1+\gamma}]$, when $0 \leq \gamma \leq 1$.

Theorem .10. $\sup_{P \in \mathcal{P}_U \cap \mathcal{P}_{\mu=0, \sigma=1}} \text{QA}(\epsilon, \gamma)$ is monotonic decreasing with respect to ϵ over the interval $[0, \frac{1}{1+\gamma}]$, when $0 \leq \gamma \leq 1$.

Proof. When $0 \leq \epsilon \leq \frac{1}{6}$, $\frac{\partial \sup \text{QA}(\epsilon, \gamma)}{\partial \epsilon} = \frac{\gamma}{\sqrt{\frac{\epsilon\gamma}{12-9\epsilon\gamma}(4-3\epsilon\gamma)^2}} - \frac{1}{3\sqrt{\frac{4}{9\epsilon}-9\epsilon^2}}$. When $\gamma = 0$, $\frac{\partial \sup \text{QA}(\epsilon, \gamma)}{\partial \epsilon} = -\frac{1}{3\sqrt{\frac{4}{9\epsilon}-9\epsilon^2}} \leq 0$. When $\epsilon \rightarrow 0^+$, $\lim_{\epsilon \rightarrow 0^+} \left(\frac{\gamma}{(4-3\gamma\epsilon)^2 \sqrt{\frac{\epsilon\gamma}{12-9\epsilon\gamma}}} - \frac{1}{3\sqrt{\frac{4}{9\epsilon}-9\epsilon^2}} \right) =$

$\lim_{\epsilon \rightarrow 0^+} \left(\frac{\gamma\sqrt{3}}{\sqrt{43\epsilon\gamma}} - \frac{1}{6\sqrt{\epsilon^3}} \right) \rightarrow -\infty$. Assuming $\epsilon > 0$, when $0 < \gamma \leq 1$, to prove $\frac{\partial \sup \text{QA}(\epsilon, \gamma)}{\partial \epsilon} \leq 0$, it is equivalent to showing $\frac{\sqrt{\frac{\epsilon\gamma}{12-9\epsilon\gamma}(4-3\epsilon\gamma)^2}}{\gamma} \geq 3\sqrt{\frac{4}{9\epsilon}-9\epsilon^2}$. Define $L(\epsilon, \gamma) = \frac{\sqrt{\frac{\epsilon\gamma}{12-9\epsilon\gamma}(4-3\epsilon\gamma)^2}}{\gamma}$, $R(\epsilon, \gamma) = 3\sqrt{\frac{4}{9\epsilon}-9\epsilon^2}$. $\frac{L(\epsilon, \gamma)}{\epsilon^2} = \frac{\sqrt{\frac{\epsilon\gamma}{12-9\epsilon\gamma}(4-3\epsilon\gamma)^2}}{\gamma\epsilon^2} = \frac{1}{\gamma} \left(\frac{4}{\epsilon} - 3\gamma \right)^2 \sqrt{\frac{1}{12-9\epsilon\gamma}}$, $\frac{R(\epsilon, \gamma)}{\epsilon^2} = 3\sqrt{\frac{4}{9\epsilon}-9\epsilon^2}$. Then, $\frac{L(\epsilon, \gamma)}{\epsilon^2} \geq \frac{R(\epsilon, \gamma)}{\epsilon^2} \Leftrightarrow \frac{1}{\gamma} \sqrt{\frac{1}{12-9\epsilon\gamma}} \left(\frac{4}{\epsilon} - 3\gamma \right)^2 \geq 3\sqrt{\frac{4}{9\epsilon}-9\epsilon^2} \Leftrightarrow \frac{1}{\gamma} \left(\frac{4}{\epsilon} - 3\gamma \right)^2 \geq 3\sqrt{\frac{12}{\epsilon\gamma} - 9\sqrt{\frac{4}{9\epsilon}-9\epsilon^2}}$. Let $LmR\left(\frac{1}{\epsilon}\right) = \frac{1}{\gamma^2} \left(\frac{4}{\epsilon} - 3\gamma \right)^4 - 9 \left(\frac{12}{\epsilon\gamma} - 9 \right) \left(\frac{4}{\epsilon} - 9 \right)$. $\frac{\partial LmR(1/\epsilon)}{\partial (1/\epsilon)} = \frac{16(\frac{4}{\epsilon}-3\gamma)^3}{\gamma^2} - 36 \left(\frac{12}{\epsilon\gamma} - 9 \right) - \frac{108(4\frac{4}{\epsilon}-9)}{\gamma} = \frac{4(4(\frac{4}{\epsilon}-3\gamma)^3 - 27\gamma(\frac{4}{\epsilon}-3\gamma) + 27(9-\frac{4}{\epsilon})\gamma)}{\gamma^2} = \frac{4(256\frac{1}{\epsilon^3} - 576\frac{1}{\epsilon^2}\gamma + 432\frac{1}{\epsilon}\gamma^2 - 216\frac{1}{\epsilon}\gamma - 108\gamma^3 + 81\gamma^2 + 243\gamma)}{\gamma^2}$. Since $256\frac{1}{\epsilon^3} - 576\frac{1}{\epsilon^2}\gamma + 432\frac{1}{\epsilon}\gamma^2 - 216\frac{1}{\epsilon}\gamma - 108\gamma^3 + 81\gamma^2 + 243\gamma \geq 1536\frac{1}{\epsilon^2} - 576\frac{1}{\epsilon} + 432\frac{1}{\epsilon}\gamma^2 - 216\frac{1}{\epsilon}\gamma - 108\gamma^3 + 81\gamma^2 + 243\gamma \geq 924\frac{1}{\epsilon^2} + 36\frac{1}{\epsilon} - 216\frac{1}{\epsilon} + 432\frac{1}{\epsilon}\gamma^2 - 108\gamma^3 + 81\gamma^2 + 243\gamma \geq 924\frac{1}{\epsilon^2} + 36\frac{1}{\epsilon} - 216\frac{1}{\epsilon} + 513\gamma^2 - 108\gamma^3 + 243\gamma > 0$, $\frac{\partial LmR(1/\epsilon)}{\partial (1/\epsilon)} > 0$. Also, $LmR(6) = \frac{81(\gamma-8)((\gamma-8)^3+15\gamma)}{\gamma^2} > 0 \Leftrightarrow \gamma^4 - 32\gamma^3 + 399\gamma^2 - 2168\gamma + 4096 > 0$. Since $\gamma^4 > 0$, if $0 < \gamma \leq 1$, then $32\gamma^3 < 256$, it suffices to prove that $399\gamma^2 - 2168\gamma + 4096 > 256$. Applying the quadratic formula demonstrates the validity of this inequality. Hence, $LmR\left(\frac{1}{\epsilon}\right) \geq 0$ for $\epsilon \in (0, \frac{1}{6}]$, provided that $0 < \gamma \leq 1$. The first part is finished.

When $\frac{1}{6} < \epsilon \leq \frac{1}{1+\gamma}$, $\frac{\partial \sup \text{QA}(\epsilon, \gamma)}{\partial \epsilon} = \frac{\gamma}{\sqrt{\frac{\epsilon\gamma}{12-9\epsilon\gamma}(4-3\epsilon\gamma)^2}} - \frac{1}{\sqrt{1-\epsilon}(3\epsilon+1)^{\frac{3}{2}}} > 0$. When $\gamma = 0$, $\frac{\partial \sup \text{QA}(\epsilon, \gamma)}{\partial \epsilon} < 0$. For other cases, to determine whether $\frac{\partial \sup \text{QA}(\epsilon, \gamma)}{\partial \epsilon} \leq 0$, since $\sqrt{1-\epsilon}(3\epsilon+1)^{\frac{3}{2}} > 0$ and $\sqrt{\gamma\epsilon}(4-3\gamma\epsilon)^{\frac{3}{2}} > 0$, showing $\frac{\sqrt{\gamma\epsilon}(4-3\gamma\epsilon)^{\frac{3}{2}}}{\gamma} \geq \sqrt{1-\epsilon}(3\epsilon+1)^{\frac{3}{2}} \Leftrightarrow \frac{\gamma\epsilon(4-3\gamma\epsilon)^3}{\gamma^2} \geq (1-\epsilon)(3\epsilon+1)^3 \Leftrightarrow -27\gamma^2\epsilon^4 + 108\gamma\epsilon^3 + \frac{64\epsilon}{\gamma} + 27\epsilon^4 - 162\epsilon^2 - 8\epsilon - 1 \geq 0$ is sufficient. When $0 < \gamma \leq 1$, the inequality can be further simplified to $108\gamma\epsilon^3 + \frac{64\epsilon}{\gamma} - 162\epsilon^2 - 8\epsilon - 1 \geq 0$. Since $\epsilon \leq \frac{1}{1+\gamma}$, $\gamma \leq \frac{1}{\epsilon} - 1$. Also, as $0 < \gamma \leq 1$, $0 < \gamma \leq \min(1, \frac{1}{\epsilon} - 1)$. When $\frac{1}{6} < \epsilon \leq \frac{1}{2}$, $\frac{1}{\epsilon} - 1 > 1$, so $0 < \gamma \leq 1$. When $\frac{1}{2} \leq \epsilon < 1$, $0 < \gamma \leq \frac{1}{\epsilon} - 1$. Let $h(\gamma) = 108\gamma\epsilon^3 + \frac{64\epsilon}{\gamma}$, $\frac{\partial h(\gamma)}{\partial \gamma} = 108\epsilon^3 - \frac{64\epsilon}{\gamma^2}$. When $\gamma \leq \sqrt{\frac{64\epsilon}{18\epsilon^3}}$, $\frac{\partial h(\gamma)}{\partial \gamma} \geq 0$, when $\gamma \geq \sqrt{\frac{64\epsilon}{18\epsilon^3}}$, $\frac{\partial h(\gamma)}{\partial \gamma} \leq 0$, therefore, the minimum of $h(\gamma)$ must be when γ is equal to the boundary point of the domain. When $\frac{1}{6} < \epsilon \leq \frac{1}{2}$, $0 < \gamma \leq 1$, since $h(0) \rightarrow \infty$, $h(1) = 108\epsilon^3 + 64\epsilon$, the minimum occurs at the boundary point $\gamma = 1$, $108\gamma\epsilon^3 + \frac{64\epsilon}{\gamma} - 162\epsilon^2 - 8\epsilon - 1 > 108\epsilon^3 + 56\epsilon - 162\epsilon^2 - 1$. Let $g(\epsilon) = 108\epsilon^3 + 56\epsilon - 162\epsilon^2 - 1$. $g'(\epsilon) = 324\epsilon^2 - 324\epsilon + 56$, when $\epsilon \leq \frac{2}{9}$, $g'(\epsilon) \geq 0$, when $\frac{2}{9} \leq \epsilon \leq \frac{1}{2}$, $g'(\epsilon) \leq 0$, since $g(\frac{1}{6}) = \frac{13}{3}$, $g(\frac{1}{2}) = 0$, so $g(\epsilon) \geq 0$, the simplified inequality is satisfied. When $\frac{1}{2} \leq \epsilon < 1$, $0 < \gamma \leq \frac{1}{\epsilon} - 1$. Since $h(\frac{1}{\epsilon} - 1) = 108(\frac{1}{\epsilon} - 1)\epsilon^3 + \frac{64\epsilon}{\frac{1}{\epsilon} - 1} - 162\epsilon^2 - 8\epsilon - 1 > 108(\frac{1}{\epsilon} - 1)\epsilon^3 + \frac{64\epsilon}{\frac{1}{\epsilon} - 1} - 162\epsilon^2 - 8\epsilon - 1 = \frac{-108\epsilon^4 + 54\epsilon^3 - 18\epsilon^2 + 7\epsilon + 1}{\epsilon - 1}$. Let $nu(\epsilon) = -108\epsilon^4 + 54\epsilon^3 - 18\epsilon^2 + 7\epsilon + 1$, then $nu'(\epsilon) = -432\epsilon^3 + 162\epsilon^2 - 36\epsilon + 7$, $nu''(\epsilon) = -1296\epsilon^2 + 324\epsilon - 36 < 0$. Since $nu'(\epsilon = \frac{1}{2}) = -\frac{49}{2} < 0$, $nu'(\epsilon) < 0$. Also, $nu(\epsilon = \frac{1}{2}) = 0$,

so $nu(\epsilon) \leq 0$, the simplified inequality is also satisfied. As a result, the simplified inequality is also valid within the range of $\frac{1}{6} < \epsilon \leq \frac{1}{1+\gamma}$, provided that $0 < \gamma \leq 1$. Then, it validates $\frac{\partial \sup_{\epsilon} QA(\epsilon, \gamma)}{\partial \epsilon} \leq 0$ for the same range of ϵ and γ .

The first and second formulae, when $\epsilon = \frac{1}{6}$, are all equal

to $\frac{1}{2} \left(\frac{\sqrt{\frac{\gamma}{4-\gamma}}}{\sqrt{2}} + \sqrt{\frac{5}{3}} \right)$. It follows that $\sup QA(\epsilon, \gamma)$ is continuous over $[0, \frac{1}{1+\gamma}]$. Hence, $\frac{\partial \sup_{\epsilon} QA(\epsilon, \gamma)}{\partial \epsilon} \leq 0$ holds for the entire range $0 \leq \epsilon \leq \frac{1}{1+\gamma}$, when $0 \leq \gamma \leq 1$, which leads to the assertion of this theorem. \square

Let \mathcal{P}_γ^k denote the set of all continuous distributions whose moments, from the first to the k th, are all finite. For a right-skewed distribution, it suffices to consider the upper bound. The monotonicity of $\sup_{P \in \mathcal{P}_\gamma^2} QA$ with respect to ϵ implies that the extent of any violations of the γ -orderliness, if $0 \leq \gamma \leq 1$, is bounded for any distribution with a finite second moment, e.g., for a right-skewed distribution in \mathcal{P}_γ^2 , if $\exists 0 \leq \epsilon_1 \leq \epsilon_2 \leq \epsilon_3 \leq \frac{1}{1+\gamma}$, $QA_{\epsilon_2, \gamma} \geq QA_{\epsilon_3, \gamma} \geq QA_{\epsilon_1, \gamma}$, $QA_{\epsilon_2, \gamma}$ will not be too far away from $QA_{\epsilon_1, \gamma}$, since $\sup_{P \in \mathcal{P}_\gamma^2} QA_{\epsilon_1, \gamma} > \sup_{P \in \mathcal{P}_\gamma^2} QA_{\epsilon_2, \gamma} > \sup_{P \in \mathcal{P}_\gamma^2} QA_{\epsilon_3, \gamma}$. Moreover, a stricter bound can be established for unimodal distributions. The violation of ν th γ -orderliness, when $\nu \geq 2$, is also bounded as it corresponds to the higher-order derivatives of the QA function with respect to ϵ .

Data Availability. Data for Figure ?? are given in SI Dataset S1. All codes have been deposited in [GitHub](#).

ACKNOWLEDGMENTS. I sincerely acknowledge the insightful comments from the editor which considerably elevated the lucidity and merit of this paper.

1. CF Gauss, *Theoria combinationis observationum erroribus minimis obnoxiae*. (Henricus Dieterich), (1823).
2. C Bernard, R Kazzi, S Vanduffel, Range value-at-risk bounds for unimodal distributions under partial information. *Insur. Math. Econ.* **94**, 9–24 (2020).
3. P Daniell, Observations weighted according to order. *Am. J. Math.* **42**, 222–236 (1920).
4. JW Tukey, A survey of sampling from contaminated distributions in *Contributions to probability and statistics*. (Stanford University Press), pp. 448–485 (1960).
5. WJ Dixon, Simplified Estimation from Censored Normal Samples. *The Annals Math. Stat.* **31**, 385–391 (1960).
6. K Danielak, T Rychlik, Theory & methods: Exact bounds for the bias of trimmed means. *Aust. & New Zealand J. Stat.* **45**, 83–96 (2003).
7. M Bieniek, Comparison of the bias of trimmed and winsorized means. *Commun. Stat. Methods* **45**, 6641–6650 (2016).
8. J Hodges Jr, E Lehmann, Estimates of location based on rank tests. *The Annals Math. Stat.* **34**, 598–611 (1963).
9. F Wilcoxon, Individual comparisons by ranking methods. *Biom. Bull.* **1**, 80–83 (1945).
10. PJ Huber, Robust estimation of a location parameter. *Ann. Math. Stat.* **35**, 73–101 (1964).
11. Q Sun, WX Zhou, J Fan, Adaptive huber regression. *J. Am. Stat. Assoc.* **115**, 254–265 (2020).
12. T Mathieu, Concentration study of m-estimators using the influence function. *Electron. J. Stat.* **16**, 3695–3750 (2022).
13. AS Nemirovskij, DB Yudin, *Problem complexity and method efficiency in optimization*. (Wiley-Interscience), (1983).
14. D Hsu, S Sabato, Heavy-tailed regression with a generalized median-of-means in *International Conference on Machine Learning*. (PMLR), pp. 37–45 (2014).
15. L Devroye, M Lerasle, G Lugosi, RI Oliveira, Sub-gaussian mean estimators. *The Annals Stat.* **44**, 2695–2725 (2016).
16. P Laforgue, S Cléménçon, P Bertail, On medians of (randomized) pairwise means in *International Conference on Machine Learning*. (PMLR), pp. 1272–1281 (2019).
17. G Lecué, M Lerasle, Robust machine learning by median-of-means: theory and practice. (2020).
18. B Efron, Bootstrap methods: Another look at the jackknife. *The Annals Stat.* **7**, 1–26 (1979).
19. PJ Bickel, DA Freedman, Some asymptotic theory for the bootstrap. *The annals statistics* **9**, 1196–1217 (1981).
20. PJ Bickel, DA Freedman, Asymptotic normality and the bootstrap in stratified sampling. *The annals statistics* **12**, 470–482 (1984).
21. R Helmers, P Janssen, N Veraverbeke, *Bootstrapping U-quantiles*. (CWI. Department of Operations Research, Statistics, and System Theory [BS]), (1990).

22. J Neyman, On the two different aspects of the representative method: The method of stratified sampling and the method of purposive selection. *J. Royal Stat. Soc.* **97**, 558–606 (1934).
23. G McIntyre, A method for unbiased selective sampling, using ranked sets. *Aust. journal agricultural research* **3**, 385–390 (1952).
24. CM Stein, Efficient nonparametric testing and estimation in *Proceedings of the third Berkeley symposium on mathematical statistics and probability*. Vol. 1, pp. 187–195 (1956).
25. P Bickel, CA Klaassen, Y Ritov, JA Wellner, *Efficient and adaptive estimation for semiparametric models*. (Springer) Vol. 4, (1993).
26. JT Runnenburg, Mean, median, mode. *Stat. Neerlandica* **32**, 73–79 (1978).
27. Wv Zwet, Mean, median, mode ii. *Stat. Neerlandica* **33**, 1–5 (1979).
28. H Rietz, On certain properties of frequency distributions of the powers and roots of the variates of a given distribution. *Proc. Natl. Acad. Sci.* **13**, 817–820 (1927).
29. WR van Zwet, *Convex Transformations of Random Variables: Nebst Stellingen*. (1964).
30. K Pearson, X. contributions to the mathematical theory of evolution.—ii. skew variation in homogeneous material. *Philos. Transactions Royal Soc. London. (A.)* **186**, 343–414 (1895).
31. AL Bowley, *Elements of statistics*. (King) No. 8, (1926).
32. RA Groeneveld, G Meeden, Measuring skewness and kurtosis. *J. Royal Stat. Soc. Ser. D (The Stat.)* **33**, 391–399 (1984).
33. L Li, H Shao, R Wang, J Yang, Worst-case range value-at-risk with partial information. *SIAM J. on Financial Math.* **9**, 190–218 (2018).