

# Semiparametric robust mean estimations based on the orderliness of quantile averages

Tuban Lee

This manuscript was compiled on May 22, 2023

**As one of the most fundamental problems in statistics, robust location estimation has many prominent solutions, such as the symmetric trimmed mean, symmetric Winsorized mean, Hodges–Lehmann estimator, Huber  $M$ -estimator, and median of means. Recent studies suggest that their maximum biases concerning the mean can be quite different in asymmetric distributions, but the underlying mechanisms and average performance remain largely unclear. In this article, similar to the mean-median-mode inequality, it is proven that within the context of nearly all common unimodal distributions, there is an orderliness of symmetric quantile averages with varying breakdown points. Further deductions explain why the Winsorized mean and median of means typically have smaller biases compared to the trimmed mean. Building on the  $U$ -orderliness, the superiority of the median Hodges–Lehmann mean is discussed.**

semiparametric | mean-median-mode inequality | asymptotic | unimodal  
| Hodges–Lehmann estimator

In 1823, Gauss (1) proved that for any unimodal distribution with a finite second moment,  $|m - \mu| \leq \sqrt{\frac{3}{4}}\omega$ , where  $\mu$  is the population mean,  $m$  is the population median, and  $\omega$  is the root mean square deviation from the mode,  $M$ . This pioneering work revealed that despite potential bias with respect to the mean in robust estimates, the deviation remains bounded in unit of a scale parameter under certain assumptions. Bernard, Kazzi, and Vanduffel (2020) (2) further derived asymptotic bias bounds of any quantile for unimodal distributions with finite second moments by reducing this optimization problem to a parametric one, which can be solved analytically. They showed that the population median,  $m$ , has the smallest maximum distance to the population mean,  $\mu$ , among all symmetric quantile averages (SQA $_{\epsilon}$ ). Daniell, in 1920, (3) analyzed a class of estimators, linear combinations of order statistics, and identified that  $\epsilon$ -symmetric trimmed mean (STM $_{\epsilon}$ ) belongs to this class. Another popular choice, the  $\epsilon$ -symmetric Winsorized mean (SWM $_{\epsilon}$ ), named after Winsor and introduced by Tukey (4) and Dixon (5) in 1960, is also an  $L$ -estimator. Bieniek (2016) derived exact bias upper bounds of the Winsorized mean based on Danielak and Rychlik's work (2003) on the trimmed mean for any distribution with a finite second moment and confirmed that the former is smaller than the latter (6, 7). In 1963, Hodges and Lehmann (8) proposed a class of nonparametric location estimators based on rank tests and, from the Wilcoxon signed-rank statistic (9), deduced the median of pairwise means as a robust location estimator for a symmetric population. Both  $L$ -statistics and  $R$ -statistics achieve robustness essentially by removing a certain proportion of extreme values. In 1964, Huber (10) generalized maximum likelihood estimation to the minimization of the sum of a specific loss function, which measures the residuals between the data points and the model's parameters. Some  $L$ -estimators are also  $M$ -estimators, e.g., the sample mean is an  $M$ -estimator with a squared error loss function, the sample median is an

$M$ -estimator with an absolute error loss function (10). The Huber  $M$ -estimator is obtained by applying the Huber loss function that combines elements of both squared error and absolute error to achieve robustness against gross errors and high efficiency for contaminated Gaussian distributions (10). Sun, Zhou, and Fan (2020) examined the concentration bounds of Huber  $M$ -estimator (11). Mathieu (2022) (12) further derived the concentration bounds of  $M$ -estimators and demonstrated that, by selecting the tuning parameter which depends on the variance, Huber  $M$ -estimator can also be a sub-Gaussian estimator. The concept of median of means (MoM $_{k,b} = \frac{n}{k}$ ,  $k$  is the number of size in each block,  $b$  is the number of blocks) was implicitly introduced several times in Nemirovsky and Yudin (1983) (13), Jerrum, Valiant, and Vazirani (1986), (14) and Alon, Matias and Szegedy (1996) (15)'s works. Given its good performance even for distributions with infinite second moments, MoM has received increasing attention over the past decade (16–18). Devroye, Lerasle, Lugosi, and Oliveira (2016) showed that MoM nears the optimum of sub-Gaussian mean estimation with regards to concentration bounds when the distribution has a heavy tail (17). For a comparison of concentration bounds of trimmed mean, Huber  $M$ -estimator, median of means and other relevant estimators, readers are directed to Gobet, Lerasle, and Métivier's paper (2022) (19). Laforgue, Clemencon, and Bertail (2019) proposed the median of randomized means (MoRM $_{k,b}$ ) (18), wherein, rather than partitioning, an arbitrary number,  $b$ , of blocks are built independently from the sample, and showed that MoRM has a better non-asymptotic sub-Gaussian property compared to MoM. In fact, asymptotically, the Hodges–Lehmann (H-L) estimator is equivalent to MoM $_{k=2,b=\frac{n}{k}}$  and MoRM $_{k=2,b}$ , and they can be seen as the pairwise mean distribution is approximated by the sampling without replacement and bootstrap, respectively. For the asymptotic validity, readers are referred to the foundational works of Efron (1979) (20), Bickel and

## Significance Statement

In 1964, van Zwet introduced the convex transformation order for comparing the skewness of two distributions. This paradigm shift played a fundamental role in defining robust measures of distributions, from spread to kurtosis. Here, instead of examining the stochastic ordering between two distributions, the orderliness of quantile averages within a distribution is investigated. By classifying distributions through the signs of derivatives, a series of sophisticated robust mean estimators are deduced. Nearly all common nonparametric robust location estimators are found to be special cases thereof.

T.L. designed research, performed research, analyzed data, and wrote the paper.

The author declares no competing interest.

<sup>1</sup>To whom correspondence should be addressed. E-mail: tl@biomathematics.org

71 Freedman (1981, 1984) (21, 22), and Helmers, Janssen, and  
 72 Veraverbeke (1990) (23).

Here, the  $\epsilon, b$ -stratified mean is defined as

$$SM_{\epsilon, b, n} := \frac{b}{n} \left( \sum_{j=1}^{\frac{b-1}{2b\epsilon}} \sum_{i_j=\frac{(2bj-b-1)n\epsilon}{b-1}+1}^{\frac{(2bj-b+1)n\epsilon}{b-1}} X_{i_j} \right),$$

73 where  $X_1 \leq \dots \leq X_n$  denote the order statistics of a sample  
 74 of  $n$  independent and identically distributed random variables  
 75  $X_1, \dots, X_n$ .  $b \in \mathbb{N}$ ,  $b \geq 3$ . The definition was further refined to  
 76 guarantee the continuity of the breakdown point by incorporat-  
 77 ing an additional block in the center when  $\lfloor \frac{b-1}{2b\epsilon} \rfloor \bmod 2 = 0$ ,  
 78 or by adjusting the central block when  $\lfloor \frac{b-1}{2b\epsilon} \rfloor \bmod 2 = 1$  (SI  
 79 Text). If the subscript  $n$  is omitted, only the asymptotic  
 80 behavior is considered. If  $b$  is omitted,  $b = 3$  is assumed.  
 81  $SM_{\epsilon, b=3}$  is equivalent to  $STM_{\epsilon}$ , when  $\epsilon > \frac{1}{6}$ . The basic idea  
 82 of the stratified mean, when  $\frac{b-1}{2\epsilon} \in \mathbb{N}$ ,  $b \bmod 2 = 1$ , is to dis-  
 83 tribute the data into  $\frac{b-1}{2\epsilon}$  equal-sized non-overlapping blocks  
 84 according to their order, then further sequentially group these  
 85 blocks into  $b$  equal-sized strata and compute the mean of the  
 86 middle stratum, which is the median of means of each stratum.  
 87 In situations where  $i \bmod 1 \neq 0$ , a potential solution is to  
 88 generate multiple smaller samples that satisfy the equality  
 89 by sampling without replacement, and subsequently calculate  
 90 the mean of all estimations. The details of determining the  
 91 sample size and sampling times are provided in the SI Text.  
 92 Although the principle resembles that of the median of means,  
 93 without the random shift,  $SM_{\epsilon, b, n}$  is different from  $MoM_{k=\frac{n}{b}, b}$ .  
 94 Additionally, the stratified mean differs from the mean of the  
 95 sample obtained through stratified sampling methods, intro-  
 96 duced by Neyman (1934) (24) or ranked set sampling (25),  
 97 introduced by McIntyre in 1952, as these sampling methods  
 98 aim to obtain more representative samples or improve the  
 99 efficiency of sample estimates, but the sample means based  
 100 on them are not robust. When  $b \bmod 2 = 1$ , the stratified  
 101 mean can be regarded as replacing the other equal-sized strata  
 102 with the middle stratum, which, in principle, is analogous to  
 103 the Winsorized mean that replaces extreme values with less  
 104 extreme percentiles. Furthermore, while the bounds confirm  
 105 that the Winsorized mean and median of means outperform  
 106 the trimmed mean (6, 7, 17, 19) in worst-case performance,  
 107 the complexity of bound analysis makes it difficult to achieve a  
 108 complete and intuitive understanding of these results. Also, a  
 109 clear explanation for the average performance of them remains  
 110 elusive. The aim of this paper is to define a series of semi-  
 111 parametric models using the signs of derivatives, reveal their  
 112 elegant interrelations and connections to parametric models,  
 113 and show that by exploiting these models, a set of sophisti-  
 114 cated mean estimators can be deduced, which exhibit strong  
 115 robustness to departures from assumptions.

## 116 Quantile average and weighted average

The symmetric trimmed mean, symmetric Winsorized mean, and stratified mean are all  $L$ -estimators. More specifically, they are symmetric weighted averages, which are defined as

$$SWA_{\epsilon, n} := \frac{\sum_{i=1}^{\lceil \frac{n}{2} \rceil} \frac{X_i + X_{n-i+1}}{2} w_i}{\sum_{i=1}^{\lceil \frac{n}{2} \rceil} w_i},$$

where  $w_i$ s are the weights applied to the symmetric quantile  
 averages according to the definition of the corresponding  $L$ -  
 estimators. For example, for the  $\epsilon$ -symmetric trimmed mean,  
 $w_i = \begin{cases} 0, & i < n\epsilon \\ 1, & i \geq n\epsilon \end{cases}$ , provided that  $n\epsilon \in \mathbb{N}$ . The mean and  
 median are indeed two special cases of the symmetric trimmed  
 mean.

To extend the symmetric quantile average to the asymmet-  
 ric case, there are two possible definitions for the  $\epsilon, \gamma$ -quantile  
 average (QA( $\epsilon, \gamma, n$ )), i.e.,

$$\frac{1}{2}(\hat{Q}_n(\gamma\epsilon) + \hat{Q}_n(1 - \epsilon)), \quad [1]$$

and

$$\frac{1}{2}(\hat{Q}_n(\epsilon) + \hat{Q}_n(1 - \gamma\epsilon)), \quad [2]$$

where  $\gamma \geq 0$  and  $0 \leq \epsilon \leq \frac{1}{1+\gamma}$ ,  $\hat{Q}_n(p)$  is the empirical quantile  
 function. For trimming from both sides, [1] and [2] are essen-  
 tially equivalent. [1] is assumed in this article unless otherwise  
 specified, since many common asymmetric distributions are  
 right-skewed, and [1] allows trimming only from the right side  
 by setting  $\gamma = 0$ .

Analogously, the weighted average can be defined as

$$WA_{\epsilon, \gamma, n} := \frac{\int_{\epsilon_0=0}^{\frac{1}{1+\gamma}} QA(\epsilon_0, \gamma, n) w_{\epsilon_0}}{\int_{\epsilon_0=0}^{\frac{1}{1+\gamma}} w_{\epsilon_0}}.$$

For instance, the  $\epsilon, \gamma$ -trimmed mean ( $TM_{\epsilon, \gamma, n}$ ) is a weighted  
 average with a left trim size of  $\gamma\epsilon n$  and a right trim size of  $\epsilon n$ ,  
 where  $w_{\epsilon_0} = \begin{cases} 0, & \epsilon_0 < \epsilon \\ 1, & \epsilon_0 \geq \epsilon \end{cases}$ . Using this definition, even  $\gamma\epsilon n \notin \mathbb{N}$   
 or  $\epsilon n \notin \mathbb{N}$ , the TM computation remains unaltered since this  
 definition is based on the empirical quantile function. However,  
 considering the computational cost in practice, here, the non-  
 asymptotic definitions of various types of weighted averages,  
 in most cases, are essentially based on order statistics. The  
 solution to the decimal issue of them is the same as that in  
 SM, unless stated otherwise.

## Classifying distributions by the signs of derivatives

Let  $\mathcal{P}_k$  denote the set of all distributions over  $\mathbb{R}$  whose mo-  
 ments, from the first to the  $k$ th, are all finite. Without loss  
 of generality, the discussion of all the classes outlined below  
 is restricted to the intersection with the nonparametric class  
 of distributions  $\mathcal{P}_\gamma^k := \{\text{All continuous distribution } P \in \mathcal{P}_k\}$ .  
 Besides fully and smoothly parameterizing by a Euclidean pa-  
 rameter or just assuming regularity conditions, there are many  
 ways to classify distributions. In 1956, Stein initiated the  
 problem of estimating parameters in the presence of an infinite  
 dimensional nuisance shape parameter (26). A notable exam-  
 ple discussed in his groundbreaking work was the estimation  
 of the center of symmetry for an unknown symmetric distribu-  
 tion. In 1993, Bickel, Klaassen, Ritov, and Wellner published  
 an influential semiparametrics textbook (27) which system-  
 atically categorized many common models into three classes:  
 parametric, nonparametric, and semiparametric. Yet, there  
 is another old and commonly encountered class of distribu-  
 tions that receives little attention in semiparametric literature:  
 the unimodal distribution. It is a very unique semiparametric

model because its definition is based on the signs of derivatives, i.e., for a continuous distribution,  $(f'(x) > 0 \text{ for } x \leq M) \wedge (f'(x) < 0 \text{ for } x \geq M)$ . Let  $\mathcal{P}_U$  denote the set of all unimodal distributions. There was a widespread misbelief that the median of an arbitrary unimodal distribution always lies between its mean and mode until Runnenburg (1978) and van Zwet (1979) (28, 29) endeavored to determine sufficient conditions for the inequality to hold, thereby implying the possibility of its violation. The class of distributions that satisfy the mean-median-mode inequality constitutes a subclass of  $\mathcal{P}_U$ . By analogy, a right-skewed distribution is called  $\gamma$ -ordered, if and only if

$$\forall 0 \leq \epsilon_1 \leq \epsilon_2 \leq \frac{1}{1+\gamma}, \text{QA}_{\epsilon_1, \gamma} \geq \text{QA}_{\epsilon_2, \gamma}.$$

The necessary and sufficient condition below hints at the relation between the mean-median-mode inequality and the  $\gamma$ -orderliness.

**Theorem .1.** Let  $P_\gamma^k$  represent an arbitrary distribution in the set  $\mathcal{P}_\gamma^k$ .  $P_\gamma^k$  is  $\gamma$ -ordered if and only if the probability density function (pdf) satisfies the inequality  $f(Q(\gamma\epsilon)) \geq f(Q(1-\epsilon))$  for all  $0 \leq \epsilon \leq \frac{1}{1+\gamma}$  or  $f(Q(\gamma\epsilon)) \leq f(Q(1-\epsilon))$  for all  $0 \leq \epsilon \leq \frac{1}{1+\gamma}$ , where  $\gamma \geq 0$ .

*Proof.* Without loss of generality, consider the case of right-skewed continuous distribution. From the definition of  $\gamma$ -orderliness, it is deduced that  $\frac{Q(\gamma\epsilon-\delta)+Q(1-\epsilon+\delta)}{2} \geq \frac{Q(\gamma\epsilon)+Q(1-\epsilon)}{2} \Leftrightarrow Q(\gamma\epsilon-\delta)-Q(\gamma\epsilon) \geq Q(1-\epsilon)-Q(1-\epsilon+\delta) \Leftrightarrow Q'(1-\epsilon) \geq Q'(\gamma\epsilon)$ , where  $\delta$  is an infinitesimal positive quantity. Observing that the quantile function is the inverse function of the cumulative distribution function (cdf),  $Q'(1-\epsilon) \geq Q'(\gamma\epsilon) \Leftrightarrow F'(Q(\gamma\epsilon)) \geq F'(Q(1-\epsilon))$ , thereby completing the proof, given that the derivative of cdf is pdf.  $\square$

According to Theorem .1, if a probability distribution is right-skewed and monotonic, it will always be  $\gamma$ -ordered, provided  $\gamma \geq 0$ . For a right-skewed continuous unimodal distribution, if  $Q(\gamma\epsilon) > M$ , the inequality  $f(Q(\gamma\epsilon)) \geq f(Q(1-\epsilon))$  holds. The principle is extendable to unimodal-like distributions. Suppose there is a right-skewed continuous multimodal distribution following the mean- $\gamma$ -median-first mode inequality with several smaller modes on the right side, with the first mode,  $M$ , having the greatest probability density, and the  $\gamma$ -median,  $Q(\frac{\gamma}{1+\gamma})$ , falling within the first dominant mode (i.e., if  $x > Q(\frac{\gamma}{1+\gamma})$ ,  $f(Q(\frac{\gamma}{1+\gamma})) \geq f(x)$ ), then if  $Q(\gamma\epsilon) > M$ , the inequality  $f(Q(\gamma\epsilon)) \geq f(Q(1-\epsilon))$  also holds. In other words, while a distribution following the mean- $\gamma$ -median-mode inequality may not be strictly  $\gamma$ -ordered, the inequality that defines  $\gamma$ -orderliness remains valid for most quantile averages. The mean- $\gamma$ -median-mode inequality can also indicate possible bounds for  $\gamma$  in practice, e.g., for any distributions, when  $\gamma \rightarrow \infty$ , the  $\gamma$ -median will be greater than the mean and the mode, when  $\gamma \rightarrow 0$ , the  $\gamma$ -median will be smaller than the mean and the mode.

Consider the sign of the derivative of the quantile average with respect to the breakdown point, the above definition of  $\gamma$ -orderliness can also be expressed as

$$\forall 0 \leq \epsilon \leq \frac{1}{1+\gamma}, \frac{\partial \text{QA}_{\epsilon, \gamma}}{\partial \epsilon} \leq 0.$$

The left-skewed case can be obtained by reversing the inequality  $\frac{\partial \text{QA}_{\epsilon, \gamma}}{\partial \epsilon} \leq 0$  to  $\frac{\partial \text{QA}_{\epsilon, \gamma}}{\partial \epsilon} \geq 0$  and employing the second definition of QA, as given in [2]; for simplicity, it will be omitted in the following discussion. If  $\gamma = 1$ , the  $\gamma$ -ordered distribution is referred to as ordered.

Furthermore, many common right-skewed distributions are partial bounded, indicating a convex behavior of the QA function when  $\epsilon \rightarrow 0$ . By further assuming convexity, the second  $\gamma$ -orderliness can be defined as follows for a right-skewed distribution,

$$\forall 0 \leq \epsilon \leq \frac{1}{1+\gamma}, \frac{\partial^2 \text{QA}_{\epsilon, \gamma}}{\partial \epsilon^2} \geq 0 \wedge \frac{\partial \text{QA}_{\epsilon, \gamma}}{\partial \epsilon} \leq 0.$$

Analogously, the  $\nu$ th  $\gamma$ -orderliness of a right-skewed distribution can be defined as  $(-1)^\nu \frac{\partial^\nu \text{QA}_{\epsilon, \gamma}}{\partial \epsilon^\nu} \geq 0 \wedge \dots \wedge -\frac{\partial \text{QA}_{\epsilon, \gamma}}{\partial \epsilon} \geq 0$ . If  $\gamma = 1$ , the  $\nu$ th  $\gamma$ -orderliness is referred as  $\nu$ th orderliness. Let  $\mathcal{P}_O$  denote the set of all distributions that are ordered and  $\mathcal{P}_{O_\nu}$  and  $\mathcal{P}_{\gamma O_\nu}$  represent the sets of all distributions that are  $\nu$ th ordered and  $\nu$ th  $\gamma$ -ordered, respectively. When the shape parameter of the Weibull distribution,  $\alpha$ , is smaller than 3.258, it can be shown that the Weibull distribution belong to  $\mathcal{P}_U \cap \mathcal{P}_O \cap \mathcal{P}_{O_2} \cap \mathcal{P}_{O_3}$  (SI Text). At  $\alpha \approx 3.602$ , the Weibull distribution is symmetric, and as  $\alpha \rightarrow \infty$ , the skewness of the Weibull distribution reaches 1. Therefore, the parameters that let it not be included in the set correspond to cases when it is near-symmetric, as shown in the SI Text. Nevertheless, computing the derivatives of the QA function is often intricate and, at times, challenging. The following theorems establish the relationship between  $\mathcal{P}_O$ ,  $\mathcal{P}_{O_\nu}$ , and  $\mathcal{P}_{\gamma O_\nu}$ , and a wide range of other semi-parametric distributions. They can be used to quickly identify some parametric distributions in  $\mathcal{P}_O$ ,  $\mathcal{P}_{O_\nu}$ , and  $\mathcal{P}_{\gamma O_\nu}$ .

**Theorem .2.** For any random variable  $X$  whose probability distribution function belongs to a location-scale family, the distribution is  $\nu$ th  $\gamma$ -ordered if and only if the family of probability distributions is  $\nu$ th  $\gamma$ -ordered.

*Proof.* Let  $Q_0$  denote the quantile function of the standard distribution without any shifts or scaling. After a location-scale transformation, the quantile function is  $Q(p) = \lambda Q_0(p) + \mu$ , where  $\lambda$  is the scale parameter and  $\mu$  is the location parameter. According to the definition of the  $\nu$ th  $\gamma$ -orderliness, the signs of derivatives of the QA function are invariant after this transformation. As the location-scale transformation is reversible, the proof is complete.  $\square$

Theorem .2 demonstrates that in the analytical proof of the  $\nu$ th  $\gamma$ -orderliness of a parametric distribution, both the location and scale parameters can be regarded as constants. It is also instrumental in proving other theorems, as illustrated below.

**Theorem .3.** Any symmetric distribution with a finite second moment is  $\nu$ th ordered.

*Proof.* Without loss of generality, assuming continuity and  $m = 0$ . A symmetric distribution is a probability distribution such that for all  $x$ ,  $f(x) = f(-x)$ . Its cdf satisfies  $F(x) = 1 - F(-x)$ . Let  $x = Q(p)$ , then,  $F(Q(p)) = p = 1 - F(-Q(p))$  and  $F(Q(1-p)) = 1 - p \Leftrightarrow p = 1 - F(Q(1-p))$ . Therefore,  $F(-Q(p)) = F(Q(1-p))$ . Since the cdf is monotonic,  $-Q(p) = Q(1-p) \Leftrightarrow Q(p) + Q(1-p) = 0$ . As a result, all symmetric



quantile averages coincide; the  $\nu$ th order derivative is zero. The case of  $m \neq 0$  follows directly from Theorem 2.  $\square$

As a consequence of Theorem 3 and the fact that generalized Gaussian distribution is symmetric around the median, it is  $\nu$ th ordered.

**Theorem 4.** Any continuous right-skewed distribution whose quantile function  $Q$  satisfies  $Q^{(\nu)}(p) \geq 0 \wedge \dots \wedge Q^{(i)}(p) \geq 0 \wedge \dots \wedge Q^{(2)}(p) \geq 0$ ,  $i \bmod 2 = 0$ , is  $\nu$ th  $\gamma$ -ordered, provided that  $0 \leq \gamma \leq 1$ .

*Proof.* Since  $(-1)^i \frac{\partial^i Q_{A_{\epsilon}, \gamma}}{\partial \epsilon^i} = \frac{1}{2}((- \gamma)^i Q^i(\gamma \epsilon) + Q^i(1 - \epsilon))$ , for  $0 \leq \epsilon \leq \frac{1}{1+\gamma}$  and  $1 \leq i \leq \nu$ , when  $i \bmod 2 = 0$ ,  $(-1)^i \frac{\partial^i Q_{A_{\epsilon}, \gamma}}{\partial \epsilon^i} \geq 0$  for all  $\gamma \geq 0$ . When  $i \bmod 2 = 1$ , if further assuming  $0 \leq \gamma \leq 1$ ,  $(-1)^i \frac{\partial^i Q_{A_{\epsilon}, \gamma}}{\partial \epsilon^i} \geq 0$ , since  $Q^{(i+1)}(p) \geq 0$ .  $\square$

It is now straightforward to prove that the Pareto distribution follows the  $\nu$ th  $\gamma$ -orderliness, provided that  $0 \leq \gamma \leq 1$ , since the quantile function of the Pareto distribution is  $Q(p) = x_m(1-p)^{-\frac{1}{\alpha}}$ , where  $x_m > 0$ ,  $\alpha > 0$ , and so  $Q^{(\nu)}(p) \geq 0$  for all  $\nu \in \mathbb{N}$  according to the chain rule.

**Theorem 5.** A right-skewed continuous distribution with a monotonic decreasing pdf is second  $\gamma$ -ordered.

*Proof.* A monotonic decreasing pdf implies  $f'(x) = F^{(2)}(x) \leq 0$ . Since  $Q'(p) \geq 0$ , let  $x = Q(F(x))$ , then by differentiating both sides of the equation twice, one can obtain  $0 = Q^{(2)}(F(x))(F'(x))^2 + Q'(F(x))F^{(2)}(x) \Leftrightarrow Q^{(2)}(F(x)) = -\frac{Q'(F(x))F^{(2)}(x)}{(F'(x))^2} \geq 0$ . The desired result is derived from Theorem 1 and 4.  $\square$

Theorem 5 provides valuable insights into the relation between modality and orderliness. The conventional definition states that a distribution with a monotonic pdf is still considered unimodal. However, within its supported interval, the mode number is zero. The number of modes and their magnitudes within a distribution are closely related to the possibility of orderliness being valid, although counterexamples can always be constructed for non-monotonic distributions. A proof of the second  $\gamma$ -orderliness, if  $\gamma > 0$ , can be easily established for the gamma distributions when  $\alpha \leq 1$  as the pdf of the gamma distribution is  $f(x) = \frac{\lambda^{-\alpha} x^{\alpha-1} e^{-\frac{x}{\lambda}}}{\Gamma(\alpha)}$ , where  $x \geq 0$ ,  $\lambda > 0$ ,  $\alpha > 0$ ,  $\Gamma$  is the gamma function, it is a product of two monotonic decreasing functions under constraints. For  $\alpha > 1$ , the proof becomes challenging. Numerical results show that the orderliness is valid until  $\alpha > 140$ , the second orderliness is valid until  $\alpha > 78$ , and the third orderliness is valid until  $\alpha > 55$  (SI Text). It is instructive to consider that when  $\alpha \rightarrow \infty$  the gamma distribution converges to a Gaussian distribution with mean  $\mu = \alpha\lambda$  and variance  $\sigma = \alpha\lambda^2$ . The skewness of the gamma distribution,  $\frac{\alpha+2}{\sqrt{\alpha(\alpha+1)}}$ , is monotonic with respect to  $\alpha$ , since  $\frac{\partial \mu_3(\alpha)}{\partial \alpha} = \frac{-3\alpha-2}{2(\alpha(\alpha+1))^{3/2}} < 0$ . When  $\alpha = 55$ ,  $\mu_3(\alpha) = 1.027$ . Therefore, similar to the Weibull distribution, the parameters that let the distribution not be included in  $\mathcal{P}_U \cap \mathcal{P}_O \cap \mathcal{P}_{O_2} \cap \mathcal{P}_{O_3}$  also correspond to cases when it is near-symmetric.

**Theorem 6.** Consider a symmetric random variable  $X$ . Let it be transformed using a function  $\phi(x)$  such that  $\phi^{(2)}(x) \geq 0 \wedge \phi'(x) \geq 0$  over the interval supported, the resulting convex transformed distribution is ordered. Moreover, if the quantile function of  $X$  satisfies  $Q^{(2)}(\epsilon) \leq 0$ , the convex transformed distribution is second ordered.

*Proof.* Let  $\phi\text{SQA}(\epsilon) = \frac{1}{2}(\phi(Q(\epsilon)) + \phi(Q(1-\epsilon)))$ ,  $0 \leq \epsilon \leq \frac{1}{2}$ . Then,  $\frac{d\phi\text{SQA}}{d\epsilon} = \frac{1}{2}(\phi'(Q(\epsilon))Q'(\epsilon) - \phi'(Q(1-\epsilon))Q'(1-\epsilon)) = \frac{1}{2}Q'(\epsilon)(\phi'(Q(\epsilon)) - \phi'(Q(1-\epsilon))) \leq 0$ , since for a symmetric distribution,  $m - Q(\epsilon) = Q(1-\epsilon) - m$ , differentiating both sides,  $-Q'(\epsilon) = -Q'(1-\epsilon)$ , where  $Q'(\epsilon) \geq 0$ ,  $\phi^{(2)}(x) \geq 0$ . If further differentiating the equality,  $Q^{(2)}(\epsilon) = -Q^{(2)}(1-\epsilon)$ . Since  $\frac{d^{(2)}\phi\text{SQA}}{d\epsilon^{(2)}} = \frac{1}{2}(\phi^2(Q(\epsilon))(Q'(\epsilon))^2 + \phi^2(Q(1-\epsilon))(Q'(1-\epsilon))^2) + \frac{1}{2}(\phi'(Q(\epsilon))(Q^2(\epsilon)) + \phi'(Q(1-\epsilon))(Q^2(1-\epsilon))) = \frac{1}{2}((\phi^{(2)}(Q(\epsilon)) + \phi^{(2)}(Q(1-\epsilon)))(Q'(\epsilon))^2) + \frac{1}{2}((\phi'(Q(\epsilon)) - \phi'(Q(1-\epsilon)))Q^{(2)}(\epsilon))$ . If  $Q^{(2)}(\epsilon) \leq 0$ ,  $\frac{d^{(2)}\phi\text{SQA}}{d\epsilon^{(2)}} \geq 0$ .  $\square$

The mean-median-mode inequality for distributions of the powers and roots of the variates of a given distribution was investigated by Henry Rietz in 1927 (30), but the most straightforward solution is the exponential transformation since the derivatives are invariably positive. An application of Theorem 6 is that the lognormal distribution is ordered as it is exponentially transformed from the Gaussian distribution. The quantile function of the Gaussian distribution meets the condition  $Q^{(2)}(\epsilon) = -2\sqrt{2}\pi\sigma e^{2\text{erfc}^{-1}(2\epsilon)^2}\text{erfc}^{-1}(2\epsilon) \leq 0$ , where  $\sigma$  is the standard deviation,  $\text{erfc}$  denotes the complementary error function. Thus, the lognormal distribution is second ordered. Numerical results suggest that it is also third ordered, although an analytical proof is challenging.

Theorem 6 also reveals a relation between convex transformation and orderliness, since  $\phi$  is the non-decreasing convex function in van Zwet's trailblazing work *Convex transformations of random variables* (31). Consider a near-symmetric distribution  $S$ , such that  $\text{SQA}_\epsilon$  as a function of  $\epsilon$  fluctuates from 0 to  $\frac{1}{2}$ , with  $\mu = m$ . By definition,  $S$  is not ordered. Let  $s$  be the pdf of  $S$ . Applying the transformation  $\phi(x)$  to  $S$  decreases  $s(Q_S(\epsilon))$ , and the decrease rate, due to the order, is much smaller for  $s(Q_S(1-\epsilon))$ . As a consequence, as the second derivative of  $\phi(x)$  increases, eventually, after a point,  $s(Q_S(\epsilon))$  becomes greater than  $s(Q_S(1-\epsilon))$  even if it was not previously. Thus, the  $\text{SQA}_\epsilon$  function becomes monotonically decreasing, and  $S$  becomes ordered. Accordingly, in a family of distributions that differ by a skewness-increasing transformation in van Zwet's sense, violations of orderliness typically occur only when the distribution is near-symmetric.

Pearson proposed using the mean-median difference  $\mu - m$  as a measure of skewness after standardization in 1895 (32). Bowley (1926) proposed a measure of skewness based on the  $\text{SQA}$ -median difference  $\text{SQA}_\epsilon - m$  (33). Groeneveld and Meeden (1984) (34) generalized these measures of skewness based on van Zwet's convex transformation (31) while exploring their properties. A distribution is called monotonically right-skewed if and only if  $\forall 0 \leq \epsilon_1 \leq \epsilon_2 \leq \frac{1}{2}$ ,  $\text{SQA}_{\epsilon_1} - m \geq \text{SQA}_{\epsilon_2} - m$ . Since  $m$  is a constant, the monotonic skewness is equivalent to the orderliness. For a nonordered distribution, the signs of

SQA $_{\epsilon} - m$  with different breakdown points might be different, implying that some skewness measures indicate left-skewed distribution, while others suggest right-skewed distribution. Although it seems reasonable that such a distribution is likely to be generally near-symmetric, however, counterexamples can be constructed. For example, consider the Weibull distribution, when  $\alpha > \frac{1}{1-\ln(2)}$ , it is near-symmetric and nonordered, the non-monotonicity of the SQA function arises when  $\epsilon$  is close to  $\frac{1}{2}$ . Replacing the third quartile with one from a right-skewed heavy-tailed distribution leads to a right-skewed, heavy-tailed, and nonordered distribution. Therefore, the validity of robust measures of skewness based on the SQA-median difference is closely related to the orderliness of the distribution.

Remarkably, in 2020, Bernard et al. (2) proved the bias bounds of any quantile for  $P \in \mathcal{P}_U \cap \mathcal{P}_T^2$ . They further derived the bias bound of the symmetric quantile average. Here, let  $\mathcal{P}_{\mu, \sigma}$  denotes the set of continuous distributions whose mean is  $\mu$  and standard deviation is  $\sigma$ , the bias upper bound of the quantile average,  $0 \leq \gamma < 5$ , is given as

$$\sup_{P \in \mathcal{P}_U \cap \mathcal{P}_{\mu=0, \sigma=1}} \text{QA}(\epsilon, \gamma) = \begin{cases} \frac{1}{2} \left( \sqrt{\frac{4}{9} - 1} + \sqrt{\frac{3\gamma\epsilon}{4-3\gamma\epsilon}} \right) & 0 \leq \epsilon \leq \frac{1}{6} \\ \frac{1}{2} \left( \sqrt{\frac{3(1-\epsilon)}{4-3(1-\epsilon)}} + \sqrt{\frac{3\gamma\epsilon}{4-3\gamma\epsilon}} \right) & \frac{1}{6} < \epsilon \leq \frac{1}{1+\gamma} \end{cases}$$

The proof based on the bias bounds of any quantile (2) and the  $\gamma \geq 5$  case are given in the SI Text. The next theorem highlights its safeguarding role in defining estimators based on  $\nu$ th  $\gamma$ -orderliness.

**Theorem .7.** *The above bias upper bound function,  $\sup_{P \in \mathcal{P}_U \cap \mathcal{P}_{\mu=0, \sigma=1}} \text{QA}(\epsilon, \gamma)$ , is monotonic decreasing with respect to  $\epsilon$  over the interval  $[0, \frac{1}{1+\gamma}]$ , when  $0 \leq \gamma \leq 1$ .*

*Proof.* When  $0 \leq \epsilon \leq \frac{1}{6}$ ,  $\frac{\partial \sup \text{QA}(\epsilon, \gamma)}{\partial \epsilon} = \frac{\gamma}{\sqrt{\frac{\epsilon\gamma}{12-9\epsilon\gamma}(4-3\epsilon\gamma)^2}} - \frac{1}{3\sqrt{\frac{4}{9} - 9\epsilon^2}}$ . When  $\gamma = 0$ ,  $\frac{\partial \sup \text{QA}(\epsilon, \gamma)}{\partial \epsilon} = -\frac{1}{3\sqrt{\frac{4}{9} - 9\epsilon^2}} \leq 0$ .

When  $\epsilon \rightarrow 0^+$ ,  $\lim_{\epsilon \rightarrow 0^+} \left( \frac{\gamma}{(4-3\gamma\epsilon)^2 \sqrt{\frac{\epsilon\gamma}{12-9\epsilon\gamma}}} - \frac{1}{3\sqrt{\frac{4}{9} - 9\epsilon^2}} \right) =$

$\lim_{\epsilon \rightarrow 0^+} \left( \frac{\gamma\sqrt{3}}{4^{3/2}\epsilon\gamma} - \frac{1}{6\sqrt{\epsilon^3}} \right) \rightarrow -\infty$ . Assuming  $\epsilon > 0$ , when

$0 < \gamma \leq 1$ , to prove  $\frac{\partial \sup \text{QA}(\epsilon, \gamma)}{\partial \epsilon} \leq 0$ , it is equivalent to showing  $\sqrt{\frac{\epsilon\gamma}{12-9\epsilon\gamma}(4-3\epsilon\gamma)^2} \geq 3\sqrt{\frac{4}{9} - 9\epsilon^2}$ . Define

$$L(\epsilon, \gamma) = \frac{\sqrt{\frac{\epsilon\gamma}{12-9\epsilon\gamma}(4-3\epsilon\gamma)^2}}{\gamma}, R(\epsilon, \gamma) = 3\sqrt{\frac{4}{9} - 9\epsilon^2}. \quad \frac{L(\epsilon, \gamma)}{\epsilon^2} = \frac{\sqrt{\frac{\epsilon\gamma}{12-9\epsilon\gamma}(4-3\epsilon\gamma)^2}}{\gamma\epsilon^2} = \frac{1}{\gamma} \left( \frac{4}{\epsilon} - 3\gamma \right)^2 \sqrt{\frac{1}{\frac{12}{\epsilon\gamma} - 9}}, \quad \frac{R(\epsilon, \gamma)}{\epsilon^2} = 3\sqrt{\frac{4}{9} - 9\epsilon^2}.$$

$$\text{Then, } \frac{L(\epsilon, \gamma)}{\epsilon^2} \geq \frac{R(\epsilon, \gamma)}{\epsilon^2} \Leftrightarrow \frac{1}{\gamma} \sqrt{\frac{1}{\frac{12}{\epsilon\gamma} - 9}} \left( \frac{4}{\epsilon} - 3\gamma \right)^2 \geq 3\sqrt{\frac{4}{9} - 9\epsilon^2} \Leftrightarrow$$

$$\frac{1}{\gamma} \left( \frac{4}{\epsilon} - 3\gamma \right)^2 \geq 3\sqrt{\frac{12}{\epsilon\gamma} - 9} \sqrt{\frac{4}{9} - 9\epsilon^2}. \quad \text{Let } LmR\left(\frac{1}{\epsilon}\right) =$$

$$\frac{1}{\gamma^2} \left( \frac{4}{\epsilon} - 3\gamma \right)^4 - 9 \left( \frac{12}{\epsilon\gamma} - 9 \right) \left( \frac{4}{\epsilon} - 9 \right). \quad \frac{\partial LmR(1/\epsilon)}{\partial (1/\epsilon)} = \frac{16 \left( \frac{4}{\epsilon} - 3\gamma \right)^3}{\gamma^2} -$$

$$36 \left( \frac{12}{\epsilon\gamma} - 9 \right) - \frac{108 \left( \frac{4}{\epsilon} - 9 \right)}{\gamma} = \frac{4 \left( 4 \left( \frac{4}{\epsilon} - 3\gamma \right)^3 - 27\gamma \left( \frac{4}{\epsilon} - 3\gamma \right) + 27 \left( 9 - \frac{4}{\epsilon} \right) \gamma \right)}{\gamma^2} =$$

$$\frac{4 \left( 256 \frac{1}{\epsilon^3} - 576 \frac{1}{\epsilon^2} \gamma + 432 \frac{1}{\epsilon} \gamma^2 - 216 \frac{1}{\epsilon} \gamma - 108 \gamma^3 + 81 \gamma^2 + 243 \gamma \right)}{\gamma^2}. \quad \text{Since}$$

$$256 \frac{1}{\epsilon^3} - 576 \frac{1}{\epsilon^2} \gamma + 432 \frac{1}{\epsilon} \gamma^2 - 216 \frac{1}{\epsilon} \gamma - 108 \gamma^3 + 81 \gamma^2 + 243 \gamma \geq$$

$$1536 \frac{1}{\epsilon^2} - 576 \frac{1}{\epsilon} \gamma^2 + 432 \frac{1}{\epsilon} \gamma^2 - 216 \frac{1}{\epsilon} \gamma - 108 \gamma^3 + 81 \gamma^2 + 243 \gamma \geq$$

$$924 \frac{1}{\epsilon^2} + 36 \frac{1}{\epsilon} \gamma^2 - 216 \frac{1}{\epsilon} \gamma + 432 \frac{1}{\epsilon} \gamma^2 - 108 \gamma^3 + 81 \gamma^2 + 243 \gamma \geq$$

$$924 \frac{1}{\epsilon^2} + 36 \frac{1}{\epsilon} \gamma^2 - 216 \frac{1}{\epsilon} \gamma + 513 \gamma^2 - 108 \gamma^3 + 243 \gamma > 0, \quad \frac{\partial LmR(1/\epsilon)}{\partial (1/\epsilon)} > 0. \quad \text{Also, } LmR(6) = \frac{81(\gamma-8)((\gamma-8)^3+15\gamma)}{\gamma^2} >$$

$0 \Leftrightarrow \gamma^4 - 32\gamma^3 + 399\gamma^2 - 2168\gamma + 4096 > 0$ . Since  $\gamma^4 > 0$ , if  $0 < \gamma \leq 1$ , then  $32\gamma^3 < 256$ , it suffices to prove that  $399\gamma^2 - 2168\gamma + 4096 > 256$ . Applying the quadratic formula demonstrates the validity of this inequality. Hence,  $LmR\left(\frac{1}{\epsilon}\right) \geq 0$  for  $\epsilon \in (0, \frac{1}{6}]$ , provided that  $0 < \gamma \leq 1$ . The first part is finished.

When  $\frac{1}{6} < \epsilon \leq \frac{1}{1+\gamma}$ ,  $\frac{\partial \sup \text{QA}(\epsilon, \gamma)}{\partial \epsilon} = \sqrt{3} \left( \frac{\gamma}{\sqrt{\gamma\epsilon(4-3\gamma\epsilon)^{\frac{3}{2}}}} - \frac{1}{\sqrt{1-\epsilon(3\epsilon+1)^{\frac{3}{2}}}} \right)$ . When  $\gamma = 0$ ,

$$\frac{\gamma}{\sqrt{\gamma\epsilon(4-3\gamma\epsilon)^{\frac{3}{2}}}} = \frac{\sqrt{\gamma}}{\sqrt{\epsilon(4-3\gamma\epsilon)^{\frac{3}{2}}}} = 0, \quad \frac{\partial \sup \text{QA}(\epsilon, \gamma)}{\partial \epsilon} < 0. \quad \text{For}$$

other cases, to determine whether  $\frac{\partial \sup \text{QA}(\epsilon, \gamma)}{\partial \epsilon} \leq 0$ , since  $\sqrt{1-\epsilon(3\epsilon+1)^{\frac{3}{2}}} > 0$  and  $\sqrt{\gamma\epsilon(4-3\gamma\epsilon)^{\frac{3}{2}}} > 0$ , showing

$$\frac{\sqrt{\gamma\epsilon(4-3\gamma\epsilon)^{\frac{3}{2}}}}{\gamma} \geq \sqrt{1-\epsilon(3\epsilon+1)^{\frac{3}{2}}} \Leftrightarrow \frac{\gamma\epsilon(4-3\gamma\epsilon)^3}{\gamma^2} \geq (1-\epsilon)(3\epsilon+1)^3 \Leftrightarrow -27\gamma^2\epsilon^4 + 108\gamma\epsilon^3 + \frac{64\epsilon}{\gamma} + 27\epsilon^4 - 162\epsilon^2 - 8\epsilon - 1 \geq 0$$

is sufficient. When  $0 < \gamma \leq 1$ , the inequality can be further simplified to  $108\gamma\epsilon^3 + \frac{64\epsilon}{\gamma} - 162\epsilon^2 - 8\epsilon - 1 \geq 0$ . Since  $\epsilon \leq \frac{1}{1+\gamma}$ ,

$\gamma \leq \frac{1}{\epsilon} - 1$ . Also, as  $0 < \gamma \leq 1$ ,  $0 < \gamma \leq \min(1, \frac{1}{\epsilon} - 1)$ . When  $\frac{1}{6} < \epsilon \leq \frac{1}{2}$ ,  $\frac{1}{\epsilon} - 1 > 1$ , so  $0 < \gamma \leq 1$ . When

$\frac{1}{2} \leq \epsilon < 1$ ,  $0 < \gamma \leq \frac{1}{\epsilon} - 1$ . Let  $h(\gamma) = 108\gamma\epsilon^3 + \frac{64\epsilon}{\gamma}$ ,

$\frac{1}{2} \leq \epsilon < 1$ ,  $0 < \gamma \leq \frac{1}{\epsilon} - 1$ . Let  $h(\gamma) = 108\gamma\epsilon^3 + \frac{64\epsilon}{\gamma}$ ,  $\frac{\partial h(\gamma)}{\partial \gamma} = 108\epsilon^3 - \frac{64\epsilon}{\gamma^2}$ . When  $\gamma \leq \sqrt{\frac{64\epsilon}{18\epsilon^3}}$ ,  $\frac{\partial h(\gamma)}{\partial \gamma} \geq 0$ , when

$\gamma \geq \sqrt{\frac{64\epsilon}{18\epsilon^3}}$ ,  $\frac{\partial h(\gamma)}{\partial \gamma} \leq 0$ , therefore, the minimum of  $h(\gamma)$  must be when  $\gamma$  is equal to the boundary point of the domain. When  $\frac{1}{6} < \epsilon \leq \frac{1}{2}$ ,  $0 < \gamma \leq 1$ , since  $h(0) \rightarrow \infty$ ,

$h(1) = 108\epsilon^3 + 64\epsilon$ , the minimum occurs at the boundary point  $\gamma = 1$ ,  $108\gamma\epsilon^3 + \frac{64\epsilon}{\gamma} - 162\epsilon^2 - 8\epsilon - 1 > 108\epsilon^3 + 56\epsilon - 162\epsilon^2 - 1$ .

Let  $g(\epsilon) = 108\epsilon^3 + 56\epsilon - 162\epsilon^2 - 1$ .  $g'(\epsilon) = 324\epsilon^2 - 324\epsilon + 56$ , when  $\epsilon \leq \frac{2}{9}$ ,  $g'(\epsilon) \geq 0$ , when  $\frac{2}{9} \leq \epsilon \leq \frac{1}{2}$ ,  $g'(\epsilon) \leq 0$ , since

$g(\frac{1}{6}) = \frac{13}{3}$ ,  $g(\frac{1}{2}) = 0$ , so  $g(\epsilon) \geq 0$ , the simplified inequality is satisfied. When  $\frac{1}{2} \leq \epsilon < 1$ ,  $0 < \gamma \leq \frac{1}{\epsilon} - 1$ . Since

$h(\frac{1}{\epsilon} - 1) = 108(\frac{1}{\epsilon} - 1)\epsilon^3 + \frac{64\epsilon}{\frac{1}{\epsilon} - 1}$ ,  $108\gamma\epsilon^3 + \frac{64\epsilon}{\gamma} - 162\epsilon^2 - 8\epsilon - 1 >$

$$108 \left( \frac{1}{\epsilon} - 1 \right) \epsilon^3 + \frac{64\epsilon}{\frac{1}{\epsilon} - 1} - 162\epsilon^2 - 8\epsilon - 1 = \frac{-108\epsilon^4 + 54\epsilon^3 - 18\epsilon^2 + 7\epsilon + 1}{\epsilon - 1}.$$

Let  $nu(\epsilon) = -108\epsilon^4 + 54\epsilon^3 - 18\epsilon^2 + 7\epsilon + 1$ , then  $nu'(\epsilon) = -432\epsilon^3 + 162\epsilon^2 - 36\epsilon + 7$ ,  $nu''(\epsilon) = -1296\epsilon^2 + 324\epsilon - 36 < 0$ .

Since  $nu'(\epsilon = \frac{1}{2}) = -\frac{49}{2} < 0$ ,  $nu'(\epsilon) < 0$ . Also,  $nu(\epsilon = \frac{1}{2}) = 0$ , so  $nu(\epsilon) \leq 0$ , the simplified inequality is also satisfied. As a result, the simplified inequality is also valid within the range of  $\frac{1}{6} < \epsilon \leq \frac{1}{1+\gamma}$ , provided that  $0 < \gamma \leq 1$ . Then, it validates

$\frac{\partial \sup \text{QA}(\epsilon, \gamma)}{\partial \epsilon} \leq 0$  for the same range of  $\epsilon$  and  $\gamma$ .

The first and second formulae, when  $\epsilon = \frac{1}{6}$ , are all equal to  $\frac{1}{2} \left( \sqrt{\frac{4}{9} - \frac{\gamma}{2}} + \sqrt{\frac{5}{3}} \right)$ . It follows that  $\sup \text{QA}(\epsilon, \gamma)$  is continuous over  $[0, \frac{1}{1+\gamma}]$ . Hence,  $\frac{\partial \sup \text{QA}(\epsilon, \gamma)}{\partial \epsilon} \leq 0$  holds for the entire range  $0 \leq \epsilon \leq \frac{1}{1+\gamma}$ , when  $0 \leq \gamma \leq 1$ , which leads to the assertion of this theorem.  $\square$

For a right-skewed distribution, considering the upper bound is enough. The monotonicity of  $\sup_{P \in \mathcal{P}_U \cap \mathcal{P}_T^2} \text{QA}$  implies that the extent of any violations of the  $\gamma$ -orderliness, if  $0 \leq \gamma \leq 1$ , is bounded for a unimodal distribution with a finite second moment, e.g., for a right-skewed unimodal distribution in  $\mathcal{P}_T^2$ , if  $\exists 0 \leq \epsilon_1 \leq \epsilon_2 \leq \epsilon_3 \leq \frac{1}{1+\gamma}$ ,  $\text{QA}_{\epsilon_2} \geq \text{QA}_{\epsilon_3} \geq \text{QA}_{\epsilon_1}$ ,  $\text{QA}_{\epsilon_2}$  will not be too far away from  $\text{QA}_{\epsilon_1}$ , since  $\sup_{P \in \mathcal{P}_U \cap \mathcal{P}_T^2} \text{QA}_{\epsilon_1} > \sup_{P \in \mathcal{P}_U \cap \mathcal{P}_T^2} \text{QA}_{\epsilon_2} > \sup_{P \in \mathcal{P}_U \cap \mathcal{P}_T^2} \text{QA}_{\epsilon_3}$ . The  $\nu$ th  $\gamma$ -orderliness, when  $\nu \geq 2$ , corresponds to the higher order derivatives of the QA function, so

its violation is also bounded by  $\sup_{P \in \mathcal{P}_U \cap \mathcal{P}_Y^2} \text{QA}$  for  $\mathcal{P}_U \cap \mathcal{P}_Y^2$ .

### Inequalities related to weighted averages

The bias bound of the  $\epsilon$ -symmetric trimmed mean also exhibits monotonicity for  $\mathcal{P}_U \cap \mathcal{P}_Y^2$ , as proven in the SI Text using the formulae provided in Bernard et al.'s paper (2). So far, it appears clear that the bias of an estimator is closely related to its degree of robustness. In a right-skewed unimodal distribution, it is often observed that  $\mu \geq \text{STM}_\epsilon \geq m$ . Analogous to the  $\gamma$ -orderliness, the  $\gamma$ -trimming inequality is defined as  $\forall 0 \leq \epsilon_1 \leq \epsilon_2 \leq \frac{1}{1+\gamma}$ ,  $\text{TM}_{\epsilon_1, \gamma} \geq \text{TM}_{\epsilon_2, \gamma}$ . A necessary and sufficient condition for the  $\gamma$ -trimming inequality is the monotonic decrease of the bias of the  $\epsilon, \gamma$ -trimmed mean as a function of the breakdown point  $\epsilon$  for a right-skewed distribution, as proven in the SI Text. While  $\gamma$ -orderliness is a sufficient condition for the  $\gamma$ -trimming inequality, as proven in the SI Text, it is not necessary.

**Theorem .8.** *For a distribution that is right-skewed and follows the  $\gamma$ -trimming inequality, it is asymptotically true that the quantile average is always greater or equal to the corresponding trimmed mean with the same  $\epsilon$  and  $\gamma$ , provided that  $0 \leq \epsilon \leq \frac{1}{1+\gamma}$  and  $\gamma \geq 0$ .*

*Proof.* Assume, without loss of generality, that the distribution is continuous. According to the definition of the  $\gamma$ -trimming inequality:  $\frac{1}{1-\epsilon-\gamma\epsilon+2\delta} \int_{\gamma\epsilon-\delta}^{1-\epsilon+\delta} Q(u) du \geq \frac{1}{1-\epsilon-\gamma\epsilon} \int_{\gamma\epsilon}^{1-\epsilon} Q(u) du$ , if  $0 \leq \epsilon \leq \frac{1}{1+\gamma}$  and  $\gamma \geq 0$ , where  $\delta$  is an infinitesimal positive quantity. Subsequently, rewriting the inequality gives  $\int_{\gamma\epsilon-\delta}^{1-\epsilon+\delta} Q(u) du - \frac{1-\epsilon-\gamma\epsilon+2\delta}{1-\epsilon-\gamma\epsilon} \int_{\gamma\epsilon}^{1-\epsilon} Q(u) du \geq 0 \Leftrightarrow \int_{1-\epsilon}^{1-\epsilon+\delta} Q(u) du + \int_{\gamma\epsilon-\delta}^{\gamma\epsilon} Q(u) du - \frac{2\delta}{1-\epsilon-\gamma\epsilon} \int_{\gamma\epsilon}^{1-\epsilon} Q(u) du \geq 0$ . Since  $\delta \rightarrow 0^+$ ,  $\frac{1}{2\delta} \left( \int_{1-\epsilon}^{1-\epsilon+\delta} Q(u) du + \int_{\gamma\epsilon-\delta}^{\gamma\epsilon} Q(u) du \right) = \frac{Q(\gamma\epsilon) + Q(1-\epsilon)}{2} \geq \frac{1}{1-\epsilon-\gamma\epsilon} \int_{\gamma\epsilon}^{1-\epsilon} Q(u) du$ , the proof is complete.  $\square$

An analogous result can be obtained in the following theorem.

**Theorem .9.** *For a right-skewed continuous distribution following the  $\gamma$ -trimming inequality, asymptotically, the Winsorized mean is always greater or equal to the corresponding trimmed mean with the same  $\epsilon$  and  $\gamma$ , provided that  $0 \leq \epsilon \leq \frac{1}{1+\gamma}$  and  $0 \leq \gamma \leq 1$ .*

*Proof.* According to Theorem .8,  $\frac{Q(\gamma\epsilon) + Q(1-\epsilon)}{2} \geq \frac{1}{1-\epsilon-\gamma\epsilon} \int_{\gamma\epsilon}^{1-\epsilon} Q(u) du \Leftrightarrow \gamma\epsilon(Q(\gamma\epsilon) + Q(1-\epsilon)) \geq \left( \frac{2\gamma\epsilon}{1-\epsilon-\gamma\epsilon} \right) \int_{\gamma\epsilon}^{1-\epsilon} Q(u) du$ . Then, if  $1 \geq \gamma \geq 0$ ,  $\left( 1 - \frac{1}{1-\epsilon-\gamma\epsilon} \right) \int_{\gamma\epsilon}^{1-\epsilon} Q(u) du + \gamma\epsilon(Q(\gamma\epsilon) + Q(1-\epsilon)) \geq 0 \Rightarrow \int_{\gamma\epsilon}^{1-\epsilon} Q(u) du + \gamma\epsilon Q(\gamma\epsilon) + \epsilon Q(1-\epsilon) \geq \int_{\gamma\epsilon}^{1-\epsilon} Q(u) du + \gamma\epsilon(Q(\gamma\epsilon) + Q(1-\epsilon)) \geq \frac{1}{1-\epsilon-\gamma\epsilon} \int_{\gamma\epsilon}^{1-\epsilon} Q(u) du$ , the proof is complete.  $\square$

Assuming  $\gamma$ -orderliness, the result in Theorem .9 can be extended to the  $\gamma > 1$  case, as proven in the SI Text. Replacing the trimmed mean in the  $\gamma$ -trimming inequality with Winsorized mean forms the definition of the  $\gamma$ -Winsorization inequality.  $\gamma$ -orderliness also implies the  $\gamma$ -Winsorization inequality when  $0 \leq \gamma \leq 1$ , as proven in the SI Text.

To construct weighted averages based on the  $\gamma$ -orderliness, let  $\mathcal{B}_i = \int_{i\epsilon}^{(i+1)\epsilon} \text{QA}(u, \gamma) du$ ,  $ka = k\epsilon + c$ . It follows from the  $\gamma$ -orderliness that,  $-\frac{\partial \text{QA}_{\epsilon, \gamma}}{\partial \epsilon} \geq 0 \Rightarrow \forall 0 \leq a \leq 2a \leq \frac{1}{1+\gamma}$ ,  $-\frac{(\text{QA}(2a, \gamma) - \text{QA}(a, \gamma))}{a} \geq 0 \Rightarrow \mathcal{B}_i - \mathcal{B}_{i+1} \geq 0$ . Suppose  $\mathcal{B}_i = \mathcal{B}_0$ , then, based on the  $\gamma$ -orderliness, the  $\epsilon, \gamma$ -block Winsorized mean, is defined here for comparison in the SI Dataset S1 as

$$\text{BWM}_{\epsilon, \gamma, n} := \frac{1}{n} \left( \sum_{i=n\gamma\epsilon+1}^{(1-\epsilon)n} X_i + \sum_{i=n\gamma\epsilon+1}^{2n\gamma\epsilon+1} X_i + \sum_{i=(1-2\epsilon)n}^{(1-\epsilon)n} X_i \right),$$

which is double weighting the leftest and rightest blocks having the size  $\gamma\epsilon n$  and  $\epsilon n$ . Since their sizes are different, the  $0 \leq \gamma \leq 1$  is still necessary for the  $\gamma$ -block Winsorization inequality. If  $\gamma$  is omitted,  $\gamma = 1$  is assumed. This terminology is the same for other weighted averages. The solutions for the  $i \bmod 1 \neq 0$  case are the same as that in the SM. From the second  $\gamma$ -orderliness,  $\frac{\partial^2 \text{QA}_{\epsilon, \gamma}}{\partial^2 \epsilon} \geq 0 \Rightarrow \forall 0 \leq a \leq 2a \leq 3a \leq \frac{1}{1+\gamma}$ ,  $\frac{1}{a} \left( \frac{(\text{QA}(3a, \gamma) - \text{QA}(2a, \gamma))}{a} - \frac{(\text{QA}(2a, \gamma) - \text{QA}(a, \gamma))}{a} \right) \geq 0 \Rightarrow \mathcal{B}_i - 2\mathcal{B}_{i+1} + \mathcal{B}_{i+2} \geq 0$ . So, based on the second orderliness,  $\text{SM}_\epsilon$  can be seen as assuming  $\gamma = 1$ , replacing the two blocks,  $\mathcal{B}_i + \mathcal{B}_{i+2}$  with one block  $2\mathcal{B}_{i+1}$ . From the  $\nu$ th  $\gamma$ -orderliness, the recurrence relation of the derivatives naturally produces the alternating binomial coefficients,

$$\begin{aligned} (-1)^\nu \frac{\partial^\nu \text{QA}_{\epsilon, \gamma}}{\partial \epsilon^\nu} &\geq 0 \Rightarrow \forall 0 \leq a \leq \dots \leq (\nu+1)a \leq \frac{1}{1+\gamma}, \\ \frac{(-1)^\nu}{a} \left( \frac{\text{QA}(\nu a + a, \gamma)}{a} - \dots - \frac{\text{QA}(2a, \gamma)}{a} - \frac{\text{QA}(\nu a, \gamma)}{a} - \dots - \frac{\text{QA}(a, \gamma)}{a} \right) \\ &\geq 0 \Leftrightarrow \frac{(-1)^\nu}{a^\nu} \left( \sum_{j=0}^{\nu} (-1)^j \binom{\nu}{j} \text{QA}((\nu-j+1)a, \gamma) \right) \geq 0 \\ &\Rightarrow \sum_{j=0}^{\nu} (-1)^j \binom{\nu}{j} \mathcal{B}_{i+j} \geq 0. \end{aligned}$$

Based on the  $\nu$ th orderliness, the  $\epsilon$ -binomial mean is introduced as

$$\text{BM}_{\nu, \epsilon, n} := \frac{1}{n} \left( \sum_{i=1}^{\frac{1}{2}\epsilon^{-1}(\nu+1)-1} \sum_{j=0}^{\nu} \left( 1 - (-1)^j \binom{\nu}{j} \right) \mathfrak{B}_{i_j} \right),$$

where  $\mathfrak{B}_{i_j} = \sum_{l=n\epsilon(j+(i-1)(\nu+1)+1)}^{n\epsilon(j+(i-1)(\nu+1)+1)} (X_l + X_{n-l+1})$ . If  $\nu$  is not indicated, it is default as  $\nu = 3$ . Since the alternating sum of binomial coefficients equals zero, when  $\nu \ll \epsilon^{-1}$ ,  $\epsilon \rightarrow 0$ ,  $\text{BM} \rightarrow \mu$ . If  $\frac{1}{2}\epsilon^{-1}(\nu+1) \in \mathbb{N}$ , the asymmetry case is dividing the sample into  $\epsilon^{-1}$  blocks in the same way as SM and then further weighting each block using binomial coefficients ( $0 \leq \gamma \leq 1$  is needed). The solutions for the continuity of the breakdown point and  $l \bmod 1 \neq 0$  are the same as that in SM and not repeated here. The equality  $\text{BM}_{\nu=1, \epsilon} = \text{BWM}_\epsilon$  holds, and similarly,  $\text{BM}_{\nu=2, \epsilon} = \text{SM}_{\epsilon, b=3}$ , when  $\gamma = 1$  and their respective  $\epsilon$ s are identical. Interestingly, the biases of the  $\text{SM}_{\epsilon=\frac{1}{3}, b=3}$  and the  $\text{WM}_{\epsilon=\frac{1}{3}}$  are nearly indistinguishable in common asymmetric unimodal distributions such as Weibull, gamma, lognormal, and Pareto (SI Text), indicating that their robustness to departures from the symmetry assumption is practically similar. The reason is that the Winsorized mean is

506 using two single quantiles to replace the trimmed parts, not  
 507 two blocks. The subsequent theorem provides an explanation  
 508 for this difference.

509 **Data Availability.** Data for Figure ?? are given in SI Dataset  
 510 S1. All codes have been deposited in [GitHub](#).

511 **ACKNOWLEDGMENTS.** I sincerely acknowledge the insightful  
 512 comments from the editor which considerably elevated the lucidity  
 513 and merit of this paper.

- 514 1. CF Gauss, *Theoria combinationis observationum erroribus minimis obnoxiae*. (Henricus  
 515 Dieterich), (1823).
- 516 2. C Bernard, R Kazzi, S Vanduffel, Range value-at-risk bounds for unimodal distributions under  
 517 partial information. *Insur. Math. Econ.* **94**, 9–24 (2020).
- 518 3. P Daniell, Observations weighted according to order. *Am. J. Math.* **42**, 222–236 (1920).
- 519 4. JW Tukey, A survey of sampling from contaminated distributions in *Contributions to probability*  
 520 *and statistics*. (Stanford University Press), pp. 448–485 (1960).
- 521 5. WJ Dixon, Simplified Estimation from Censored Normal Samples. *The Annals Math. Stat.* **31**,  
 522 385 – 391 (1960).
- 523 6. K Danielak, T Rychlik, Theory & methods: Exact bounds for the bias of trimmed means. *Aust.*  
 524 *& New Zealand J. Stat.* **45**, 83–96 (2003).
- 525 7. M Bieniek, Comparison of the bias of trimmed and winsorized means. *Commun. Stat. Methods*  
 526 **45**, 6641–6650 (2016).
- 527 8. J Hodges Jr, E Lehmann, Estimates of location based on rank tests. *The Annals Math. Stat.*  
 528 **34**, 598–611 (1963).
- 529 9. F Wilcoxon, Individual comparisons by ranking methods. *Biom. Bull.* **1**, 80–83 (1945).
- 530 10. PJ Huber, Robust estimation of a location parameter. *Ann. Math. Stat.* **35**, 73–101 (1964).
- 531 11. Q Sun, WX Zhou, J Fan, Adaptive huber regression. *J. Am. Stat. Assoc.* **115**, 254–265 (2020).
- 532 12. T Mathieu, Concentration study of m-estimators using the influence function. *Electron. J. Stat.*  
 533 **16**, 3695–3750 (2022).
- 534 13. AS Nemirovskij, DB Yudin, *Problem complexity and method efficiency in optimization*. (Wiley-  
 535 Interscience), (1983).
- 536 14. MR Jerrum, LG Valiant, VV Vazirani, Random generation of combinatorial structures from a  
 537 uniform distribution. *Theor. computer science* **43**, 169–188 (1986).
- 538 15. N Alon, Y Matias, M Szegedy, The space complexity of approximating the frequency moments  
 539 in *Proceedings of the twenty-eighth annual ACM symposium on Theory of computing*. pp.  
 540 20–29 (1996).
- 541 16. PL Bühlmann, Bagging, subagging and bragging for improving some prediction algorithms  
 542 in *Research report/Seminar für Statistik, Eidgenössische Technische Hochschule (ETH)*.  
 543 (Seminar für Statistik, Eidgenössische Technische Hochschule (ETH), Zürich), Vol. 113,  
 544 (2003).
- 545 17. L Devroye, M Lerasle, G Lugosi, RI Oliveira, Sub-gaussian mean estimators. *The Annals Stat.*  
 546 **44**, 2695–2725 (2016).
- 547 18. P Laforge, S Cléménçon, P Bertail, On medians of (randomized) pairwise means in *Interna-*  
 548 *tional Conference on Machine Learning*. (PMLR), pp. 1272–1281 (2019).
- 549 19. E Gobet, M Lerasle, D Métivier, Mean estimation for Randomized Quasi Monte Carlo method.  
 550 working paper or preprint (2022).
- 551 20. B Efron, Bootstrap methods: Another look at the jackknife. *The Annals Stat.* **7**, 1–26 (1979).
- 552 21. PJ Bickel, DA Freedman, Some asymptotic theory for the bootstrap. *The annals statistics* **9**,  
 553 1196–1217 (1981).
- 554 22. PJ Bickel, DA Freedman, Asymptotic normality and the bootstrap in stratified sampling. *The*  
 555 *annals statistics* **12**, 470–482 (1984).
- 556 23. R Helmers, P Janssen, N Veraverbeke, *Bootstrapping U-quantiles*. (CWI. Department of  
 557 Operations Research, Statistics, and System Theory [BS]), (1990).
- 558 24. J Neyman, On the two different aspects of the representative method: The method of stratified  
 559 sampling and the method of purposive selection. *J. Royal Stat. Soc.* **97**, 558–606 (1934).
- 560 25. G McIntyre, A method for unbiased selective sampling, using ranked sets. *Aust. journal*  
 561 *agricultural research* **3**, 385–390 (1952).
- 562 26. C Stein, , et al., Efficient nonparametric testing and estimation in *Proceedings of the third*  
 563 *Berkeley symposium on mathematical statistics and probability*. Vol. 1, pp. 187–195 (1956).
- 564 27. P Bickel, CA Klaassen, Y Ritov, JA Wellner, *Efficient and adaptive estimation for semiparamet-*  
 565 *ric models*. (Springer) Vol. 4, (1993).
- 566 28. JT Runnenburg, Mean, median, mode. *Stat. Neerlandica* **32**, 73–79 (1978).
- 567 29. Wv Zwet, Mean, median, mode ii. *Stat. Neerlandica* **33**, 1–5 (1979).
- 568 30. H Rietz, On certain properties of frequency distributions of the powers and roots of the variates  
 569 of a given distribution. *Proc. Natl. Acad. Sci.* **13**, 817–820 (1927).
- 570 31. WR van Zwet, *Convex Transformations of Random Variables: Nebst Stellingen*. (1964).
- 571 32. K Pearson, X. contributions to the mathematical theory of evolution.—ii. skew variation in  
 572 homogeneous material. *Philos. Transactions Royal Soc. London.(A.)* **186**, 343–414 (1895).
- 573 33. AL Bowley, *Elements of statistics*. (King) No. 8, (1926).
- 574 34. RA Groeneveld, G Meeden, Measuring skewness and kurtosis. *J. Royal Stat. Soc. Ser. D*  
 575 *(The Stat.* **33**, 391–399 (1984).