

Semiparametric robust mean estimations based on the orderliness of quantile averages

Tuban Lee

This manuscript was compiled on May 24, 2023

1 As one of the most fundamental problems in statistics, robust loca-
2 tion estimation has many prominent solutions, such as the symmetric
3 trimmed mean, symmetric Winsorized mean, Hodges–Lehmann es-
4 timator, Huber M-estimator, and median of means. Recent studies
5 suggest that their maximum biases concerning the mean can be quite
6 different in asymmetric distributions, but the underlying mechanisms
7 and average performance remain largely unclear. In this article, simi-
8 lar to the mean-median-mode inequality, it is proven that within the
9 context of nearly all common unimodal distributions, there is an order-
10 liness of symmetric quantile averages with varying breakdown points.
11 Further deductions explain why the Winsorized mean and median of
12 means typically have smaller biases compared to the trimmed mean.
13 Building on the Hodges–Lehmann inequality, the superiority of the
14 median Hodges–Lehmann mean is discussed.

semiparametric | mean-median-mode inequality | asymptotic | unimodal
| Hodges–Lehmann estimator

1 **Data Availability.** Data for Figure ?? are given in SI Dataset
2 S1. All codes have been deposited in [GitHub](#).

3 **ACKNOWLEDGMENTS.** I sincerely acknowledge the insightful
4 comments from the editor which considerably elevated the lucidity
5 and merit of this paper.