

# Semiparametric robust mean estimations based on the orderliness of quantile averages

Tuban Lee

This manuscript was compiled on May 20, 2023

**As one of the most fundamental problems in statistics, robust location estimation has many prominent solutions, such as the symmetric trimmed mean, symmetric Winsorized mean, Hodges–Lehmann estimator, Huber  $M$ -estimator, and median of means. Recent studies suggest that their maximum biases concerning the mean can be quite different in asymmetric distributions, but the underlying mechanisms and average performance remain largely unclear. In this article, similar to the mean-median-mode inequality, it is proven that within the context of nearly all common unimodal distributions, there is an orderliness of symmetric quantile averages with varying breakdown points. Further deductions explain why the Winsorized mean and median of means typically have smaller biases compared to the trimmed mean. Building on the  $U$ -orderliness, the superiority of the median Hodges–Lehmann mean is discussed.**

semiparametric | mean-median-mode inequality | asymptotic | unimodal  
| Hodges–Lehmann estimator

In 1823, Gauss (1) proved that for any unimodal distribution with a finite second moment,  $|m - \mu| \leq \sqrt{\frac{3}{4}}\omega$ , where  $\mu$  is the population mean,  $m$  is the population median, and  $\omega$  is the root mean square deviation from the mode,  $M$ . This pioneering work revealed that despite potential bias with respect to the mean in robust estimates, the deviation remains bounded in unit of a scale parameter under certain assumptions. Bernard, Kazzi, and Vanduffel (2020) (2) further derived asymptotic bias bounds of any quantile for unimodal distributions with finite second moments by reducing this optimization problem to a parametric one, which can be solved analytically. They showed that the population median,  $m$ , has the smallest maximum distance to the population mean,  $\mu$ , among all symmetric quantile averages (SQA $_{\epsilon}$ ). Daniell, in 1920, (3) analyzed a class of estimators, linear combinations of order statistics, and identified that  $\epsilon$ -symmetric trimmed mean (STM $_{\epsilon}$ ) belongs to this class. Another popular choice, the  $\epsilon$ -symmetric Winsorized mean (SWM $_{\epsilon}$ ), named after Winsor and introduced by Tukey (4) and Dixon (5) in 1960, is also an  $L$ -estimator. Bieniek (2016) derived exact bias upper bounds of the Winsorized mean based on Danielak and Rychlik's work (2003) on the trimmed mean for any distribution with a finite second moment and confirmed that the former is smaller than the latter (6, 7). In 1963, Hodges and Lehmann (8) proposed a class of nonparametric location estimators based on rank tests and, from the Wilcoxon signed-rank statistic (9), deduced the median of pairwise means as a robust location estimator for a symmetric population. Both  $L$ -statistics and  $R$ -statistics achieve robustness essentially by removing a certain proportion of extreme values. In 1964, Huber (10) generalized maximum likelihood estimation to the minimization of the sum of a specific loss function, which measures the residuals between the data points and the model's parameters. Some  $L$ -estimators are also  $M$ -estimators, e.g., the sample mean is an  $M$ -estimator with a squared error loss function, while the sample median is

an  $M$ -estimator with an absolute error loss function (10). The Huber  $M$ -estimator is obtained by applying the Huber loss function that combines elements of both squared error and absolute error to achieve robustness against gross errors and high efficiency for contaminated Gaussian distributions (10). Sun, Zhou, and Fan (2020) examined the concentration bounds of Huber  $M$ -estimator (11). Mathieu (2022) (12) further derived the concentration bounds of  $M$ -estimators and demonstrated that, by selecting the tuning parameter which depends on the variance, Huber  $M$ -estimator can also be a sub-Gaussian estimator. The concept of median of means (MoM $_{k,b} = \frac{n}{k}$ ,  $k$  is the number of size in each block,  $b$  is the number of blocks) was implicitly introduced several times in Nemirovsky and Yudin (1983) (13), Jerrum, Valiant, and Vazirani (1986), (14) and Alon, Matias and Szegedy (1996) (15)'s works. Given its good performance even for distributions with infinite second moments, MoM has received increasing attention over the past decade (16–18). Devroye, Lerasle, Lugosi, and Oliveira (2016) showed that MoM nears the optimum of sub-Gaussian mean estimation with regards to concentration bounds when the distribution has a heavy tail (17). For a comparison of concentration bounds of trimmed mean, Huber  $M$ -estimator, median of means and other relevant estimators, readers are directed to Gobet, Lerasle, and Métivier's paper (2022) (19). Laforgue, Clemencon, and Bertail (2019) proposed the median of randomized means (MoRM $_{k,b}$ ) (18), wherein, rather than partitioning, an arbitrary number,  $b$ , of blocks are built independently from the sample, and showed that MoRM has a better non-asymptotic sub-Gaussian property compared to MoM. In fact, asymptotically, the Hodges–Lehmann (H-L) estimator is equivalent to MoM $_{k=2,b=\frac{n}{k}}$  and MoRM $_{k=2,b}$ , and they can be seen as the pairwise mean distribution is approximated by the sampling without replacement and bootstrap, respectively. For the asymptotic validity, readers are referred to the foundational works of Efron (1979) (20), Bickel and

## Significance Statement

In 1964, van Zwet introduced the convex transformation order for comparing the skewness of two distributions. This paradigm shift played a fundamental role in defining robust measures of distributions, from spread to kurtosis. Here, rather than examining the stochastic ordering between two distributions, the orderliness of quantile averages within a distribution is investigated. By classifying distributions through the signs of derivatives, a series of sophisticated robust mean estimators are deduced. Nearly all common nonparametric robust location estimators are found to be special cases thereof.

T.L. designed research, performed research, analyzed data, and wrote the paper.

The author declares no competing interest.

<sup>1</sup>To whom correspondence should be addressed. E-mail: tl@biomathematics.org

71 Freedman (1981, 1984) (21, 22), and Helmers, Janssen, and  
 72 Veraverbeke (1990) (23).

Here, the  $\epsilon, b$ -stratified mean is defined as

$$SM_{\epsilon, b, n} := \frac{b}{n} \left( \sum_{j=1}^{\lfloor \frac{b-1}{2b\epsilon} \rfloor} \sum_{i_j = \frac{(2bj-b-1)n\epsilon}{b-1}}^{\frac{(2bj-b+1)n\epsilon}{b-1}} X_{i_j} \right),$$

73 where  $X_1 \leq \dots \leq X_n$  denote the order statistics of a sample  
 74 of  $n$  independent and identically distributed random variables  
 75  $X_1, \dots, X_n$ .  $b \in \mathbb{N}$ ,  $b \geq 3$ . The definition was further refined to  
 76 guarantee the continuity of the breakdown point by incorporat-  
 77 ing an additional block in the center when  $\lfloor \frac{b-1}{2b\epsilon} \rfloor \bmod 2 = 0$ ,  
 78 or by adjusting the central block when  $\lfloor \frac{b-1}{2b\epsilon} \rfloor \bmod 2 = 1$  (SI  
 79 Text). If the subscript  $n$  is omitted, only the asymptotic  
 80 behavior is considered. If  $b$  is omitted,  $b = 3$  is assumed.  
 81  $SM_{\epsilon, b=3}$  is equivalent to  $STM_{\epsilon}$ , when  $\epsilon > \frac{1}{6}$ . The basic idea  
 82 of the stratified mean, when  $\frac{b-1}{2\epsilon} \in \mathbb{N}$ ,  $b \bmod 2 = 1$ , is to dis-  
 83 tribute the data into  $\frac{b-1}{2\epsilon}$  equal-sized non-overlapping blocks  
 84 according to their order, then further sequentially group these  
 85 blocks into  $b$  equal-sized strata and compute the mean of the  
 86 middle stratum, which is the median of means of each stratum.  
 87 In situations where  $i \bmod 1 \neq 0$ , a potential solution is to  
 88 generate multiple smaller samples that satisfy the equality by  
 89 sampling without replacement, and subsequently calculate the  
 90 mean of all estimations. The details of determining the sample  
 91 size and sampling times are provided in the SI Text. Although  
 92 the principle resembles that of the median of means, without  
 93 the random shift, the result is different from  $MoM_{k=\frac{n}{b}, b}$ .

94 **Data Availability.** Data for Figure ?? are given in SI Dataset  
 95 S1. All codes have been deposited in [GitHub](#).

96 **ACKNOWLEDGMENTS.** I sincerely acknowledge the insightful  
 97 comments from the editor which considerably elevated the lucidity  
 98 and merit of this paper.

- 99 1. CF Gauss, *Theoria combinationis observationum erroribus minimis obnoxiae*. (Henricus  
 100 Dieterich), (1823).
- 101 2. C Bernard, R Kazzi, S Vanduffel, Range value-at-risk bounds for unimodal distributions under  
 102 partial information. *Insur. Math. Econ.* **94**, 9–24 (2020).
- 103 3. P Daniell, Observations weighted according to order. *Am. J. Math.* **42**, 222–236 (1920).
- 104 4. JW Tukey, A survey of sampling from contaminated distributions in *Contributions to probability*  
 105 *and statistics*. (Stanford University Press), pp. 448–485 (1960).
- 106 5. WJ Dixon, Simplified Estimation from Censored Normal Samples. *The Annals Math. Stat.* **31**,  
 107 385 – 391 (1960).
- 108 6. M Bieniek, Comparison of the bias of trimmed and winsorized means. *Commun. Stat. Methods*  
 109 **45**, 6641–6650 (2016).
- 110 7. K Danielak, T Rychlik, Theory & methods: Exact bounds for the bias of trimmed means. *Aust.*  
 111 *& New Zealand J. Stat.* **45**, 83–96 (2003).
- 112 8. J Hodges Jr, E Lehmann, Estimates of location based on rank tests. *The Annals Math. Stat.*  
 113 **34**, 598–611 (1963).
- 114 9. F Wilcoxon, Individual comparisons by ranking methods. *Biom. Bull.* **1**, 80–83 (1945).
- 115 10. PJ Huber, Robust estimation of a location parameter. *Ann. Math. Stat.* **35**, 73–101 (1964).
- 116 11. Q Sun, WX Zhou, J Fan, Adaptive huber regression. *J. Am. Stat. Assoc.* **115**, 254–265 (2020).
- 117 12. T Mathieu, Concentration study of  $m$ -estimators using the influence function. *Electron. J. Stat.*  
 118 **16**, 3695–3750 (2022).
- 119 13. AS Nemirovskij, DB Yudin, *Problem complexity and method efficiency in optimization*. (Wiley-  
 120 Interscience), (1983).
- 121 14. MR Jerrum, LG Valiant, VV Vazirani, Random generation of combinatorial structures from a  
 122 uniform distribution. *Theor. computer science* **43**, 169–188 (1986).
- 123 15. N Alon, Y Matias, M Szegedy, The space complexity of approximating the frequency moments  
 124 in *Proceedings of the twenty-eighth annual ACM symposium on Theory of computing*. pp.  
 125 20–29 (1996).
- 126 16. PL Bühlmann, Bagging, subbagging and bragging for improving some prediction algorithms  
 127 in *Research report/Seminar für Statistik, Eidgenössische Technische Hochschule (ETH)*.  
 128 (Seminar für Statistik, Eidgenössische Technische Hochschule (ETH), Zürich), Vol. 113,  
 129 (2003).
- 130 17. L Devroye, M Lerasle, G Lugosi, RI Oliveira, Sub-gaussian mean estimators. *The Annals Stat.*  
 131 **44**, 2695–2725 (2016).
- 132 18. P Laforgue, S Cléménçon, P Bertail, On medians of (randomized) pairwise means in *Interna-*  
 133 *tional Conference on Machine Learning*. (PMLR), pp. 1272–1281 (2019).

- 134 19. E Gobet, M Lerasle, D Métivier, Mean estimation for Randomized Quasi Monte Carlo method. 134  
 working paper or preprint (2022). 135
20. B Efron, Bootstrap methods: Another look at the jackknife. *The Annals Stat.* **7**, 1–26 (1979). 136
21. PJ Bickel, DA Freedman, Some asymptotic theory for the bootstrap. *The annals statistics* **9**, 137  
 1196–1217 (1981). 138
22. PJ Bickel, DA Freedman, Asymptotic normality and the bootstrap in stratified sampling. *The*  
*annals statistics* **12**, 470–482 (1984). 139
23. R Helmers, P Janssen, N Veraverbeke, *Bootstrapping U-quantiles*. (CWI. Department of 140  
 Operations Research, Statistics, and System Theory [BS]), (1990). 141  
 142