

# Semiparametric robust mean estimations based on the orderliness of quantile averages

Tuban Lee

This manuscript was compiled on June 9, 2023

semiparametric | mean-median-mode inequality | asymptotic | unimodal  
| Hodges–Lehmann estimator

## Hodges–Lehmann inequality and $\gamma$ - $U$ -orderliness

The Hodges–Lehmann estimator stands out as a unique robust location estimator due to its definition being substantially dissimilar from conventional  $L$ -estimators,  $R$ -estimators, and  $M$ -estimators. In their landmark paper, *Estimates of location based on rank tests*, Hodges and Lehmann (1) proposed two methods for computing the H-L estimator: the Wilcoxon score  $R$ -estimator and the median of pairwise means. The Wilcoxon score  $R$ -estimator is a location estimator based on signed-rank test, or  $R$ -estimator, (1) and was later independently discovered by Sen (1963) (2, 3). However, the median of pairwise means is a generalized  $L$ -statistic and a trimmed  $U$ -statistic, as classified by Serfling in his novel conceptualized study in 1984 (4). Serfling further advanced the understanding by generalizing the H-L kernel function as  $hl_k(x_1, \dots, x_k) = \frac{1}{k} \sum_{i=1}^k x_i$ , where  $k \in \mathbb{N}$  (4). Here, the weighted H-L kernel function is defined as  $whl_k(x_1, \dots, x_k) = \frac{\sum_{i=1}^k x_i w_i}{\sum_{i=1}^k w_i}$ , where  $w_i$ s are the weights applied to each element.

**Data Availability.** Data for Figure ?? are given in SI Dataset S1. All codes have been deposited in [GitHub](#).

**ACKNOWLEDGMENTS.** I sincerely acknowledge the insightful comments from the editor which considerably elevated the lucidity and merit of this paper.

1. J Hodges Jr, E Lehmann, Estimates of location based on rank tests. *The Annals Math. Stat.* **34**, 598–611 (1963).
2. PK Sen, On the estimation of relative potency in dilution (-direct) assays by distribution-free methods. *Biometrics* pp. 532–552 (1963).
3. M Ghosh, MJ Schell, PK Sen, A conversation with pranab kumar sen. *Stat. Sci.* pp. 548–564 (2008).
4. RJ Serfling, Generalized  $L$ -,  $m$ -, and  $r$ -statistics. *The Annals Stat.* **12**, 76–86 (1984).