

# Semiparametric robust mean estimations based on the orderliness of quantile averages

Tuban Lee

This manuscript was compiled on July 9, 2023

semiparametric | mean-median-mode inequality | asymptotic | unimodal  
| Hodges–Lehmann estimator

## Weighted Inequalities and Binomial Mean

Analogous to the  $\gamma$ -orderliness, the  $\gamma$ -trimming inequality for a right-skewed distribution is defined as  $\forall 0 \leq \epsilon_1 \leq \epsilon_2 \leq \frac{1}{1+\gamma}$ ,  $\text{TM}_{\epsilon_1, \gamma} \geq \text{TM}_{\epsilon_2, \gamma}$ .  $\gamma$ -orderliness is a sufficient condition for the  $\gamma$ -trimming inequality, as proven in the SI Text. The next theorem shows a relation between the  $\epsilon, \gamma$ -quantile average and the  $\epsilon, \gamma$ -trimmed mean under the  $\gamma$ -trimming inequality, suggesting the  $\gamma$ -orderliness is not a necessary condition for the  $\gamma$ -trimming inequality.

**Theorem .1.** *For a distribution that is right-skewed and follows the  $\gamma$ -trimming inequality, it is asymptotically true that the quantile average is always greater or equal to the corresponding trimmed mean with the same  $\epsilon$  and  $\gamma$ , for all  $0 \leq \epsilon \leq \frac{1}{1+\gamma}$ .*

*Proof.* According to the definition of the  $\gamma$ -trimming inequality:  $\forall 0 \leq \epsilon \leq \frac{1}{1+\gamma}$ ,  $\frac{1}{1-\epsilon-\gamma\epsilon} \int_{\gamma\epsilon-\delta}^{1-\epsilon+\delta} Q(u) du \geq \frac{1}{1-\epsilon-\gamma\epsilon} \int_{\gamma\epsilon}^{1-\epsilon} Q(u) du$ , where  $\delta$  is an infinitesimal positive quantity. Subsequently, rewriting the inequality gives  $\int_{\gamma\epsilon-\delta}^{1-\epsilon+\delta} Q(u) du - \frac{1-\epsilon-\gamma\epsilon+2\delta}{1-\epsilon-\gamma\epsilon} \int_{\gamma\epsilon}^{1-\epsilon} Q(u) du \geq 0 \Leftrightarrow \int_{1-\epsilon}^{1-\epsilon+\delta} Q(u) du + \int_{\gamma\epsilon-\delta}^{\gamma\epsilon} Q(u) du - \frac{2\delta}{1-\epsilon-\gamma\epsilon} \int_{\gamma\epsilon}^{1-\epsilon} Q(u) du \geq 0$ . Since  $\delta \rightarrow 0^+$ ,  $\frac{1}{2\delta} \left( \int_{1-\epsilon}^{1-\epsilon+\delta} Q(u) du + \int_{\gamma\epsilon-\delta}^{\gamma\epsilon} Q(u) du \right) = \frac{Q(\gamma\epsilon)+Q(1-\epsilon)}{2} \geq \frac{1}{1-\epsilon-\gamma\epsilon} \int_{\gamma\epsilon}^{1-\epsilon} Q(u) du$ , the proof is complete.  $\square$

An analogous result about the relation between the  $\epsilon, \gamma$ -trimmed mean and the  $\epsilon, \gamma$ -Winsorized mean can be obtained in the following theorem.

**Theorem .2.** *For a right-skewed distribution following the  $\gamma$ -trimming inequality, asymptotically, the Winsorized mean is always greater or equal to the corresponding trimmed mean with the same  $\epsilon$  and  $\gamma$ , for all  $0 \leq \epsilon \leq \frac{1}{1+\gamma}$ , provided that  $0 \leq \gamma \leq 1$ . If assuming  $\gamma$ -orderliness, the inequality is valid for any non-negative  $\gamma$ .*

*Proof.* According to Theorem .1,  $\frac{Q(\gamma\epsilon)+Q(1-\epsilon)}{2} \geq \frac{1}{1-\epsilon-\gamma\epsilon} \int_{\gamma\epsilon}^{1-\epsilon} Q(u) du \Leftrightarrow \gamma\epsilon(Q(\gamma\epsilon)+Q(1-\epsilon)) \geq \left( \frac{2\gamma\epsilon}{1-\epsilon-\gamma\epsilon} \right) \int_{\gamma\epsilon}^{1-\epsilon} Q(u) du$ . Then, if  $0 \leq \gamma \leq 1$ ,  $1, \left(1 - \frac{1}{1-\epsilon-\gamma\epsilon}\right) \int_{\gamma\epsilon}^{1-\epsilon} Q(u) du + \gamma\epsilon(Q(\gamma\epsilon)+Q(1-\epsilon)) \geq 0 \Rightarrow \int_{\gamma\epsilon}^{1-\epsilon} Q(u) du + \gamma\epsilon Q(\gamma\epsilon) + \epsilon Q(1-\epsilon) \geq \int_{\gamma\epsilon}^{1-\epsilon} Q(u) du + \gamma\epsilon(Q(\gamma\epsilon)+Q(1-\epsilon)) \geq \frac{1}{1-\epsilon-\gamma\epsilon} \int_{\gamma\epsilon}^{1-\epsilon} Q(u) du$ , the proof of the first assertion is complete. The second assertion is established in Theorem 0.3. in the SI Text.  $\square$

Replacing the TM in the  $\gamma$ -trimming inequality with WA forms the definition of the  $\gamma$ -weighted inequality. The  $\gamma$ -orderliness also implies the  $\gamma$ -Winsorization inequality when  $0 \leq \gamma \leq 1$ , as proven in the SI Text. The same rationale as presented in Theorem ??, for a location-scale distribution characterized by a location parameter  $\mu$  and a scale parameter  $\lambda$ , asymptotically, any  $\text{WA}(\epsilon, \gamma)$  can be expressed as  $\lambda \text{WA}_0(\epsilon, \gamma) + \mu$ , where  $\text{WA}_0(\epsilon, \gamma)$  is an function of  $Q_0(p)$  according to the definition of the weighted average. Adhering to the rationale present in Theorem ??, for any probability distribution within a location-scale family, a necessary and sufficient condition for whether it follows the  $\gamma$ -weighted inequality is whether the family of probability distributions also adheres to the  $\gamma$ -weighted inequality.

## Hodges–Lehmann inequality and $\gamma$ -U-orderliness

The Hodges–Lehmann estimator stands out as a unique robust location estimator due to its definition being substantially dissimilar from conventional  $L$ -estimators,  $R$ -estimators, and  $M$ -estimators. In their landmark paper, *Estimates of location based on rank tests*, Hodges and Lehmann (1) proposed two methods for computing the H-L estimator: the Wilcoxon score  $R$ -estimator and the median of pairwise means. The Wilcoxon score  $R$ -estimator is a location estimator based on signed-rank test, or  $R$ -estimator, (1) and was later independently discovered by Sen (1963) (2, 3). However, the median of pairwise means is a generalized  $L$ -statistic and a trimmed  $U$ -statistic, as classified by Serfling in his novel conceptualized study in 1984 (4). Serfling further advanced the understanding by generalizing the H-L kernel as  $hl_k(x_1, \dots, x_k) = \frac{1}{k} \sum_{i=1}^k x_i$ , where  $k \in \mathbb{N}$  (4). Here, the weighted H-L kernel is defined as  $whl_k(x_1, \dots, x_k) = \frac{\sum_{i=1}^k x_i w_i}{\sum_{i=1}^k w_i}$ , where  $w_i$ s are the weights applied to each element.

By using the weighted H-L kernel and the  $L$ -estimator, it is now clear that the Hodges–Lehmann estimator is an  $LL$ -statistic, the definition of which is provided as follows:

$$LL_{k, \epsilon, \gamma, n} := L_{\epsilon_0, \gamma, n} \left( \text{sort} \left( (whl_k(X_{N_1}, \dots, X_{N_k}))_{N=1}^{(n)} \right) \right),$$

where  $L_{\epsilon_0, \gamma, n}(Y)$  represents the  $\epsilon_0, \gamma$ - $L$ -estimator that uses the sorted sequence,  $\text{sort} \left( (whl_k(X_{N_1}, \dots, X_{N_k}))_{N=1}^{(n)} \right)$ , as input. The upper asymptotic breakdown point of  $LL_{k, \epsilon, \gamma}$  is  $\epsilon = 1 - (1 - \epsilon_0)^{\frac{1}{k}}$ , as proven in DSSM II. There are two ways to adjust the breakdown point: either by setting  $k$  as a constant and adjusting  $\epsilon_0$ , or by setting  $\epsilon_0$  as a constant and adjusting  $k$ . In the above definition,  $k$  is discrete, but the bootstrap

T.L. designed research, performed research, analyzed data, and wrote the paper.

The author declares no competing interest.

<sup>1</sup>To whom correspondence should be addressed. E-mail: tl@biomathematics.org

method can be applied to ensure the continuity of  $k$ , also making the breakdown point continuous. Specifically, if  $k \in \mathbb{R}$ , let the bootstrap size be denoted by  $b$ , then first sampling the original sample  $(1 - k + \lfloor k \rfloor)b$  times with each sample size of  $\lfloor k \rfloor$ , and then subsequently sampling  $(1 - \lfloor k \rfloor + k)b$  times with each sample size of  $\lfloor k \rfloor$ ,  $(1 - k + \lfloor k \rfloor)b \in \mathbb{N}$ ,  $(1 - \lfloor k \rfloor + k)b \in \mathbb{N}$ . The corresponding kernels are computed separately, and the pooled sorted sequence is used as the input for the  $L$ -estimator. Let  $\mathbf{S}_k$  represent the sorted sequence. Indeed, for any finite sample,  $X$ , when  $k = n$ ,  $\mathbf{S}_k$  becomes a single point,  $whl_{k=n}(X_1, \dots, X_n)$ . When  $\mathbf{w}_i = 1$ , the minimum of  $\mathbf{S}_k$  is  $\frac{1}{k} \sum_{i=1}^k X_i$ , due to the property of order statistics. The maximum of  $\mathbf{S}_k$  is  $\frac{1}{k} \sum_{i=1}^k X_{n-i+1}$ . The monotonicity of the order statistics implies the monotonicity of the extrema with respect to  $k$ , i.e., the support of  $\mathbf{S}_k$  shrinks monotonically. For unequal  $\mathbf{w}_i$ s, the shrinkage of the support of  $\mathbf{S}_k$  might not be strictly monotonic, but the general trend remains, since all  $LL$ -statistics converge to the same point, as  $k \rightarrow n$ . Therefore, if  $\sum_{i=1}^n \frac{X_i \mathbf{w}_i}{\mathbf{w}_i}$  approaches the population mean when  $n \rightarrow \infty$ , all  $LL$ -statistics based on such consistent kernel function approach the population mean as  $k \rightarrow \infty$ . For example, if  $whl_k = \text{BM}_{\nu, \epsilon_k, n=k}$ ,  $\nu \ll \epsilon_k^{-1}$ ,  $\epsilon_k \rightarrow 0$ , such kernel function is consistent. These cases are termed the  $LL$ -mean ( $\text{LLM}_{k, \epsilon, \gamma, n}$ ). By substituting the  $\text{WA}_{\epsilon_0, \gamma, n}$  for the  $L_{\epsilon_0, \gamma, n}$  in  $LL$ -statistic, the resulting statistic is referred to as the weighted  $L$ -statistic ( $\text{WLM}_{k, \epsilon, \gamma, n}$ ). The case having a consistent kernel function is termed as the weighted  $L$ -mean ( $\text{WLM}_{k, \epsilon, \gamma, n}$ ). The  $w_i = 1$  case of  $\text{WLM}_{k, \epsilon, \gamma, n}$  is termed the weighted Hodges-Lehmann mean ( $\text{WHLM}_{k, \epsilon, \gamma, n}$ ). The  $\text{WHLM}_{k=1, \epsilon, \gamma, n}$  is the weighted average. If  $k \geq 2$  and the  $\text{WA}$  in  $\text{WHLM}$  is set as  $\text{TM}_{\epsilon_0}$ , it is called the trimmed H-L mean (Figure ??,  $k = 2$ ,  $\epsilon_0 = \frac{15}{64}$ ). The  $\text{THLM}_{k=2, \epsilon, \gamma=1, n}$  appears similar to the Wilcoxon's one-sample statistic investigated by Saleh in 1976 (5), which involves first censoring the sample, and then computing the mean of the number of events that the pairwise mean is greater than zero. The  $\text{THLM}_{k=2, \epsilon=1-(1-\frac{1}{2})^{\frac{1}{2}}, \gamma=1, n}$  is the Hodges-Lehmann estimator, or more generally, a special case of the median Hodges-Lehmann mean ( $m\text{HLM}_{k, n}$ ).  $m\text{HLM}_{k, n}$  is asymptotically equivalent to the  $\text{MoM}_{k, b=\frac{n}{k}}$  as discussed previously. Therefore, it is possible to define a series of location estimators, analogous to the  $\text{WHLM}$ , based on  $\text{MoM}$ . For example, the  $\gamma$ -median of means,  $\gamma\text{moM}_{k, b=\frac{n}{k}, n}$ , is defined by replacing the median in  $\text{MoM}_{k, b=\frac{n}{k}, n}$  with the  $\gamma$ -median.

The  $hl_k$  kernel distribution, denoted as  $F_{hl_k}$ , can be defined as the probability distribution of the sorted sequence  $\text{sort}\left((hl_k(X_{N_1}, \dots, X_{N_k}))_{N=1}^n\right)$ . For any real value  $y$ , the cdf of the  $hl_k$  kernel distribution is given by:  $F_{hk}(y) = \Pr(Y_i \leq y)$ , where  $Y_i$  represents an individual element from the sorted sequence. The overall  $hl_k$  kernel distributions possess a two-dimensional structure, encompassing  $n$  kernel distributions with varying  $k$  values, from 1 to  $n$ , where one dimension is inherent to each individual kernel distribution, while the other is formed by the alignment of the same percentiles across all kernel distributions. As  $k$  increases, all percentiles converge to  $\bar{X}$ , leading to the concept of  $\gamma$ - $U$ -orderliness:

$$(\forall k_2 \geq k_1 \geq 1, \gamma\text{mHLM}_{k_2, \epsilon=1-(\frac{\gamma}{1+\gamma})^{\frac{1}{k_2}}, \gamma} \geq \gamma\text{mHLM}_{k_1, \epsilon=1-(\frac{\gamma}{1+\gamma})^{\frac{1}{k_1}}, \gamma})$$

$$(\forall k_2 \geq k_1 \geq 1, \gamma\text{mHLM}_{k_2, \epsilon=1-(\frac{\gamma}{1+\gamma})^{\frac{1}{k_2}}, \gamma} \leq \gamma\text{mHLM}_{k_1, \epsilon=1-(\frac{\gamma}{1+\gamma})^{\frac{1}{k_1}}, \gamma})$$

where  $\gamma\text{mHLM}_k$  sets the  $\text{WA}$  in  $\text{WHLM}$  as  $\gamma$ -median, with  $\gamma$  being constant. The direction of the inequality depends on the relative magnitudes of  $\gamma\text{mHLM}_{k=1, \epsilon, \gamma} = \gamma m$  and  $\gamma\text{mHLM}_{k=\infty, \epsilon, \gamma} = \mu$ . The Hodges-Lehmann inequality can be defined as a special case of the  $\gamma$ - $U$ -orderliness when  $\gamma = 1$ . When  $\gamma \in \{0, \infty\}$ , the  $\gamma$ - $U$ -orderliness is valid for any distribution as previously shown. If  $\gamma \notin \{0, \infty\}$ , analytically proving the validity of the  $\gamma$ - $U$ -orderliness for a parametric distribution is pretty challenging. As an example, the  $hl_2$  kernel distribution has a probability density function  $f_{hl_2}(x) = \int_0^{2x} 2f(t)f(2x-t)dt$  (a result after the transformation of variables); the support of the original distribution is assumed to be  $[0, \infty)$  for simplicity. The expected value of the H-L estimator is the positive solution of  $\int_0^{\text{H-L}} (f_{hl_2}(s))ds = \frac{1}{2}$ . For the exponential distribution,  $f_{hl_2, \text{exp}}(x) = 4\lambda^{-2}xe^{-2\lambda^{-1}x}$ ,  $\lambda$  is a scale parameter,  $E[\text{H-L}] = \frac{-W_{-1}(-\frac{1}{2e})-1}{2}\lambda \approx 0.839\lambda$ , where  $W_{-1}$  is a branch of the Lambert  $W$  function which cannot be expressed in terms of elementary functions. However, the violation of the  $\gamma$ - $U$ -orderliness is bounded under certain assumptions, as shown below.

**Theorem .3.** For any distribution with a finite second central moment,  $\sigma^2$ , the following concentration bound can be established for the  $\gamma$ -median of means,

$$\mathbb{P}\left(\gamma\text{moM}_{k, b=\frac{n}{k}, n} - \mu > \frac{t\sigma}{\sqrt{k}}\right) \leq e^{-\frac{2n}{k}\left(\frac{1}{1+\gamma} - \frac{1}{k+t^2}\right)^2}.$$

*Proof.* Denote the mean of each block as  $\hat{\mu}_i$ ,  $1 \leq i \leq b$ . Observe that the event  $\left\{\gamma\text{moM}_{k, b=\frac{n}{k}, n} - \mu > \frac{t\sigma}{\sqrt{k}}\right\}$  necessitates the condition that there are at least  $b(1 - \frac{\gamma}{1+\gamma})$  of  $\hat{\mu}_i$ s larger than  $\mu$  by more than  $\frac{t\sigma}{\sqrt{k}}$ , i.e.,  $\left\{\gamma\text{moM}_{k, b=\frac{n}{k}, n} - \mu > \frac{t\sigma}{\sqrt{k}}\right\} \subset \left\{\sum_{i=1}^b \mathbf{1}_{(\hat{\mu}_i - \mu) > \frac{t\sigma}{\sqrt{k}}} \geq b\left(1 - \frac{\gamma}{1+\gamma}\right)\right\}$ , where  $\mathbf{1}_A$  is the indicator of event  $A$ . Assuming a finite second central moment,  $\sigma^2$ , it follows from one-sided Chebeshev's inequality that  $\mathbb{E}\left(\mathbf{1}_{(\hat{\mu}_i - \mu) > \frac{t\sigma}{\sqrt{k}}}\right) = \mathbb{P}\left((\hat{\mu}_i - \mu) > \frac{t\sigma}{\sqrt{k}}\right) \leq \frac{\sigma^2}{k\sigma^2 + t^2\sigma^2}$ .

Given that  $\mathbf{1}_{(\hat{\mu}_i - \mu) > \frac{t\sigma}{\sqrt{k}}} \in [0, 1]$  are independent and identically distributed random variables, according to the aforementioned inclusion relation, the one-sided Chebeshev's inequality and the one-sided Hoeffding's inequality,  $\mathbb{P}\left(\gamma\text{moM}_{k, b=\frac{n}{k}, n} - \mu > \frac{t\sigma}{\sqrt{k}}\right) \leq \mathbb{P}\left(\sum_{i=1}^b \mathbf{1}_{(\hat{\mu}_i - \mu) > \frac{t\sigma}{\sqrt{k}}} \geq b\left(1 - \frac{\gamma}{1+\gamma}\right)\right) = \mathbb{P}\left(\frac{1}{b} \sum_{i=1}^b \left(\mathbf{1}_{(\hat{\mu}_i - \mu) > \frac{t\sigma}{\sqrt{k}}} - \mathbb{E}\left(\mathbf{1}_{(\hat{\mu}_i - \mu) > \frac{t\sigma}{\sqrt{k}}}\right)\right) \geq \left(1 - \frac{\gamma}{1+\gamma}\right) - \mathbb{E}\left(\mathbf{1}_{(\hat{\mu}_i - \mu) > \frac{t\sigma}{\sqrt{k}}}\right)\right) \leq e^{-2b\left(\left(1 - \frac{\gamma}{1+\gamma}\right) - \mathbb{E}\left(\mathbf{1}_{(\hat{\mu}_i - \mu) > \frac{t\sigma}{\sqrt{k}}}\right)\right)^2} = e^{-2b\left(1 - \frac{\gamma}{1+\gamma} - \frac{\sigma^2}{k\sigma^2 + t^2\sigma^2}\right)^2} = e^{-2b\left(\frac{1}{1+\gamma} - \frac{1}{k+t^2}\right)^2}.$   $\square$

**Theorem .4.** Let  $B(k, \gamma, t, n) = e^{-\frac{2n}{k}\left(\frac{1}{1+\gamma} - \frac{1}{k+t^2}\right)^2}$ . If  $k \in \mathbb{N}$ ,  $\gamma \geq 0$ ,  $0 \leq t^2 < \gamma + 1$ , and  $\gamma - t^2 + 1 \leq k \leq \frac{1}{2}\sqrt{9\gamma^2 + 18\gamma - 8\gamma t^2 - 8t^2 + 9} + \frac{1}{2}(3\gamma - 2t^2 + 3)$ ,  $B$  is monotonic decreasing with respect to  $k$ .

165 *Proof.* Since  $\frac{\partial B}{\partial k} = \left( \frac{2n\left(\frac{1}{\gamma+1} - \frac{1}{k+t^2}\right)^2}{k^2} - \frac{4n\left(\frac{1}{\gamma+1} - \frac{1}{k+t^2}\right)}{k(k+t^2)^2} \right)$   
166  $e^{-\frac{2n\left(\frac{1}{\gamma+1} - \frac{1}{k+t^2}\right)^2}{k^2}}$  and  $n \in \mathbb{N}$ ,  $\frac{\partial B}{\partial k} \leq 0 \Leftrightarrow$   
167  $\frac{2n\left(\frac{1}{\gamma+1} - \frac{1}{k+t^2}\right)^2}{k^2} - \frac{4n\left(\frac{1}{\gamma+1} - \frac{1}{k+t^2}\right)}{k(k+t^2)^2} \leq 0 \Leftrightarrow$   
168  $\frac{2n(-\gamma+k+t^2-1)(k^2-3(\gamma+1)k+2kt^2+t^2(-\gamma+t^2-1))}{(\gamma+1)^2 k^2 (k+t^2)^3} \leq 0 \Leftrightarrow$   
169  $(-\gamma+k+t^2-1)(k^2-3(\gamma+1)k+2kt^2+t^2(-\gamma+t^2-1))$   
170  $\leq 0$ . When the factors are expanded, it yields a cubic inequality in terms of  $k$ :  $k^3+k^2(3t^2-4(\gamma+1))+3k(\gamma-t^2+1)^2+t^2(\gamma-t^2+1)^2 \leq 0$ . Assuming  $0 \leq t^2 < \gamma+1$  and  $\gamma \geq 0$ ,  
171 using the factored form and subsequently applying the quadratic formula, the inequality is valid if  $\gamma-t^2+1 \leq k \leq$   
172  $\frac{1}{2}\sqrt{9\gamma^2+18\gamma-8\gamma t^2-8t^2+9} + \frac{1}{2}(3\gamma-2t^2+3)$ .  $\square$

176 Let  $X$  be a random variable and  $\bar{Y} = \frac{1}{k}(Y_1 + \dots + Y_k)$  be  
177 the average of  $k$  independent, identically distributed copies  
178 of  $X$ . Applying the variance operation gives:  $\text{Var}(\bar{Y}) =$   
179  $\text{Var}\left(\frac{1}{k}(Y_1 + \dots + Y_k)\right) = \frac{1}{k^2}(\text{Var}(Y_1) + \dots + \text{Var}(Y_k)) =$   
180  $\frac{1}{k^2}(k\sigma^2) = \frac{\sigma^2}{k}$ , since the variance operation is a linear op-  
181 erator for independent variables, and the variance of a scaled  
182 random variable is the square of the scale times the vari-  
183 ance of the variable, i.e.,  $\text{Var}(cX) = E[(cX - E[cX])^2] =$   
184  $E[(cX - cE[X])^2] = E[c^2(X - E[X])^2] = c^2 E[(X - E[X])^2] =$   
185  $c^2 \text{Var}(X)$ . Thus, the standard deviation of the  $hl_k$  kernel  
186 distribution, asymptotically, is  $\frac{\sigma}{\sqrt{k}}$ . By utilizing the asymp-  
187 totic bias bound of any quantile for any continuous distribu-  
188 tion with a finite second central moment,  $\sigma^2$ , (6), a conserva-  
189 tive asymptotic bias bound of  $\gamma \text{moM}_{k,b=\frac{n}{k}}$  can be estab-  
190 lished as  $\gamma \text{moM}_{k,b=\frac{n}{k}} - \mu \leq \sqrt{\frac{\gamma}{1-\gamma}} \sigma_{hl_k} = \sqrt{\frac{\gamma}{k}} \sigma$ . That  
191 implies in Theorem .3,  $t < \sqrt{\gamma}$ , so when  $\gamma = 1$ , the upper  
192 bound of  $k$ , subject to the monotonic decreasing constraint,  
193 is  $2 + \sqrt{5} < \frac{1}{2}\sqrt{9+18-8t^2-8t^2+9} + \frac{1}{2}(3-2t^2+3) \leq 6$ ,  
194 the lower bound is  $1 < 2 - t^2 \leq 2$ . These analyses elucidate a  
195 surprising result: although the conservative asymptotic bound  
196 of  $\text{MoM}_{k,b=\frac{n}{k}}$  is monotonic with respect to  $k$ , its concentration  
197 bound is optimal when  $k \in (2 + \sqrt{5}, 6]$ .

198 Then consider the structure within each individual  $hl_k$  ker-  
199 nel distribution. The sorted sequence  $\mathbf{S}_k$ , when  $k = n - 1$ ,  
200 has  $n$  elements and the corresponding  $hl_k$  kernel distribu-  
201 tion can be seen as a location-scale transformation of the  
202 original distribution, so the corresponding  $hl_k$  kernel distri-  
203 bution is  $\nu$ th  $\gamma$ -ordered if and only if the original distri-  
204 bution is  $\nu$ th  $\gamma$ -ordered according to Theorem ???. Ana-  
205 lytically proving other cases is challenging. For example,  
206  $f'_{hl_2}(x) = 4f(2x)f(0) + \int_0^{2x} 4f(t)f'(2x-t)dt$ , the strict nega-  
207 tive of  $f'_{hl_2}(x)$  is not guaranteed if just assuming  $f'(x) < 0$ ,  
208 so, even if the original distribution is monotonic decreasing,  
209 the  $hl_2$  kernel distribution might be non-monotonic. Also,  
210 unlike the pairwise difference distribution, if the original dis-  
211 tribution is unimodal, the pairwise mean distribution might  
212 be non-unimodal, as demonstrated by a counterexample given  
213 by Chung in 1953 and mentioned by Hodges and Lehmann  
214 in 1954 (7, 8). Theorem ??? implies that the violation of  
215  $\nu$ th  $\gamma$ -orderliness within the  $hl_k$  kernel distribution is also  
216 bounded, and the bound monotonically shrinks as  $k$  increases  
217 because the bound is in unit of the standard deviation of the

$hl_k$  kernel distribution. If all  $hl_k$  kernel distributions are  $\nu$ th  
 $\gamma$ -ordered and the distribution itself is  $\nu$ th  $\gamma$ -ordered and  $\gamma$ - $U$ -  
ordered, then the distribution is called  $\nu$ th  $\gamma$ - $U$ -ordered. The  
following theorems highlight the significance of  $\gamma$ -symmetric  
distributions.

**Theorem .5.** Any  $\gamma$ -symmetric distribution is  $\nu$ th  $\gamma$ - $U$ -  
ordered, provided that the  $\gamma$  is the same.

The succeeding theorem shows that the  $whl_k$  kernel distri-  
bution is invariably a location-scale distribution if the original  
distribution belongs to a location-scale family with the same  
location and scale parameters.

**Theorem .6.**  $whl_k(x_1 = \lambda x_1 + \mu, \dots, x_k = \lambda x_k + \mu) =$   
 $\lambda whl_k(x_1, \dots, x_k) + \mu$ .

*Proof.*  $whl_k(x_1 = \lambda x_1 + \mu, \dots, x_k = \lambda x_k + \mu) =$   
 $\frac{\sum_{i=1}^k (\lambda x_i + \mu) w_i}{\sum_{i=1}^k w_i} = \frac{\sum_{i=1}^k \lambda x_i w_i + \sum_{i=1}^k \mu w_i}{\sum_{i=1}^k w_i} = \lambda \frac{\sum_{i=1}^k x_i w_i}{\sum_{i=1}^k w_i} +$   
 $\frac{\sum_{i=1}^k \mu w_i}{\sum_{i=1}^k w_i} = \lambda \frac{\sum_{i=1}^k x_i w_i}{\sum_{i=1}^k w_i} + \mu = \lambda whl_k(x_1, \dots, x_k) + \mu$ .  $\square$

According to Theorem .6, the  $\gamma$ -weighted inequality for a  
right-skewed distribution can be modified as  $\forall 0 \leq \epsilon_{01} \leq \epsilon_{02} \leq$   
 $\frac{1}{1+\gamma}$ ,  $\text{WLM}_{k,\epsilon=1-(1-\epsilon_{01})^{\frac{1}{k}},\gamma} \geq \text{WLM}_{k,\epsilon=1-(1-\epsilon_{02})^{\frac{1}{k}},\gamma}$ , which  
holds the same rationale as the  $\gamma$ -weighted inequality defined  
in the last section. If the  $\nu$ th  $\gamma$ -orderliness is valid for the  
 $whl_k$  kernel distribution, then all results in the last section can  
be directly implemented. From that, the binomial H-L mean  
(set the WA as BM) can be constructed (Figure ??), while its  
maximum breakdown point is  $\approx 0.065$  if  $\nu = 3$ . A compar-  
ison of the biases of  $\text{BM}_{\nu=3,\epsilon=\frac{1}{8}}$ ,  $\text{SQM}_{\epsilon=\frac{1}{8}}$ ,  $\text{THLM}_{k=2,\epsilon=\frac{1}{8}}$ ,  
 $\text{WHLM}_{k=2,\epsilon=\frac{1}{8}}$ ,  $\text{MHLM}_{k=\frac{2\ln(2)-\ln(3)}{3\ln(2)-\ln(7)},\epsilon=\frac{1}{8}}$  (midhinge  
H-L mean),  $\text{mHLM}_{k=\frac{\ln(2)}{3\ln(2)-\ln(7)},\epsilon=\frac{1}{8}}$ ,  $\text{THLM}_{k=5,\epsilon=\frac{1}{8}}$ ,  
and  $\text{WHLM}_{k=5,\epsilon=\frac{1}{8}}$  is appropriate (Figure ??, SI  
Dataset S1), given their same breakdown points, with  
 $\text{mHLM}_{k=\frac{\ln(2)}{3\ln(2)-\ln(7)},\epsilon=\frac{1}{8}}$  exhibiting the smallest biases.  
Another comparison among the H-L estimator, the trimmed  
mean, and the Winsorized mean, all with the same breakdown  
point, yields the same result that the H-L estimator has the  
smallest biases (SI Dataset S1). This aligns with Devroye et  
al.(2016)'s seminal work that MoM is nearly optimal with  
regards to concentration bounds for heavy-tailed distributions  
(9).

In 1958, Richtmyer introduced the concept of quasi-Monte  
Carlo simulation that utilizes low-discrepancy sequences, re-  
sulting in a significant reduction in computational expenses for  
large sample simulation (10). Among various low-discrepancy  
sequences, Sobol sequences are often favored in quasi-Monte  
Carlo methods (11). Building upon this principle, in 1991,  
Do and Hall extended it to bootstrap and found that the  
quasi-random approach resulted in lower variance compared  
to other bootstrap Monte Carlo procedures (12). By using  
a deterministic approach, the variance of  $\text{mHLM}_{k,n}$  is much  
lower than that of  $\text{MoM}_{k,b=\frac{n}{k}}$  (SI Dataset S1), when  $k$  is small.  
This highlights the superiority of the median Hodges-Lehmann  
mean over the median of means, as it not only can provide an  
accurate estimate for moderate sample sizes, but also allows  
the use of quasi-bootstrap, where the bootstrap size can be  
adjusted as needed.

**Data Availability.** Data for Figure ?? are given in SI Dataset S1. All codes have been deposited in [GitHub](#).

**ACKNOWLEDGMENTS.** I sincerely acknowledge the insightful comments from the editor which considerably elevated the lucidity and merit of this paper.

1. J Hodges Jr, E Lehmann, Estimates of location based on rank tests. *The Annals Math. Stat.* **34**, 598–611 (1963).
2. PK Sen, On the estimation of relative potency in dilution (-direct) assays by distribution-free methods. *Biometrics* pp. 532–552 (1963).
3. M Ghosh, MJ Schell, PK Sen, A conversation with pranab kumar sen. *Stat. Sci.* pp. 548–564 (2008).
4. RJ Serfling, Generalized L-, m-, and r-statistics. *The Annals Stat.* **12**, 76–86 (1984).
5. A Ehsanes Saleh, Hodges-lehmann estimate of the location parameter in censored samples. *Annals Inst. Stat. Math.* **28**, 235–247 (1976).
6. L Li, H Shao, R Wang, J Yang, Worst-case range value-at-risk with partial information. *SIAM J. on Financial Math.* **9**, 190–218 (2018).
7. J Hodges, E Lehmann, Matching in paired comparisons. *The Annals Math. Stat.* **25**, 787–791 (1954).
8. K Chung, Sur les lois de probabilité unimodales. *COMPTES RENDUS HEBDOMADAIRES DES SEANCES DE L ACADEMIE DES SCIENCES* **236**, 583–584 (1953).
9. L Devroye, M Lerasle, G Lugosi, RI Oliveira, Sub-gaussian mean estimators. *The Annals Stat.* **44**, 2695–2725 (2016).
10. RD Richtmyer, A non-random sampling method, based on congruences, for" monte carlo" problems, (New York Univ., New York. Atomic Energy Commission Computing and Applied . . . ), Technical report (1958).
11. IM Sobol', On the distribution of points in a cube and the approximate evaluation of integrals. *Zhurnal Vychislitel'noi Matematiki i Matematicheskoi Fiziki* **7**, 784–802 (1967).
12. KA Do, P Hall, Quasi-random resampling for the bootstrap. *Stat. Comput.* **1**, 13–22 (1991).