Semiparametric robust mean estimations based on the orderliness of quantile averages

Tuban Lee

This manuscript was compiled on June 9, 2023

semiparametric | mean-median-mode inequality | asymptotic | unimodal | Hodges—Lehmann estimator

Classifying Distributions by the Signs of Derivatives

- Let $\mathcal{P}_{\mathbb{R}}$ denote the set of all continuous distributions over \mathbb{R} and $\mathcal{P}_{\mathbb{X}}$ denote the set of all discrete distributions over a countable set X. The primary focus of this article will be on the class of continuous distributions, $\mathcal{P}_{\mathbb{R}}$. However, it's worth noting that most discussions and results can be extended to encompass the discrete case, $\mathcal{P}_{\mathbb{X}}$, unless explicitly specified otherwise. Besides fully and smoothly parameterizing them by a Euclidean parameter or merely assuming regularity conditions, there exist additional methods for classifying distributions based on their characteristics, such as their skewness, peakedness, modality, and supported interval. In 1956, Stein initiated the study of estimating parameters in the presence of an infinite-dimensional nuisance shape parameter (1) and proposed a necessary condition for this type of problem, a contribution later explicitly recognized as initiating the field of semiparametric statistics (2). In 1982, Bickel simplified Stein's general heuristic necessary condition (1), derived sufficient conditions, and used them in formulating adaptive estimates for the parameters in several semiparametric models (2).
- Data Availability. Data for Figure ?? are given in SI Dataset S1. All codes have been deposited in GitHub.
- ACKNOWLEDGMENTS. I sincerely acknowledge the insightful comments from the editor which considerably elevated the lucidity and merit of this paper.
- CM Stein, Efficient nonparametric testing and estimation in *Proceedings of the third Berkeley* symposium on mathematical statistics and probability. Vol. 1, pp. 187–195 (1956).
- 2. PJ Bickel, On adaptive estimation. *The Annals Stat.* **10**, 647–671 (1982).