

P8160 - Hurricane Project

Tucker Morgan - tlm2152

4/26/2022

Given Information

The following Bayesian model was suggested.

$$Y_i(t+6) = \beta_{0,i} + \beta_{1,i}Y_i(t) + \beta_{2,i}\Delta_{i,1}(t) + \beta_{3,i}\Delta_{i,2}(t) + \beta_{4,i}\Delta_{i,3}(t) + \epsilon_i(t)$$

where $Y_i(t)$ the wind speed at time t (i.e. 6 hours earlier), $\Delta_{i,1}(t)$, $\Delta_{i,2}(t)$ and $\Delta_{i,3}(t)$ are the changes of latitude, longitude and wind speed between t and $t+6$, and $\epsilon_{i,t}$ follows a normal distributions with mean zero and variance σ^2 , independent across t .

In the model, $\beta_i = (\beta_{0,i}, \beta_{1,i}, \dots, \beta_{7,i})$ are the random coefficients associated the i th hurricane, we assume that

$$\beta_i \sim N(\beta, \Sigma)$$

follows a multivariate normal distributions with mean β and covariance matrix Σ .

We assume the following non-informative or weak prior distributions for σ^2 , β and Σ .

$$P(\sigma^2) \propto \frac{1}{\sigma^2}; \quad P(\beta) \propto 1; \quad P(\Sigma^{-1}) \propto |\Sigma|^{-(d+1)} \exp\left(-\frac{1}{2}\Sigma^{-1}\right)$$

d is dimension of β .

Task 1:

Let $\mathbf{B} = (\beta_1^\top, \dots, \beta_n^\top)^\top$, derive the posterior distribution of the parameters $\Theta = (\mathbf{B}^\top, \beta^\top, \sigma^2, \Sigma)$.

Note from given Bayesian model:

$$\epsilon_i(t) = Y_i(t+6) - \left(\beta_{0,i} + \beta_{1,i}Y_i(t) + \beta_{2,i}\Delta_{i,1}(t) + \beta_{3,i}\Delta_{i,2}(t) + \beta_{4,i}\Delta_{i,3}(t)\right) \stackrel{i.i.d}{\sim} N(0, \sigma^2)$$

$$f_{Y_i(t+6)}(Y_i | \beta, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma}} \exp \left\{ -\frac{1}{2\sigma^2} (Y_i(t+6) - (\beta_{0,i} + \beta_{1,i}Y_i(t) + \beta_{2,i}\Delta_{i,1}(t) + \beta_{3,i}\Delta_{i,2}(t) + \beta_{4,i}\Delta_{i,3}(t)))^2 \right\}$$

Therefore,

$$\pi(\beta | Y_i) \propto f_{Y_i(t+6)}(Y_i | \beta, \sigma^2) \times \pi(\beta) \times \pi(\sigma^2)$$

$$\Rightarrow \pi(\boldsymbol{\beta} \mid Y_i) \propto \frac{1}{\sqrt{2\pi}\sigma^3} \exp \left\{ -\frac{1}{2\sigma^2} (Y_i(t+6) - (\beta_{0,i} + \beta_{1,i}Y_i(t) + \beta_{2,1}\Delta_{i,1}(t) + \beta_{3,i}\Delta_{i,2}(t) + \beta_{4,i}\Delta_{i,3}(t))) \right\}$$

with joint likelihood

$$\pi(\boldsymbol{\beta} \mid Y_i) \propto \prod_{i=1}^n \frac{1}{\sqrt{2\pi}\sigma^3} \exp \left\{ -\frac{1}{2\sigma^2} (Y_i(t+6) - (\beta_{0,i} + \beta_{1,i}Y_i(t) + \beta_{2,1}\Delta_{i,1}(t) + \beta_{3,i}\Delta_{i,2}(t) + \beta_{4,i}\Delta_{i,3}(t))) \right\}$$

$$\log \pi(\boldsymbol{\beta} \mid Y_i) \propto \sum_{i=1}^n -\frac{1}{2} \log(2\pi) - 3 \log \sigma - \frac{1}{2\sigma^2} (Y_i(t+6) - (\beta_{0,i} + \beta_{1,i}Y_i(t) + \beta_{2,1}\Delta_{i,1}(t) + \beta_{3,i}\Delta_{i,2}(t) + \beta_{4,i}\Delta_{i,3}(t)))$$

Please, feel free to make edits or add work to this.