

## Title in Progress

Amy Pitts, Hun Lee, Jimmy Kelliher,  
Tucker Morgan, and Waveley Qiu

2022-02-21

# Motivation

- ▶ Identifying the effect of a treatment or intervention is one of the most fundamental tasks we encounter as biostatisticians. . .

# Motivation

- ▶ Identifying the effect of a treatment or intervention is one of the most fundamental tasks we encounter as biostatisticians. . .
- ▶ . . . but confounding variables can bias our estimates of treatment effects.

# Motivation

- ▶ Identifying the effect of a treatment or intervention is one of the most fundamental tasks we encounter as biostatisticians. . .
- ▶ . . . but confounding variables can bias our estimates of treatment effects.
- ▶ Propensity score matching (PSM) is a tool that can help us mitigate the effects of confounders. . .

# Motivation

- ▶ Identifying the effect of a treatment or intervention is one of the most fundamental tasks we encounter as biostatisticians. . .
- ▶ . . . but confounding variables can bias our estimates of treatment effects.
- ▶ Propensity score matching (PSM) is a tool that can help us mitigate the effects of confounders. . .
- ▶ . . . but there is no consensus on the best way to estimate standard errors when using the PSM algorithm.

# Motivation

- ▶ Identifying the effect of a treatment or intervention is one of the most fundamental tasks we encounter as biostatisticians. . .
- ▶ . . . but confounding variables can bias our estimates of treatment effects.
- ▶ Propensity score matching (PSM) is a tool that can help us mitigate the effects of confounders. . .
- ▶ . . . but there is no consensus on the best way to estimate standard errors when using the PSM algorithm.
- ▶ How can we assess which procedures reliably estimate standard errors?

# Motivation

A simulation study!

# Enter the Bootstrap

- ▶ Bootstrapping is one of the most common procedures for estimating standard errors.



# Enter the Bootstrap

- ▶ Bootstrapping is one of the most common procedures for estimating standard errors.
- ▶ The PSM algorithm intakes an unmatched dataset and outputs a matched one.


# Enter the Bootstrap

- ▶ Bootstrapping is one of the most common procedures for estimating standard errors.
- ▶ The PSM algorithm intakes an unmatched dataset and outputs a matched one.
- ▶ When do we execute the bootstrap - before the match or after it?

# Enter the Bootstrap

- ▶ Bootstrapping is one of the most common procedures for estimating standard errors.
- ▶ The PSM algorithm intakes an unmatched dataset and outputs a matched one.
- ▶ When do we execute the bootstrap - before the match or after it?
- ▶ Let's try both!

# Roadmap of the Simulation Study



## Data Generation - Continuous Outcome

For each individual  $i \in \{1, \dots, n\}$ , we consider covariates  $L_{1i}, L_{2i}, L_{3i} \sim N(0, 1)$ . Treatments are distributed according to law  $A_i \sim B(\pi_i)$ , where  $\pi_i$  - the true propensity to be treated - is subject to the data-generating process

$$\log \left( \frac{\pi_i}{1 - \pi_i} \right) = \alpha_0 + \alpha_1 L_{1i} + \alpha_2 L_{2i}.$$

Given this, we further define the data-generating process of our continuous outcome via

$$Y_i = \beta_1 A_i + \beta_2 L_{2i} + \beta_3 L_{3i} + \varepsilon_i,$$

where  $\varepsilon_i$  denotes random error. Because  $L_{2i}$  effects both  $A_i$  and  $Y_i$ , it acts as a confounder in estimating the treatment effect.

## Data Generation - Binary Outcome

For each individual  $i \in \{1, \dots, n\}$ , we consider covariates  $L_{1i}, L_{2i}, L_{3i} \sim N(0, 1)$ . Treatments are distributed according to law  $A_i \sim B(\pi_i)$ , where  $\pi_i$  - the true propensity to be treated - is subject to the data-generating process

$$\log \left( \frac{\pi_i}{1 - \pi_i} \right) = \alpha_0 + \alpha_1 L_{1i} + \alpha_2 L_{2i}.$$

Given this, we further define the data-generating process of our binary outcome via  $Y_i \sim B(\tau_i)$  where

$$\log \left( \frac{\tau_i}{1 - \tau_i} \right) = \beta_1 A_i + \beta_2 L_{2i} + \beta_3 L_{3i}.$$

Observe that we have omitted a random error term, as realizations of  $Y_i$  are innately subject to noise.