## MATH 3310 – Midterm Experience

Fall 2026

**Group:** Tucker Phillips, Peter Stevenson, Owen Harps

**Constraints:** You may discuss the problems below with the following set of people:


$$\begin{aligned} P \;=\; &\{\text{Anyone currently enrolled in MATH 3310,} \\ &\text{David Brown, Mcrae Arnold, Gideon Walker}\}. \end{aligned}$$

The only resources we may use to solve the problems posed in this experience are our brains. The use of anything beyond that is cheating.

The solutions to this Experience should be created by groups of at most four and are to be submitted October 31, 2026, by 11:59pm.

### ✠ Sub-Experience One: *Pizza Cutting with Lasers*

**No outside resources, please** Given the difficulty of organizing and sequentially lining up cuts to get the maximum number of pieces, you invent a laser cutting system that allows you to make all cuts at once. You place $n$ of your devices on the crust with spacing that will maximize the number of pieces made and, once activated (ideally remotely because they are dangerous), *laser sabers*® shoot between every pair of devices simultaneously cutting the pizza into $R(n)$ pieces*. Below is an illustration representing 5 of your devices cutting the pizza into 16 pieces.




Please determine a formula for $R(n)$.

**Answer:** To begin we decided to gather data to see how we could determine the relationship. We drew the pizza slices out with different numbers of nodes on our papers and gathered the following data:

| $n$ | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 |
|---|---|---|---|---|---|---|---|---|
| $P(n)$ | 1 | 1 | 2 | 4 | 8 | 16 | 31 | 57 |

One issue we saw while drawing out the different pizza diagrams is that it was easy to draw three points intersecting eachother. Since we are trying to maximize the number of slices we instead offset the nodes to avoid intersections of more then two lines. This helps us to maximize the number of pieces.

We decided to use the calculus method to solve this problem. To start we created a difference table as pictured below. It pictures the first through fifth difference of $P(n)$.

* "R" for laseR.

| $n$ | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 |
|---|---|---|---|---|---|---|---|---|
| $P(n)$ | 1 | 1 | 2 | 4 | 8 | 16 | 31 | 57 |
| $\Delta P(n)$ | 0 | 1 | 2 | 4 | 8 | 15 | 26 | |
| $\Delta^2 P(n)$ | 1 | 1 | 2 | 4 | 7 | 11 | | |
| $\Delta^3 P(n)$ | 0 | 1 | 2 | 3 | 4 | | | |
| $\Delta^4 P(n)$ | 1 | 1 | 1 | 1 | | | | |
| $\Delta^5 P(n)$ | 0 | 0 | 0 | | | | | |

Based on the fact that $\delta^5 P(n)$ is all zeroes we can assume that the equation $P(n)$ may be represented by a equation of this form:

$$P(n) = An^4 + Bn^3 + Cn^2 + Dn + E$$

To solve for the coefficents $A, B, C, D, E$ we want to create a system of equations we could use to solve for them. We use the data we gathered above to create a system of equations which is pictured below.

$$\begin{bmatrix} 0 & 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 1 & 1 \\ 16 & 8 & 4 & 2 & 1 \\ 81 & 27 & 9 & 3 & 1 \\ 256 & 64 & 16 & 4 & 1 \end{bmatrix} \begin{bmatrix} A \\ B \\ C \\ D \\ E \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 2 \\ 4 \\ 8 \end{bmatrix}$$

We plugged this equation into a matrix calculator to solve for the coefficents. We got the following results.

$$\begin{bmatrix} A \\ B \\ C \\ D \\ E \end{bmatrix} = \begin{bmatrix} 0.041667 \\ -0.250000 \\ 0.958333 \\ -0.750000 \\ 1.000000 \end{bmatrix}$$

We can plug these into our initial equation for the final equation:

$$\boxed{P(n) = 0.041667n^4 - .25n^3 + .95833n^2 - .75n + 1}$$

We tested this by plugging in all the values of n we had manually calculated, and found that this equation described our function for all values from $n = 0$ to $n = 7$

As we had more answers then we needed to solve the system of equations we are assuming that this equation has been proven and is correct.

---


### ✠ Sub-Experience Two: *Strange Walks*

After the pizza, you decide to take a walk.

**Number of shortest paths using the *city block metric*.** Starting at Center and Main, you decide to walk to X North and Y East. The distance is measured in blocks and so the distance from $X_1$ North and $Y_1$ East to $X_2$ North and $Y_2$ East is $|X_2 - X_1| + |Y_2 - Y_1|$.

**SE 2.1.** For yuks, determine $\pi$ in the Logan metric. Use the fact that $\pi$ is defined to be the ratio of the circumference to the diameter of a circle, and that a circle is the set of points equidistant from a given point (that distance is typically called the *radius* of the circle and the point is called its *center*). Remember that, in this context, '*distance*' has been redefined.

**Answer:** In this context we are using the logan metric, so the way we measure distance has changed. When we create a circle (A set of points equidistant from a given center) it ends up creating a diamond shape. This is pictured in the image below.




The equation relating circumference to the diameter of a circle is

$C = \pi * D$

Where $D = Diameter$ and $C = Circumference$

To solve for $\pi$ we need a circumference and diameter that correspond to eachother. Lets use a "circle" of diameter 4. This is what we used to create the diamond pictured above. To solve for $\pi$ we need to find the circumference of a "circle" of diameter 4.

Using the Logam metric we can find that the length of each of the individual sides of the diameter 4 diamond is equal to 4 where R is our radius. We have four sides so the circumference is equal to $4 * 4$

Now knowing both the circumference and the diameter we can plug these values in and solve for $\pi$


$\pi = 16 \div (4)$

$\boxed{\pi = 4}$

This is the final answer for this problem.

**SE 2.2.** Count the number of shortest paths from Center and Main to X North and Y East. A more erudite way to pose this problem is as follows: count paths from $(0,0)$ to $(X,Y)$ using steps of the form $R:(x,y)\mapsto(x+1,y)$ or $U:(x,y)\mapsto(x,y+1)$. So paths trace rectilinear "curves" in the first quadrant of the Cartesian coordinate system that visit only integer-valued coordinates.

**Proof:** In order to arrive at $(X,Y)$ from $(0,0)$ we will always end up having to take Y steps to the East or R and X steps to the North or U. For a few smaller $(X,Y)$ coordinates we can find the number of paths easily enough. To get to $(0,1)$ there is 1 path, we just take one step to the North. To get to $(1,0)$ there is also only 1 path, we just take one step to the East. To get to $(1,1)$ we can follow either of the two previous paths with an additional move to the North or East. We can follow this logic further and we find that the number of paths to $(X,Y)$ which we'll call $P(X,Y)=P(X-1,Y)+P(X,Y-1)$. If we start to populate a graph with these path numbers we start to notice that we can see Pascal's triangle as seen below.




Now that we recognize Pascal's triangle it's not too big of a leap to assume that we will need to use choose to count the number of paths. We need to take a total of $X+Y$ steps no matter what, so if we choose how we want to take X steps and just fill in the rest of the steps with the Y steps we find that the number of paths to $\boxed{(X,Y)=P(X,Y)=\binom{X+Y}{X}}$.

**SE 2.3.** Walk in a strange way. Start at Center and Main, thinking of it as the origin $(0,0)$ and take $n$ steps, each of type R, L, or U, with R never followed by L and vice-versa. The steps are defined via $R:(x,y)\mapsto(x+1,y)$, $L:(x,y)\mapsto(x-1,y)$, and $U:(x,y)\mapsto(x,y+1)$. Determine the number of different paths with $n$ steps.

We started this equation by drawing out the different potential paths that could be taken. I started by drawing out many paths. We decided to start by examining the potential paths the system could take. We quickly see the potential for a recursive formula. The number of possible routes for any n steps is related to the previous problem.

I see that the total number of possible routes is P(n) = (number of rights/lefts from P(n-1)) * 2 + (number of ups from P(n-1) * 3). This is because each path that last moved left or right has two options to move in, which each path that last moved up has three options to move in.

From this relationship I can figure out how many rights and lefts there were in $P(n-1)$ and how many ups there were in $P(n-1)$.

The number of times $p(n-1)$ moved up is equal to the value of $p(n-2)$. This is because every line in $p(n-2)$ moved up once in $p(n-1)$

The number of times $p(n-1)$ moved left or right is equal to $p(n-1)-p(n-2)$. This is the total value of $p(n-1)$ with the total number of "ups" subtracted, since any path that did not last move up must have moved left or right.

Now we can develop the recursive relationship $P(n) = P(n-2)*3 + (P(n-1) - P(n-2))*2$. This simplifies to $P(n) = P(n-2) + 2*P(n-1)$

We want a closed form solution instead of a recursive relationship. Lets use a characteristic equation.

Turn $P(n) = P(n-2) + 2P(n-1)$ into $R^n = 2*R^{n-1} + R^{n-2}$

Dividing both sides by $R^{n-2}$ we turn that equation into

$R^2 = 2*R + 1$

Solving for the quadratic roots of this equation we find the roots are:

$R = 2.41421, -0.414214$

Now we know P(n) is of the form $P(n) = A(2.41421)^n + B(-0.414214)^n$

Lets plug in some of our known stats to find what this value will be.

$P(0) = 1 = A + B$

$P(1) = 3 = A(2.41421) + B(-0.414214)$

We can solve for $A = 1.20710685527$ and $B = -0.207106855266$.

This gives us the final explicit equation of:


$$\boxed{P(n) = 1.207106*(2.41421)^n - 0.207106*(-0.414214)^n}$$

We plugged in various values of n that we had manually found the solutions for, and found this equation lined up for each tested case, proving the equation is true.

**SE 2.4.** Given what happens when you take the red pill, you choose the blue one ... and turn into a puma ... and find yourself in the Yucatan peninsula at the base of a Mayan temple. Whoa. You find that your puma strength allows you to clear one or two terraces in a single bound. For reasons known only by the metaphysics of the blue pill, you need to

1. Compute $P_n$, the number of ways to ascend an n-terraced temple. Please determine $P_n$ by finding a recurrence relation and then using the recurrence relation find a closed formula for $P_n$. Calculate $P_{365}$ because the temple you're facing has 365 terraces.
2. Again, for reasons which clear only to you as you are under the influence of the blue pill, you must prove that

$$P_n = \sum_{i=0}^{\lfloor n/2 \rfloor} \binom{n-i}{i}.$$



**S : SINGLE LEAP, D : DOUBLE LEAP**

| n | Leap Options | Total Paths |
|---|---|---|
| 0 | [No Jump] | 1 |
| 1 | [S] | 1 |
| 2 | [SS, D] | 2 |
| 3 | [DS, SD, SSS] | 3 |
| 4 | [DD, DSS, SSD, SDS, SSSS] | 5 |
| 5 | [DDS, DSSS, DSD, SDD, SSSD, SDSS, SSDS, SSSSS] | 8 |
| ... | ... | ... |

Table 1: Initial Leap Options for n = 0 to n = 5

1. $P_n$ will continue following this pattern shown in the total paths column. This recursive patterns is idential to the Fibonnacci Sequence, so $P_n$ can be described as $P_n = P(n-1) + P(n-2)$.

$$\begin{aligned} P_n &= P(n-1) + P(n-2) \\ r^n &= r^{n-1} + r^{n-2} \\ r^2 &= r + 1 \\ 0 &= r^2 - r - 1 \\ r_{1,2} &= \frac{1 \pm \sqrt{5}}{2} \end{aligned}$$

In order to find the closed form, we will use the initial conditions to find coefficients using a system of equations:

$$\begin{aligned} 1 &= A(r_1)^1 + B(r_2)^1 \\ 2 &= A(r_1)^2 + B(r_2)^2 \\ A &= \frac{1+\sqrt{5}}{2\sqrt{5}} \\ B &= -\frac{1-\sqrt{5}}{2\sqrt{5}} \end{aligned}$$

So the closed formula for $P_n$ is:

$$P_n = \frac{1+\sqrt{5}}{2\sqrt{5}}(\frac{1+\sqrt{5}}{2})^n - \frac{1-\sqrt{5}}{2\sqrt{5}}(\frac{1-\sqrt{5}}{2})^n$$

$$\boxed{P_n = (\frac{1+\sqrt{5}}{2\sqrt{5}})^{n+1} - (\frac{1-\sqrt{5}}{2\sqrt{5}})^{n+1}}$$

$P_{365}$ can now be calculated:

$$P_{365} = (\frac{1+\sqrt{5}}{2\sqrt{5}})^{366} - (\frac{1-\sqrt{5}}{2\sqrt{5}})^{366}$$

$$\boxed{P_{365} \approx 1.380 \cdot 10^{78}}$$

2.

$$P_n = \sum_{i=0}^{\lfloor n/2 \rfloor} \binom{n-i}{i}$$

.

**Proof:** Essentially, $P_n$ counts the combinations of Single and Double leaps that add to n. If we say that d is the number of double leaps in the sequence, then each d double jumps will go up 2d terraces. subtracting that from the total n terraces gives the terraces left over for single leaps, $n-2d$ (which is also the number of single leaps). So the total number of leaps is the combination of the number of single and double leaps: $d+n-2d$ or simplified to $n-d$.

We can describe a sequence by deciding where the "double jumps" d are for a given total leap combos $n-d$. This turns out to be $\binom{n-i}{i}$ ways to place them. But there can only by $n/2$ double jumps maximum to reach the top. Summing all of these possibilitees gives us the exact formula above swapping i for d:

$$P_n = \sum_{i=0}^{\lfloor n/2 \rfloor} \binom{n-d}{d}$$

.

---

### ✠ Sub-Experience Three: The Master Table of Distributions.

Below is the table of distribution functions $f : A \to B$, where A is a finite set with n elements and B is a finite set with x elements. The goal: determine a formula for the number of functions with properties described by the table for each of the twelve possibilities. So, for example, the entry in cell 5 of the table is to be the number of injective functions that map a set of n indistinguishable object into a set of x distinguishable objects.

| A | B | unrestricted | injective | onto |
|---|---|---|---|---|
| distinguishable | distinguishable | 1. | 2. | 3. |
| indistinguishable | distinguishable | 4. | 5. | 6. |
| distinguishable | indistinguishable | 7. | 8. | 9. |
| indistinguishable | indistinguishable | 10. | 11. | 12. |

Please determine, with an argument for each, a formula for each entry in the table.

If there is labeled there wll prpobably be a fatorial
Onto column will likeely have squeezer x, k
Stars and bars could prove the indistinguishable indistinguishable case

1. When we have n indistinguishable objects being placed into x distinguishable groups without restriction every object has x group choices where it can go. With n objects this leads to the equation: $\boxed{x^n}$

2. When we have n distinguishable objects being placed into x distinguishable groups. The first object has x possible groups where it can be placed. The next object has x - 1 possible places where it can be placed. This pattern will continue where each consequtive ball will have -1 places where it could possibly be placed. The number of group choices can be represented by n. We can represent this pattern as $\boxed{n^{\underline{x}}}$

3. Because both A and B are distinguishable, and "onto" is the surjective case, the squeeze function can be used to partition n elements into exactly x groups ($\left\{ {n \atop x} \right\}$). Multiplying by x! after partitioning gives the number of functions that map the "onto" case: $\boxed{x! \cdot \left\{ {n \atop x} \right\}}$

4. We can think of this cell using the stars and bars method. We have $n$ balls and $x-1$ bars to place to create $x$ boxes. So, we have a set of $n+x-1$ objects and we can choose $n$ places to put the balls first, then place the bars. This means that there are $\left(\!\binom{x}{n}\!\right) = \boxed{\binom{n+x-1}{n}}$ ways to do this.

5. If $n \geq x$ there is one way. This is because there are more balls than boxes and with it being injective, each box can have at most one ball. If $n < x$ there are $\boxed{\binom{x}{n}}$ ways to distribute the balls. We can think of this as having x boxes and then we choose n of them to receive 1 ball each, and we leave the rest empty.

6. Our first n objects are going to be placed in x predetermined spots since we have to have at least one A object in every B group. After that we have n - x more balls to place in n boxes. Our boxes are distinguishable but our balls are not. This suggests we should use some sort of binomial, where we choose boxes for each ball to go into.

   Usings stars and bars after placing our initial x each of our n - x balls has x - 1 possible options to go to. $\binom{(n-x)+x-1}{x-1}$

   Which can be simplified to $\boxed{\binom{n-1}{x-1}}$

7. Because the elements are indistinguishable we want to look at how the n elements in A are grouped into non-empty subsets. Each of the individual groups can be represented by a stirling number partion. We want to sum the different stirling number paritions to see how many different ways we can do this for each different value of k. $\boxed{\sum_{k=1}^{x} S(n,k)}$

8. In the injective case when the distinguishable objects in A are placed in indistinguishable objects in B there is only $\boxed{1}$ possible solution. No matter which order you place each individual A object into each B object all cases will be the same.

9. This case matches the description of stirling numbers perfectly. It is a mapping of n elements onto k indistinguishable elements where each element is non-empty. $\boxed{\left\{ {n \atop k} \right\}}$

10. In the case with both A and B indistinguishable objects we want a partition of n. We care about how many times the n objects can be divided into seperarte groups. $\boxed{P(n)}$

11. When both A and B are indistinguishable in the injective case there is only $\boxed{1}$ way to organize. Whatever combination you place one ball into each box all solutions will be the same.

12. In this case with indistinguishable objects placed into indistinguishable boxes we can see that there will be some partition of possible combinations. Since the order doesn't matter and we just want a combination. Since we have a maximum of x A objects placed onto n B objects we can figure out there is a integer partition $\boxed{P_x(n)}$

✠ **Sub-Experience Four:** ***Game Time!***

**SE 4.1: Modified Towers of Hanoi.** The *Towers of Hanoi* are mythical diamond needles, three of them, with 64 gold discs impaled on one of the needles. God told the monks of Brahma to transfer all the discs to another needle with the constraints that no larger disc ever be placed atop a smaller disc, and only one disc at a time is to be moved. Once the monks finish moving the discs the World will end. Evidently the monks have not yet finished.

Label the needles of the Towers of Hanoi L, M, and R, for the left, middle, and right, respectively. Consider the original Towers of Hanoi game, but with the additional constraint that a disk can only be moved to an adjacent peg; that is, a disk can only be moved to M from L or R, and can only be moved to L or R from M. Assume all discs are initially on L.

*Determine the minimum number of moves (where a* move *is defined to be the transfer of one disc from one needle to another) required to transfer* n *discs from* L *to* R.

**Answer:** We started by manually computing the number of moves required to move each of the individual discs from the L needle to the R needle. We ran these calculations up to N = 6. We were able to find the number of moves required to move from the L needle to the R needle for the number of n pins using the chart below.

In this chart the uppermost row tells us how many discs we are trying to move. The next rows tell us how many times each discs had to be moved in order to move all discs from L to R. For example, when we are moving 2 discs, we had to move disc 1 4 times and disc 2 2 times.

| | **1** | **2** | **3** | **4** | **5** | **6** |
|---|---|---|---|---|---|---|
| **1** | 2 | 4 | 8 | 16 | 32 | 64 |
| **2** | | 2 | 4 | 8 | 16 | 32 |
| **3** | | | 2 | 4 | 8 | 16 |
| **4** | | | | 2 | 4 | 8 |
| **5** | | | | | 2 | 4 |
| **6** | | | | | | 2 |

This chart revealed a pattern that to move N discs there is a summation we can use. By observation I was able to see the following sum we can use to represent how many moves it takes to move N discs.

$$\boxed{\sum_{k=1}^{n}\sum_{w=1}^{k} 2^w}$$


*How Long?* Assume the monks of Brahma were given this game with 64 discs, and can move a disc at a rate of one per second. How long in centuries will it take for the monks to complete their task?

**Answer:** Next we want to see how long it would take the monks to move 64 discs if each move takes 1 second. We can input $n = 64$ into the equation above which tells us we will need approximately $7.3786976295 \times 10^{19}$ seconds, which is equal to $4.737994774 \times 10^{17}$ centuries. A very long time.

**SE 4.2: A Two-Player Game with no Dire Consequences.**

**The Game**

Two players, A and B, alternately select an *edge* (line segment connecting two dots) on the grid graph shown below and color it red. The loser of the game is the player who is forced to select an edge that creates a red $C_4$ — a red cycle on 4 vertices.



**The Fun**

*Confirm or deny, with proof, whether player A (the first player) can always win if she employs a particular strategy for each move.*
On a 5 by 6 point grid, there are a total of 49 placements for possible edges. In order to create a $C_4$ cycle on 4 vertices, there needs to be 4 edges placed.

**Claim:** Player A can always win if a particular strategy is used.

**Proof:**
The strategy that player A needs to employ in order to always win is as follows:

1. Begin each game with an opening move between the center two dots.
2. Allow Player B to make edges.
3. Player A should always mirror Player B's edges by 180°.

The reason this strategy will always allow player A to win is because every edge can be mirrored besides the very center, but Player A has already filled the center. Player B can either place an edge in a "safe place" (non cycle completing) or place an edge that loses them the game. Since the grid is mirrored and assuming Player B didn't lose by completing a cycle, Player A's placement will be in a similarly "safe place" on the other half of the grid. This pattern will continue until Player B is forced to be the first player to complete a cycle.

## ✠ Sub-Experience Five – A Matter of Life and Death

**No outside resources** You are among a group of $n - 1$ of your friends (pretend you have $n - 1$ friends even if you don't, where $n$ is possibly very large) and are captured by a horde of theater students who will force you and your $n - 1$ friends to enact the play *Cats* with them over and over and over and ... . Your friends choose death over this fate and decide to form a circle and have every other person commit suicide (someone has a pistol and $n$

bullets, which is quite reasonable to assume here at USU) until only one person survives, who will supposedly kill themselves. You want no part of this suicide madness since, you figure, the theater people will tire eventually and you can make your escape by taking Jennyanydots hostage with the pistol and remaining bullet and escape.

Number you and your friends 1 to $n$ and assume you all form a vicious circle to facilitate the suicide.

Question: *In which position (call it $S(n)$) must you be in order to survive?*

Examples: $S(4) = 1$, $S(7) = 7$.

---

**Proof:** Let's start by gathering some data.

| $n$ | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| $S(n)$ | 1 | 1 | 3 | 1 | 3 | 5 | 7 | 1 | 3 | 5 | 7 | 9 | 11 | 13 | 15 | 1 |

From this data we can observe a pattern where on each power of 2 $S(n)$ resets to 1, then in between each power of 2 it counts up to the next odd number.

We can then make a function that describes the pattern we observe. It is: $S(n) = 2\left(n - 2^{\lfloor \log_2 n \rfloor}\right) + 1$.

To explain a bit more about where this function comes from, we can consider the fact that we can represent any positive integer $n$ as $n = 2^l + r$, the sum of some power of 2 and a remainder. We can write this as $r = n - 2^l$. When we do $\lfloor \log_2 n \rfloor$ it gives us the largest integer that when 2 is raised to that power the result can fit inside of $n$. Now we have $r = n - 2^{\lfloor \log_2 n \rfloor}$. To take the final step towards getting the equation above, we need to recall that the pattern we observed started at 1 on each power of two, and would then count up by two, so we get $S(n) = 2r + 1 = 2\left(n - 2^{\lfloor \log_2 n \rfloor}\right) + 1$.

We will now use this equation as the starting point for a proof by induction.

**Assumption:** $S(n) = 2\left(n - 2^{\lfloor \log_2 n \rfloor}\right) + 1$, for all $k$ such that $1 \leq k \leq n$.

As the base case $S(1) = 2\left(1 - 2^{\lfloor \log_2 (1) \rfloor}\right) + 1 = 2(1 - 1) + 1 = 2(0) + 1 = 1$. We know this is true because of how we defined the suicide circle to work, if it's just you, then you survive.

Now we consider $S(n+1)$. This produces two cases: one where $n+1$ is an even number and the other where $n+1$ is an odd number. The difference in these cases becomes important when we try to find a recurrence. For example, if we have 10 people in the circle after we go around the circle once we will have five people, so the old 9 becomes the new 5, the old 7 becomes the new 4, 5 becomes 3, 3 becomes 2, and 1 is still 1. So for the case where $n+1$ is even, we have the recurrence $S(n+1) = S(2n) = 2S(n) - 1$. For the odd case, we get the recurrence $S(n+1) = S(2n+1) = 2S(n) + 1$.

The even case: $S(n+1) = S(2k) = 2S(k) - 1$
$= 2\left(2\left(k - 2^{\lfloor \log_2 k \rfloor}\right) + 1\right) - 1$
$= 2\left(2k - 2^{\lfloor \log_2 k \rfloor + 1}\right) + 2 - 1$
$= 2\left(2k - 2^{\lfloor \log_2 (k) \rfloor + \log_2 (2)}\right) + 1$
$= 2\left(2k - 2^{\lfloor \log_2 (2k) \rfloor}\right) + 1$ using the property of log's that $\log_a b + \log_a c = \log_a bc$
$2\left(2k - 2^{\lfloor \log_2 (2k) \rfloor}\right) + 1 = 2\left(2k - 2^{\lfloor \log_2 (2k) \rfloor}\right) + 1$
The first equality follows from the induction assumption, and the others from algebraic manipulations. So, $S(n+1) = S(2k) = 2S(k) - 1$ and we know the even case works.

The Odd case: $S(n+1) = S(2k+1) = 2S(k) + 1$
$= 2\left(2\left(k - 2^{\lfloor \log_2 k \rfloor}\right) + 1\right) + 1$

$= 2\left(2k - 2^{\lfloor \log_2 k \rfloor + 1}\right) + 2 + 1$
$= 2\left(2k + 1 - 2^{\lfloor \log_2 (2k) \rfloor}\right) + 1$
$= 2\left(2k + 1 - 2^{\lfloor \log_2 (2k+1) \rfloor}\right) + 1$. For this last step we can simply add the one because adding a 1 inside the floor of this log won't change when $n + 1$ is odd.
$= 2\left(2k + 1 - 2^{\lfloor \log_2 (2k+1) \rfloor}\right) + 1 = 2\left(2k + 1 - 2^{\lfloor \log_2 (2k+1) \rfloor}\right) + 1$
The first equality follows from the induction assumption, and the others from algebraic manipulations. So, $S(n+1) = S(2k+1) = 2S(k) + 1$ and we now know that the odd case works.

With both the even and the odd case verified in this way we can say that the result therefore follows by the PSMI and $\boxed{S(n) = 2\left(\frac{1}{2} + n - 2^{\lfloor \log_2 n \rfloor}\right)}$.

■

---