

Here are some Lemmas which are referenced throughout the problems.

**Lemma 1** (Sum of Ideals is an Ideal). *If  $I, J$  are ideals of a ring  $R$ , then  $I + J = \{i + j \mid i \in I, j \in J\}$  is also an ideal of  $R$ .*

*Proof.*  $\forall (i_1 + j_1), (i_2 + j_2) \in I + J, (i_1 + j_1) - (i_2 + j_2) = (i_1 - i_2) + (j_1 - j_2) \in I + J \implies I + J \leq_+ R$ . Next for all  $r \in R$  and all  $(i + j) \in I + J, r(i + j) = ri + rj, (i + j)r = ir + jr \in I + J$  since  $ir, ri \in I$  and  $rj, jr \in J$ . Thus,

If  $I, J$  are ideals of  $R$ , then  $I + J$  is an ideal of  $R$ .

□

**Lemma 2** (Inclusion Mapping is Ring Embedding). *If  $S \subseteq R$  is a subring of a ring  $R$ , then the inclusion mapping  $\iota : S \hookrightarrow R$  defined via  $\iota(s) = s$  is a ring embedding (an injective ring homomorphism).*

*Proof.* For all  $a, b \in S$ ,

$$\begin{aligned}\iota(a + b) &= a + b = \iota(a) + \iota(b) \\ \iota(ab) &= ab = \iota(a)\iota(b) \\ \iota(1_S) &= 1_S = 1_R \\ \iota(a) &= \iota(b) \implies a = b \text{ by definition.}\end{aligned}$$

□

**Lemma 3** (Evaluation Mapping to coefficient ring is a Surjective Ring Homomorphism). *The evaluation homomorphism  $\epsilon_a : R[x] \rightarrow R$  defined by  $\epsilon_a(f(x)) = f(a)$  is a surjective ring homomorphism for any  $a \in R$ .*

*Proof.* For all  $f(x), g(x) \in R[x]$ ,

$$\begin{aligned}\epsilon_a(f(x) + g(x)) &= f(a) + g(a) = \epsilon_a(f(x)) + \epsilon_a(g(x)) \\ \epsilon_a(f(x)g(x)) &= f(a)g(a) = \epsilon_a(f(x))\epsilon_a(g(x)) \\ \epsilon_a(1_{R[x]}) &= 1_R \\ \text{For any } r \in R, \epsilon_a(r) &= r \text{ so } \epsilon_a \text{ is surjective.}\end{aligned}$$

□

**Lemma 4** (Telescoping Identity). *For any  $k \in \mathbb{N}$  and any polynomial  $x^k - y^k$  over a commutative ring  $R$  where  $x, y$  may either be indeterminates or elements of  $R$ , we have that:*

$$x^k - y^k = (x - y) \sum_{\substack{i+j=k-1 \\ i, j \in \mathbb{N}}} x^i y^j.$$

*Proof.*

$$\begin{aligned}
 x \sum_{\substack{i+j=k-1 \\ i,j \in \mathbb{N}}} x^i y^j &= x(x^{k-1} + x^{k-2}y + \cdots + xy^{k-2} + y^{k-1}) = x^k + (x^{k-1}y + \cdots + x^2y^{k-2} + xy^{k-1}) \\
 y \sum_{\substack{i+j=k-1 \\ i,j \in \mathbb{N}}} x^i y^j &= y(x^{k-1} + x^{k-2}y + \cdots + xy^{k-2} + y^{k-1}) = (yx^{k-1} + x^{k-2}y^2 + \cdots + xy^{k-1}) + y^k \\
 &= (x^{k-1}y + x^{k-2}y^2 + \cdots + xy^{k-1}) + y^k \\
 \implies (x-y) \sum_{\substack{i+j=k-1 \\ i,j \in \mathbb{N}}} x^i y^j &= x^k + (x^{k-1}y + \cdots + x^2y^{k-2} + xy^{k-1}) - (x^{k-1}y + x^{k-2}y^2 + \cdots + xy^{k-1}) - y^k \\
 &= x^k - y^k.
 \end{aligned}$$

□

**Lemma 5** (Polynomials over fields have division algorithm). *If  $\mathbb{K}$  is a field, then for all  $f(x), g(x) \in \mathbb{K}[x]$ , where  $g(x)$  is non-zero there exists  $q(x), r(x) \in \mathbb{K}[x]$  such that either  $r(x) = 0$  or  $\deg(r(x)) < \deg(g(x))$  and then  $f(x) = q(x)g(x) + r(x)$ .*

*Proof.* Let  $f(x) = \sum_{i=1}^n a_i x^i, g(x) = \sum_{j=1}^m b_j x^j \in \mathbb{K}[x]$  have degrees  $m, n$ , respectively, such that  $g(x)$  is non-zero. Since  $\mathbb{K}$  is a field, all non-zero coefficients are units. Note that leading terms must be non-zero.

If  $f(x) = 0$ , then  $f(x) = 0 = 0g(x) + 0$  and so we have  $q(x) = 0, r(x) = 0$  such that  $f(x) = q(x)g(x) + r(x)$ , which satisfies the statement. If  $\deg(f(x)) = n < m = \deg(g(x))$ , then set  $q(x) = 0, r(x) = f(x)$ , and then  $f(x) = q(x)g(x) + r(x) = 0(g(x)) + f(x)$  satisfies the statement since  $\deg(r(x)) = \deg(f(x)) < \deg(g(x))$ .

Lastly, if  $\deg(f(x)) = n \geq m = \deg(g(x))$ , set  $t(x) = a_n b_m^{-1} x^{n-m}$ . Then we have that the leading term of  $t(x)g(x)$  is  $(a_n b_m^{-1} x^{n-m})(b_m x^m) = a_n x^n$ , which is leading term of  $f(x)$ .

Now set  $f_1(x) = f(x) - t(x)g(x)$ . Then the  $n$ -coefficient will vanish;  $[x^n](f(x) - t(x)g(x)) = [x^n]f(x) - [x^n]t(x)g(x) = a_n - a_n = 0$  and so  $\deg(f_1(x)) < \deg(f(x)) = n$ . If  $\deg(f_1(x)) < m = \deg(g(x))$  then we have  $f(x) = t(x)g(x) + f_1(x)$ , which satisfies the statement.

Otherwise if  $\deg(f_1(x)) < m = \deg(g(x))$ , we repeat the process. For each step  $i \geq 1$ , let  $d_i = \deg(f_i(x))$  set  $t_i(x) = ([x^{d_i}]f_i(x))(b_m^{-1})x^{m-d_i}$ . Now set  $f_{i+1}(x) = f_i(x) - t_i(x)g(x)$ . Then

$$[x^{d_i}]f_{i+1}(x) = ([x^{d_i}]f_i(x)) - ([x^{d_i}]f_i(x))b_m^{-1} = 0 \text{ and } f_i(x) = t_i(x)g(x) + f_{i+1}$$

So we see that by recursion, at each step  $i \mapsto i+1$ ,  $\deg(f_{i+1}(x)) < \deg(f_i(x))$ . Suppose our process terminates at some  $i = k$  such that  $f_k(x)$  has degree  $\mu \geq m$ . That is  $f_{k+1}(x)$  is not an acceptable polynomial in  $\mathbb{K}[x]$ ; it has negative powers of  $x$ . So  $f(x) = t(x)g(x) + \sum_{i=1}^k f_i(x) = (t(x) + \sum_{i=1}^{k-1} t_i(x))g(x) + f_k(x)$ . Well,  $t_k(x) = ([x^\mu]f_k(x))(b_m^{-1})x^{m-\mu}$  must have positive degree, and so  $t_k(x)g(x)$  has the same leading coefficient as  $f_k(x)$ . But then  $f_{k+1}(x)$  is a polynomial in  $\mathbb{K}[x]$  with  $\deg(f_{k+1}(x)) < \deg(f_k(x))$ . So our process must terminate with at some  $i = r$  such that  $f_r(x)$  has degree strictly less than  $m$ . That is:

$$f(x) = t(x)g(x) + \sum_{i=1}^r f_i(x) = (t(x) + \sum_{i=1}^{r-1} t_i(x))g(x) + f_r(x) \text{ where } \deg(f_r(x)) < m = \deg(g(x)).$$

Which satisfies the statement. By exhaustion, the statement is proven.

□

**Lemma 6.** *Let  $R$  be a commutative ring with unity.  $R/I$  is a field if and only if  $I$  is a maximal ideal of  $R$ .*

*Proof.* (  $\Leftarrow$  ) For any non-zero  $\bar{q} \in R/I$  there exists some non-zero reduced representative  $q \in R \setminus I$  such that  $\bar{q} = [q]_I$ . If  $I$  is maximal, then it's proper and then  $I \subset I + \langle q \rangle \subseteq R \implies I + \langle q \rangle = R$ , since  $I$  is maximal. So  $1 \in I + \langle q \rangle$  and therefore  $\exists i \in I, \exists pq \in \langle q \rangle$  such that

$$1 = i + pq \implies [1]_I = [i + pq]_I = [0]_I + [p]_I [q]_I \in R/I.$$

So  $[q]_I$  is a unit. Therefore, we see all non-zero elements of  $R/I$  are units and so  $R/I$  is a field.

(  $\Rightarrow$  ) If  $R/I$  is a field, let  $J$  be an ideal of  $R$  with  $I \subseteq J \subseteq R$ . Then  $J/I = \{j + I : j \in J\}$  is an ideal of  $R/I$  by the **Third Isomorphism Theorem**. Well, since  $R/I$  is a field, its only ideals are  $\{0\}$  and  $R/I$ , so  $J/I = \{0\}$  or  $J/I = R/I$ .

If  $J/I = \{0\}$ , then  $J = I$ . Otherwise, If  $J/I = R/I$ , then  $[1]_I \in J/I$ , so there exists  $j \in J$  with  $[1]_I = [j]_I$ . That is,  $j - 1 \in I$ . So then  $j - (j - 1) = 1 \in J \implies r(1) = r \in J, \forall r \in R \implies J = R$ . So  $J = I$  or  $J = R$  and therefore  $I$  is maximal.

Thus,

If  $R$  is a commutative ring with unity, then  $R/I$  is a field if and only if  $I$  is a maximal ideal of  $R$ .

□

**Lemma 7** (Zorn's Lemma). *Let  $(P, \leq)$  be a partially ordered set. If every chain in  $P$  has an upper bound in  $P$ , then  $P$  contains a maximal element.*

**Problem 1.** In the following problems, we investigate the relationship between integral domains and fields.

- (a) Prove that every field is an integral domain.
- (b) Prove that if  $R$  is a finite integral domain, then  $R$  is a field. **Hint:** Consider the set function  $a : R \rightarrow R$  given by multiplication by an element  $a$  (it will not be a ring homomorphism). If  $a$  is a non-zero-divisor, prove this is an injective function, and so must also be surjective since  $R$  has finitely many elements.
- (c) Find an example of an integral domain which is not a field.

*Proof.* (a) Let  $\mathbb{K}$  be a field, and suppose it contains some non-zero zero divisor  $b \in \mathbb{K} \setminus \{0\}$ . Then there exists some  $a \in \mathbb{K} \setminus \{0\}$  such that  $ab = ba = 0$ . Since  $a, b \in \mathbb{K} \setminus \{0\}$ , they are both units and so there exist  $a^{-1}, b^{-1} \in \mathbb{K}$  such that  $a^{-1}a = bb^{-1} = 1 \implies a^{-1}(ab)b^{-1} = 1 = a^{-1}(0)b^{-1} = 0 \implies 1 = 0$ . But then  $\mathbb{K} = \{0\}$  which is not a field, a contradiction. Therefore, all non-zero elements are not zero divisors and so  $\mathbb{K}$  is a commutative ring with unity and no zero divisors, which is the definition of an integral domain.

(b) Let  $R$  be a finite integral domain, and let  $a$  be some non-zero zero divisor  $a \in R \setminus \{0\}$ . We show the mapping  $a : R \rightarrow R$  via  $a(x) = ax, \forall x \in R$  is a bijection. For all  $x, y \in R$ ,

$$\text{[1-1]} \quad a(x) = a(y) \implies ax = ay \implies ax - ay = a(x - y) = 0 \implies x = y$$

otherwise  $a$  is a zero divisor, and we get a contradiction.

$$\text{[Onto]} \quad a : R \hookrightarrow R \implies \text{the domain is in bijection with its image} \implies |R| = |a(R)|$$

and since  $a(R) \subseteq R$  (the domain is equal to the codomain), we must have that  $a(R) = R$ .

(This image is just the orbit of  $a$  in the canonical multiplicative monoid action of  $R$  on itself.) So then for each non-zero zero divisor  $a \in R \setminus \{0\}$ ,  $\exists x \in R$  such that  $a(x) = ax = 1$ . That is, each non-zero zero divisor, which is just all of  $R \setminus \{0\}$  is a unit. Therefore,  $R$  is a commutative ring with unity and inverses for all non-zero elements, which is the definition of a field.

(c)  $\mathbb{Z}$  is an integral domain but not a field.

□

**Problem 2.** Let  $I, J$  be ideals in a commutative ring  $R$  such that  $I + J = (1)$ . Prove that  $IJ = I \cap J$ .

*Proof.* ( $\subseteq$ ) Recall that since  $I, J$  are ideals of  $R$ ,  $ir \in I$  and  $rj \in J, \forall i \in I, \forall j \in J, \forall r \in R$ . So then for all  $ij \in IJ$ , we have that  $(i)j \in I$  and  $i(j) \in J \implies ij \in I \cap J \implies IJ \subseteq I \cap J$ .

( $\supseteq$ )  $I + J = (1) \implies \hat{i} + \hat{j} = 1$  for some  $\hat{i} \in I, \hat{j} \in J$ . Then for all  $a \in I \cap J$ ,  $a = a(1) = (1)a = a(\hat{i} + \hat{j}) = a\hat{i} + a\hat{j} \in IJ + JI = IJ$ , since commutativity in  $R$  implies  $IJ = JI$ . Therefore,  $I \cap J \subseteq IJ$ .

Thus,

$$IJ = I \cap J$$

.

□

**Problem 3.** Let  $\varphi : R \rightarrow S$  be a ring homomorphism, and let  $J$  be an ideal of  $S$ . Prove that  $I = \varphi^{-1}(J) = \{i \in R \mid \varphi(i) \in J\}$  is an ideal of  $R$ .

*Proof.*  $\forall a, b \in \varphi^{-1}(J)$ ,  $\varphi(a), \varphi(b) \in J$  and so  $\varphi(a - b) = \varphi(a) - \varphi(b) \in J$ . So then  $a - b \in \varphi^{-1}(J)$  and  $\varphi^{-1}(J) \leq_+ R$ .

Next, for any  $r \in R$  and any  $i \in \varphi^{-1}(J)$ ,  $\varphi(ri) = \varphi(r)\varphi(i) \in J$  and  $\varphi(ir) = \varphi(i)\varphi(r) \in J$ , since  $J$  is an ideal of  $S$ . So then  $ri, ir \in \varphi^{-1}(J)$ .

Thus,

If  $\varphi : R \rightarrow S$  is a ring homomorphism and  $J$  is an ideal of  $S$ , then  $\varphi^{-1}(J)$  is an ideal of  $R$ .

□

**Problem 4.** Let  $\varphi : R \rightarrow S$  be a ring homomorphism, and let  $J$  be an ideal of  $R$ .

- (a) Prove that  $\varphi(J) = \{\varphi(j) \mid j \in J\}$  need not be an ideal of  $S$ .
- (b) Prove that if  $\varphi$  is surjective, then  $\varphi(J)$  is an ideal of  $S$ .
- (c) Prove that if  $\varphi$  is surjective, and  $I = \ker \varphi$ , then  $S \cong R/I$  and if we let  $\bar{J} \subseteq R/I$  be the image of  $\varphi(J)$  under this isomorphism, then

$$(R/I)/\bar{J} \cong R/(I+J).$$

*Proof.* (a) Consider the inclusion map  $\iota : \mathbb{Z} \hookrightarrow \mathbb{R}$  defined by  $\iota(n) = n \in \mathbb{R}$ ,  $\forall n \in \mathbb{Z}$ . This is a ring embedding by **Lemma 2**. Now look at  $2 \in 2\mathbb{Z}$ , which is a well-known ideal of  $\mathbb{Z}$ . Well,  $\pi \in \mathbb{R}$  but  $2\pi = \pi 2 \notin \iota(2\mathbb{Z}) = 2\mathbb{Z}$ . So then  $\iota(2\mathbb{Z})$  is not an ideal and we see that the ring homomorphic image of an ideal need not be an ideal of the codomain.

(b) For all  $\varphi(a), \varphi(b) \in \varphi(J)$  with preimages  $a, b \in J$ ,  $\varphi(a) - \varphi(b) = \varphi(a - b) \in \varphi(J)$  since  $a - b \in J$ . Therefore  $\varphi(J) \leq_+ S$ . Next, since  $\varphi$  is surjective, for any  $s \in S$ ,  $\exists r \in R$  such that  $s = \varphi(r)$ . So then  $s\varphi(a) = \varphi(r)\varphi(a) = \varphi(ra)$  and  $\varphi(a)s = \varphi(a)\varphi(r) = \varphi(ar)$  which must both belong to  $\varphi(J)$  since  $J \subseteq R$  is an ideal  $\implies ar, ra \in J$ . Therefore,  $\varphi(J)$  is an ideal of  $S$ . That is, a surjective ring homomorphic image of an ideal is in fact an ideal of the codomain.

(c)  $\varphi$  is surjective, so by the First Isomorphism Theorem  $R/\ker \varphi = R/I \cong \varphi(R) = S$  and so  $S \cong R/\ker \varphi = R/I$ . Let  $\psi : S \rightarrow R/I$  be this pullback isomorphism. That is,

$$\underbrace{r}_R \xrightarrow{\varphi} \underbrace{\varphi(r)=s}_S \xleftarrow{\psi} \underbrace{[r]_I = [r]_{\ker \varphi}}_{R/I=R/\ker \varphi} \quad (1)$$

$$\text{So: } \psi(s) = \psi(\varphi(r)) = [r]_I \in R/I, \text{ for each } s = \varphi(r) \in \varphi(R) = S \quad (2)$$

$$\implies \psi(\varphi(j)) = [j]_I \in R/I \text{ for each } j \in J \quad (3)$$

$$\text{Now let: } \bar{J} = \psi(\varphi(J)) = \{\psi(\varphi(j)) = [j]_I \mid j \in J\} \subseteq R/I \quad (4)$$

Recall the Third Isomorphism Theorem. For ideals  $A, B$  of a ring  $R$  where  $A \subseteq B$  is a subset:

$$\textcircled{1} B/A \text{ is an ideal of } R/A \text{ and } \textcircled{2} \frac{(R/A)}{(B/A)} \cong \frac{R}{B}.$$

By **Lemma 1**,  $I+J$  is an ideal of  $R$  since  $I, J$  are ideals of  $R$ . Also,  $I \subseteq I+J$  is a subset. Well, we can simply compute that  $(I+J)/I = \{(i+j)+I \mid (i+j) \in I+J\} = \{j+I = [j]_I \mid j \in J\} = \bar{J}$  by (4).

Therefore, since  $I, (I+J)$  are ideals of  $R$  with  $I \subseteq (I+J)$ , by the Third Isomorphism Theorem we have that

$$\textcircled{1} \bar{J} = (I+J)/I \text{ is an ideal of } (R/I) \text{ and } \textcircled{2} \frac{(R/I)}{\bar{J}} = \frac{(R/I)}{(I+J)/I} \cong \frac{R}{(I+J)}.$$

□

**Problem 5.** Let  $R$  be a commutative ring,  $a \in R$ , and let  $f_1(x), \dots, f_n(x) \in R[x]$ .

(a) Prove that  $R[x]/(x-a) \cong R$ .

(b) Prove the equality of ideals

$$(f_1(x), \dots, f_n(x), x-a) = (f_1(a), \dots, f_n(a), x-a).$$

(c) Prove the useful substitution trick

$$R[x]/(f_1(x), \dots, f_n(x), x-a) \cong R/(f_1(a), \dots, f_n(a)).$$

**Hint:** Use part (c) of the previous problem.

*Proof.* (a) Consider  $\varepsilon_a : R[x] \rightarrow R$  defined by  $\varepsilon_a(f(x)) = f(a)$ , an evaluation which was proven to be a *surjective homomorphism* in **Lemma 3**. We show that  $\ker \varepsilon_a = \langle x-a \rangle = \{f(x)(x-a) \mid f(x) \in R[x]\}$ .

( $\subseteq$ ) For any  $B(x) = \sum_{k=0}^n b_k x^k \in \ker \varepsilon_a$ ,  $B(a) = 0$ . Well,  $B(x) - B(a) = (\sum_{k=0}^n b_k x^k) - (\sum_{k=0}^n b_k a^k) = \sum_{k=0}^n b_k (x^k - a^k)$ . Then by **Lemma 4**,  $(x^k - a^k) = (x-a) \sum_{i+j=k-1} x^i y^j$  for each  $1 \leq k \leq n$ . Therefore,

$$B(x) - B(a) = \sum_{k=1}^n b_k (x^k - a^k) = \sum_{k=1}^n b_k ((x-a) \sum_{i+j=k-1} x^i a^j) = (x-a) \sum_{k=1}^n b_k (\sum_{i+j=k-1} x^i a^j)$$

and since in fact  $B(a) = 0$ , we have that  $B(x) = (x-a) \sum_{k=1}^n b_k (\sum_{i+j=k-1} x^i a^j) \in \langle x-a \rangle$  since  $\sum_{k=1}^n b_k (\sum_{i+j=k-1} x^i a^j)$  belongs to  $R[x]$ . So then  $\ker \varepsilon_a \subseteq \langle x-a \rangle$ .

( $\supseteq$ )  $\forall C(x) = f(x)(x-a) \in \langle x-a \rangle$  obviously  $C(a) = f(a)(a-a) = 0 \implies C(x) \in \ker \varepsilon_a$ . So then  $\langle x-a \rangle \subseteq \ker \varepsilon_a$ . Thus, by the **First Isomorphism Theorem**,

$$R[x]/\ker \varepsilon_a = R[x]/\langle x-a \rangle \cong \varepsilon_a(R[x]) = R.$$

(b) For each  $1 \leq m \leq n$ , there exist  $b_{m,0}, \dots, b_{m,d_m} \in R$  and  $d_m \in \mathbb{N}$  such that  $f_m(x) = \sum_{k=0}^{d_m} b_{m,k} x^k$ . Now we use a result obtained in (a):

$$f_k(x) - f_k(a) = (x-a) \sum_{k=1}^{d_m} b_{m,k} (\sum_{i+j=k-1} x^i a^j) \implies f_k(x) = f_k(a) + (x-a) \sum_{k=1}^{d_m} b_{m,k} (\sum_{i+j=k-1} x^i a^j)$$

So then

$$f_k(x) \in \langle f_k(a), x-a \rangle \subseteq \langle f_1(a), \dots, f_n(a), x-a \rangle, \forall 1 \leq k \leq n$$

and

$$f_k(a) \in \langle f_k(x), x-a \rangle \subseteq \langle f_1(x), \dots, f_n(x), x-a \rangle, \forall 1 \leq k \leq n$$

Therefore, all generators of  $\langle f_1(x), \dots, f_n(x), x-a \rangle$  belong to  $\langle f_1(a), \dots, f_n(a), x-a \rangle$  and vice versa. So the ideals are equal.

□

*Proof.* (c) In (a) we used the surjective homomorphism  $\varepsilon_a : R[x] \rightarrow R$  defined by  $\varepsilon_a(h(x)) = h(a)$  to show that  $R[x]/\langle x-a \rangle = R[x]/I \cong R$  by the **First Isomorphism Theorem** where  $I = \ker \varepsilon_a = \langle x-a \rangle$ . Much like in **Problem 4** let  $\psi : R \rightarrow R[x]/I$  be the pullback isomorphism and let  $J = \langle f_1(x), \dots, f_n(x), x-a \rangle \subseteq R[x]$ . Then:

$$\overbrace{r(x)}^{R[x]} \xrightarrow{\varepsilon_a} \overbrace{\varepsilon_a(r(x)) = r(a)}^R \xleftarrow{\psi} \overbrace{[r(x)]_I = [r(x)]_{\ker \varepsilon_a}}^{R[x]/I = R[x]/\ker \varepsilon_a} \quad (1)$$

$$\text{So: } \psi(r) = \psi(\varepsilon_a(r(x))) = [r(x)]_I \in R[x]/I, \text{ for each } r = \varepsilon_a(r(x)) \in \varepsilon_a(R[x]) = R \text{ (use } r(x) = r) \quad (2)$$

$$\implies \psi(\varepsilon_a(j(x))) = [j(x)]_I \in R[x]/I \text{ for each } j(x) \in J \quad (3)$$

$$\text{Now let: } \bar{J} = \psi(\varepsilon_a(J)) = \{\psi(\varepsilon_a(j)) = [j(x)]_I \mid j(x) \in J\} = J/I \subseteq R[x]/I \text{ *an ideal of } R[x]/I^* \quad (4)$$

Well,  $(I+J)/I = (\langle x-a \rangle + \langle f_1(x), \dots, f_n(x), x-a \rangle)/\langle x-a \rangle = \langle f_1(x), \dots, f_n(x), x-a \rangle/\langle x-a \rangle = J/I = \bar{J}$ . Since  $I, (I+J)$  are ideals of  $R[x]$  with  $I \subseteq (I+J)$ , by the **Third Isomorphism Theorem** we have that

$$\textcircled{1} \bar{J} = (I+J)/I \text{ is an ideal of } (R[x]/I) \text{ and } \textcircled{2} \frac{(R[x]/I)}{\bar{J}} = \frac{(R[x]/I)}{(I+J)/I} \cong \frac{R[x]}{I+J} = R[x]/J.$$

Let  $K = \langle f_1, \dots, f_n \rangle$  and  $\Phi : R[x]/I \rightarrow \frac{R}{\langle f_1(a), \dots, f_n(a) \rangle} = R/K$  defined via  $\Phi([h(x)]_I) = \psi^{-1}([h(x)]_I) + K$ .

Obviously this is well defined since  $\psi^{-1}$  is an isomorphism, and then for all  $[A(x)]_I, [B(x)]_I \in R[x]/I$ ,

$$\begin{aligned} \Phi([1]_I) &= \psi^{-1}([1_R]_I) + K = 1_R + K = 1_{R/K} \\ \Phi([A(x)]_I) + \Phi([B(x)]_I) &= \psi^{-1}([A(x)]_I) + \psi^{-1}([B(x)]_I) + K = \psi^{-1}([A(x)]_I + [B(x)]_I) + K \\ &= \Phi([A(x)]_I + [B(x)]_I) \\ \Phi([A(x)]_I) \Phi([B(x)]_I) &= (\psi^{-1}([A(x)]_I))(\psi^{-1}([B(x)]_I)) + K = \psi^{-1}([A(x)]_I [B(x)]_I) + K \\ &= \Phi([A(x)]_I [B(x)]_I) \\ \forall (r+K) \in R/K, \exists \psi(r) &= [r]_I \in R/I \text{ such that } \Phi(\psi(r)) = \psi^{-1}(\psi(r)) + K = r + K. \end{aligned}$$

So  $\Phi$  is an surjective ring homomorphism. We show  $\ker \Phi = \bar{J}$ . ( $\subseteq$ ) For any  $j(x) \in J = \langle f_1(a), \dots, f_n(a), x-a \rangle$  is of the form  $j(x) = j_0(a)(x-a) + \sum_{i=1}^n j_i(x)f_i(a)$  for some  $j_0(x), \dots, j_n(x) \in R[x]$ , and so  $j(a) = \sum_{i=1}^n j_i(a)f_i(a)$ . Therefore, for any  $[j(x)]_I \in \bar{J}$ ,  $\Phi([j(x)]_I) = \psi^{-1}([j(x)]_I) + K = j(a) + K = K \implies [j(x)]_I \in \ker \Phi \subseteq \ker \Phi$ .

( $\supseteq$ ) On the other hand, for any  $[r(x)]_I$ ,  $r(a) \in K$  and so  $r(a) = \sum_{i=1}^n r_i f_i(a)$  for some  $r_1, \dots, r_k \in R$  and then  $r(x) = r_0(x)(x-a) + \sum_{i=1}^n r_i(x)f_i(x)$  for some  $r_0(x), r_1(x), \dots, r_n(x) \in R[x]$  such that  $r_i(a) = r_i$  for each  $1 \leq i \leq n$ . Therefore,  $r(x) \in J$  and so  $[r(x)]_I \in \bar{J} \implies \ker \Phi \subseteq \bar{J}$ . So  $\ker \Phi = \bar{J}$ , and by the **First Isomorphism Theorem**  $\textcircled{3} \frac{R[x]/I}{\ker \Phi} = \frac{R[x]/I}{\bar{J}} \cong \Phi(R[x]/I) = R/J$ .

Thus, by  $\textcircled{2}$  and  $\textcircled{3}$ ,

$$\frac{R[x]}{\langle f_1(x), \dots, f_n(x), x-a \rangle} = R[x]/J \cong \frac{R[x]/I}{\bar{J}} \cong R/K = \frac{R}{\langle f_1(a), \dots, f_n(a) \rangle}$$

□



**Problem 6.** If  $\mathbb{k}$  is an algebraic closed field, then the only maximal ideals of  $\mathbb{k}[x]$  are of the form  $(x - a)$  where  $a \in \mathbb{k}$ . In this problem, we'll see that this is not true when  $\mathbb{k}$  is not algebraically closed.

- (a) Use the first isomorphism theorem to show that  $\mathbb{R}[x]/(x^2 + 1) \cong \mathbb{C}$ .
- (b) Prove that  $(x^2 + 1)$  is a maximal ideal of  $\mathbb{R}[x]$ .

*Proof.* (a) Let  $\varepsilon_i : \mathbb{R}[x] \rightarrow \mathbb{C}$  be defined via  $\varepsilon_i(f(x)) = f(i)$  for all  $f(x) \in \mathbb{R}[x]$ . We show this is a surjective ring homomorphism. For any  $A(x), B(x) \in \mathbb{R}[x]$ ,

$$\begin{aligned}\varepsilon_i(1(x)) &= 1(a) = 1_{\mathbb{R}} = 1_{\mathbb{C}} \\ \varepsilon_i(A(x) + B(x)) &= A(i) + B(i) = (A + B)(i) \varepsilon_i((A + B)(x)) = \varepsilon_i(A(x) + B(x)) \\ \varepsilon_i(A(x))\varepsilon_i(B(x)) &= A(i)B(i) = (AB)(i) \varepsilon_i((AB)(x)) = \varepsilon_i(A(x)B(x)) \\ \forall (\alpha + \beta i) \in \mathbb{C}, \varepsilon_i(\alpha + \beta x) &= \alpha + \beta i\end{aligned}$$

Next, we show that  $\ker \varepsilon_i = \langle x^2 + 1 \rangle$ . ( $\subseteq$ ) Consider any  $C(x) \in \ker \varepsilon_i$ . By **Lemma 5** we have the division algorithm for polynomials over fields. So  $C(x) = q(x)(x^2 + 1) + r(x)$  where  $r(x) = 0$  or  $\deg(r(x)) < 2 = \deg(x^2 + 1)$ . Well,  $0 = C(i) = q(i)(i^2 + 1) + r(i) \implies r(i) = 0$  and so it can't be linear otherwise  $r(i)$  wouldn't vanish;  $r(x) = ax + b$  is linear implies  $a \neq 0$  and then  $r(i) = ai + b \neq 0$  since  $\mathbb{C}$  is an integral domain. So then by **Lemma 5** in fact  $r(x)$  must be zero, and so  $x^2 + 1$  divides  $C(x) \implies C(x) \in \langle x^2 + 1 \rangle$ . ( $\supseteq$ ) Trivially, any  $f(x) = r(x)(x^2 + 1)$  belongs to  $\ker \varepsilon_i$  since then  $f(i) = r(i)(i^2 - 1) = 0$ .

So then  $\ker \varepsilon_i = \langle x^2 + 1 \rangle$  and by the **First Isomorphism Theorem**

$$\mathbb{R}[x]/\langle x^2 + 1 \rangle = \mathbb{R}[x]/\ker \varepsilon_i \cong \mathbb{C}.$$

(b)  $\mathbb{R}[x]/\langle x^2 + 1 \rangle \cong \mathbb{C}$ , so  $\mathbb{R}[x]/\langle x^2 + 1 \rangle$  is a field and then by **Lemma 6**,  $\langle x^2 + 1 \rangle$  must be maximal. □

**Problem 7.** Let  $R$  be a commutative ring with  $0 \neq 1$ . In this problem, we will prove that every proper ideal of  $R$  is contained in some maximal ideal.

- (a) Look up Zorn's Lemma and record it here.
- (b) Define  $S = \{J \mid J \text{ is a proper ideal of } R \text{ and } J \supseteq I\}$ . Explain why  $S$  is a partially ordered set (what is the ordering?).
- (c) Given a chain  $C$  in  $S$ , prove that  $\bigcup_{J \in C} J$  is an ideal of  $R$  (you will use that  $C$  is totally ordered), and further that this ideal is in the set  $S$ .
- (d) Conclude using Zorn's Lemma that  $S$  has a maximal element.

(a) Zorn's Lemma is **Lemma 7** at the top of this document. We state it again for convenience.

**Lemma 7** (Zorn's Lemma). *Let  $(P, \leq)$  be a partially ordered set. If every chain in  $P$  has an upper bound in  $P$ , then  $P$  contains a maximal element.*

The proof for this problem is on the next page.

*Proof.* (b) (1)  $A \supseteq A$ ,  $\forall A \in S$ . (2)  $A \supseteq B$  and  $A \neq B \implies A \supset B \implies B \not\supseteq A$  (3)  $A \supseteq B$  and  $B \supseteq C \implies A \supseteq C$ . So  $(J, \supseteq)$  is a poset.

(c) Let  $C$  be a chain in  $S$  and consider  $C_\cap = \bigcap_{J \in C} J$ . We show this is an ideal of  $R$ . For any  $a, b \in C_\cap$ ,  $a, b \in J, \forall J \in C$ . So then  $a - b \in J, \forall J \in C$  since they're all additive subgroups of  $R$  and so  $C_\cap \leq_+ R$ .

Next, for any  $r \in R$  and any  $i \in I$ ,  $ri = ir \in J, \forall J \in C$  and so  $ri = ir \in C_\cap$ . Therefore,  $C_\cap$  is an ideal of  $R$ . Since  $C_\cap \subseteq J \subset R, \forall J \in C$ ,  $C_\cap$  must be a proper ideal of  $R$ . Lastly, by definition  $I$  is contained in every  $J \in C \subseteq S$ , and so of course  $I$  must be contained in their intersection  $C_\cap$ . So  $C_\cap \in S$ .

(d) Since  $J \supseteq I, \forall J \in C_\cap$  □

**Problem 8.** Let  $\mathbb{k}$  be a field. In this problem, we will prove that the only maximal ideal of  $\mathbb{k}[[x]]$  is  $(x)$ , which makes  $\mathbb{k}[[x]]$  a local ring.

- (a) Explain why  $(x) = \{f \in \mathbb{k}[[x]] \mid f \text{ has no constant term}\}$ .
- (b) Compute  $\mathbb{k}[[x]]/(x)$ , and then explain why  $(x)$  is a maximal ideal.
- (c) You may freely use the following result from the optional hint last week:  $f \in \mathbb{k}[[x]]$  is a unit if and only if  $f$  has a nonzero constant term. Use Proposition 1.41 to show that the only maximal ideal of  $\mathbb{k}[[x]]$  is  $(x)$ .

**Problem 9.** Let  $d$  be an integer which is not the square of an integer, and consider

$$\mathbb{Q}(\sqrt{d}) = \{a + b\sqrt{d} \mid a, b \in \mathbb{Q}\}.$$

- (a) Prove that  $\mathbb{Q}(\sqrt{d})$  is a subring of  $\mathbb{C}$ .
- (b) Define a function  $N : \mathbb{Q}(\sqrt{d}) \rightarrow \mathbb{Q}$  by  $N(a + b\sqrt{d}) = a^2 + b^2d$ . Prove that  $N(zw) = N(z)N(w)$  and that  $N(z) \neq 0$  if  $z \neq 0$ . This function is often called the norm.
- (c) Prove that  $\mathbb{Q}(\sqrt{d})$  is a field and is the smallest subfield of  $\mathbb{C}$  containing both  $\mathbb{Q}$  and  $\sqrt{d}$  (use  $N$ ).
- (d) Prove that  $\mathbb{Q}(\sqrt{d}) \cong \mathbb{Q}[t]/(t^2 - d)$ .