

Def. A constant solution (x_n) , $x_n \equiv x^*$ of the difference equation

$$x_{n+k} = f(x_{n+k-1}, \dots, x_n)$$

is called an *equilibrium solution*.

A point x^* is called an *equilibrium point* for the difference equation and it is a solution of the equation

$$x^* = f(x^*, \dots, x^*)$$

Linear case

Consider the k-order difference equation

$$x_{n+k} + p_1 x_{n+k-1} + p_2 x_{n+k-2} + \cdots + p_k x_n = 0$$

$$\mathbf{x}^* = \mathbf{0}$$

the characteristic equation is:

$$q^k + p_1 q^{k-1} + p_2 q^{k-2} + \cdots + p_k = 0$$

Theorem.

i) $x^* = 0$ is asymptotically stable if and only if $|q| < 1$ for all roots of characteristic equation.

ii) $x^* = 0$ is locally stable if and only if $|q| \leq 1$ for all roots of characteristic equation, $|q| = 1$ holds for simple roots.

iii) $\mathbf{x}^* = 0$ is unstable if and only if there exists a root q of the characteristic equation for which $|q| > 1$ or $|q| = 1$ and q is not a simple root.

```
> restart;
```

```
> eqd:=6*x(n+2)+x(n+1)-2*x(n)=0;
```

$$eqd := 6\,x(n+2) + x(n+1) - 2\,x(n) = 0$$

```
> ech:=6*q^2+q-2=0;
```

$$ech := 6\,q^2 + q - 2 = 0$$

```
> solve(ech,q);
```

$$\frac{1}{2}, -\frac{2}{3}$$

```
> rsolve(eqd,x(n));
```

$$\frac{1}{3} \left(\frac{9}{7} x(0) - \frac{18}{7} x(1) \right) \left(-\frac{2}{3} \right)^n - \frac{1}{2} \left(-\frac{8}{7} x(0) - \frac{12}{7} x(1) \right) \left(\frac{1}{2} \right)^n$$

```
> x[0]:=20;x[1]:=40;N:=50;
```

$$x_0 := 20$$

$$x_1 := 40$$


$$N := 50$$

```
> for i from 0 to N-2 do
```

```

    x[i+2]:=1/6*(-x[i+1]+2*x[i])
end do:
> plot([[n,x[n]]$n=0..N],style=point,symbol=circle);

```



```

> restart;
> eqd:=x(n+2)-6*x(n+1)+18*x(n)=0;
      eqd:=x(n+2)-6x(n+1)+18x(n)=0
> ech:=q^2-6*q+18=0;
      ech:=q^2-6q+18=0
> qq:=solve(ech,q);
      qq:=3+3I,3-3I
> abs(qq[1]);
      3√2
> sol:=rsolve(eqd,x(n));
sol:=⎛⎜⎝112-112I⎞⎟⎠(6Ix(0)-Ix(1)+x(1))(3+3I)n+⎛⎜⎝-112-112I⎞⎟⎠(6Ix(0)
      -Ix(1)-x(1))(3-3I)n
> evalc(sol);

```

$$x(0) e^{\frac{1}{2} n \ln(18)} \cos\left(\frac{1}{4} n \pi\right) - \left(-\frac{1}{6} x(1) + \frac{1}{2} x(0)\right) e^{\frac{1}{2} n \ln(18)} \sin\left(\frac{1}{4} n \pi\right) + \left(\frac{1}{6} x(1) - \frac{1}{2} x(0)\right) e^{\frac{1}{2} n \ln(18)} \sin\left(\frac{1}{4} n \pi\right) + I \left(\left(-\frac{1}{6} x(1) + \frac{1}{2} x(0)\right) e^{\frac{1}{2} n \ln(18)} \cos\left(\frac{1}{4} n \pi\right) + \left(\frac{1}{6} x(1) - \frac{1}{2} x(0)\right) e^{\frac{1}{2} n \ln(18)} \cos\left(\frac{1}{4} n \pi\right) \right)$$

```
> simplify(evalc(sol));
```

$$\frac{1}{3} e^{\frac{1}{2} n (\ln(2) + 2 \ln(3))} \left(3 x(0) \cos\left(\frac{1}{4} n \pi\right) - 3 \sin\left(\frac{1}{4} n \pi\right) x(0) + \sin\left(\frac{1}{4} n \pi\right) x(1) \right)$$

```
> simplify(evalc(sol), exp);
```

$$x(0) 18^{\frac{1}{2} n} \cos\left(\frac{1}{4} n \pi\right) - \left(-\frac{1}{6} x(1) + \frac{1}{2} x(0)\right) 18^{\frac{1}{2} n} \sin\left(\frac{1}{4} n \pi\right) + \left(\frac{1}{6} x(1) - \frac{1}{2} x(0)\right) 18^{\frac{1}{2} n} \sin\left(\frac{1}{4} n \pi\right) + I \left(\left(-\frac{1}{6} x(1) + \frac{1}{2} x(0)\right) 18^{\frac{1}{2} n} \cos\left(\frac{1}{4} n \pi\right) + \left(\frac{1}{6} x(1) - \frac{1}{2} x(0)\right) 18^{\frac{1}{2} n} \cos\left(\frac{1}{4} n \pi\right) \right)$$

```
> x[0]:=0.1;x[1]:=-0.3;N:=17;
```

$$x_0 := 0.1$$

$$x_1 := -0.3$$


$$N := 17$$

```
> for i from 0 to N-2 do
```

```
    x[i+2]:=6*x[i+1]-18*x[i]
```

```
end do;
```

```
> plot([[n,x[n]]$n=0..N],style=point,symbol=circle);
```



Nonlinear case

$$x_{n+k} = f(x_{n+k-1}, \dots, x_n)$$

$$f = f(u_1, \dots, u_k)$$

$$p_i = \frac{\partial}{\partial u_i} f(x^*, \dots, x^*)$$

linearized equation in (x^*, x^*, \dots, x^*)

$$y_{n+k} = p_1 y_{n+k-1} + p_2 y_{n+k-2} + \dots + p_k y_n$$

characteristic equation corresponding to the linearized equation is

$$q^k = p_1 q^{k-1} + p_2 q^{k-2} + \dots + p_k$$

Theorem (The Linearized Stability Theorem).

Suppose that f is continuously differentiable on an open neighborhood G from R^{k+1} of (x^*, x^*, \dots, x^*) , where x^* is an equilibrium point of the nonlinear difference equation. Then the following statements are true:

- (i) If all the characteristic roots of the linearized characteristic equation lie inside the unit disk in the complex plane ($|q| < 1$), then the equilibrium point x^* is locally asymptotically stable.
- (ii) If at least one characteristic root of the linearized characteristic equation is outside the unit disk ($|q| > 1$) in the complex plane, the equilibrium point x^* is unstable.
- (iii) If one characteristic root is on the unit disk and all the other characteristic roots are either inside or on the unit disk, then the equilibrium point x^* may be stable, unstable, or asymptotically stable.

The Schur-Cohn Criterion

(a) for the equation the equation

$$q^2 = p_1 q + p_2$$

we have:

$$|q_{1,2}| < 1 \text{ if and only if } |p_1| < 1 - p_2 < 2$$

(b) for the equation the equation

$$q^3 = p_1 q^2 + p_2 q + p_3$$

we have:

$$|q_{1,2,3}| < 1 \text{ if and only if } |p_3 + p_1| < 1 - p_2 \text{ and } |p_1 p_3 + p_2| < 1 - p_3^2$$

Example

The second order Beverton-Holt model

$$x_{n+1} = \frac{r K (\alpha x_n + \beta x_{n-1})}{K + (r-1) x_{n-1}}$$

has been used to model populations of bottom-feeding fish, where $\alpha + \beta = 1$, $r > 0$, $K > 0$. These species have very high fertility rates and very low survivorship to adulthood. Furthermore, recruitment is essentially unaffected by fishing. In this model, the future generation x_{n+1} depends not only on the present generation x_n but also on the previous generation x_{n-1} .

```
[ > restart;
```

$$f := (u, v) \rightarrow \frac{r K (\alpha u + \beta v)}{K + (r-1) v}$$

```
> f := (u,v) -> r*K*(alpha*u+beta*v)/(K+(r-1)*v);
```

$$eq := x = \frac{r K (\alpha x + \beta x)}{K + (r-1) x}$$

```
> eq:=x=f(x,x);
```

$$eqp := 0, \frac{K (\alpha r + \beta r - 1)}{r - 1}$$

```
> eqp:=solve(eq,x);
```

```

> x1:=eqp[1];x2:=simplify(subs(beta=1-alpha,eqp[2]));
      x1 := 0
      x2 := K
[
> p1:=D[1](f)(0,0);
      p1 := α r
> p2:=D[2](f)(x1,x1);
      p2 := β r
> lineq:=y(n+1)=p1*y(n)+p2*y(n-1);
      lineq := y(n + 1) = α r y(n) + β r y(n - 1)
> chareq:=q^2=p1*q+p2;
      chareq := q^2 = α q r + β r
> rr:=solve(chareq,q);
      rr := 1/2 α r + 1/2 √(α^2 r^2 + 4 β r), 1/2 α r - 1/2 √(α^2 r^2 + 4 β r)
> abs(subs(beta=1-alpha,rr[1]));
      |1/2 α r + 1/2 √(α^2 r^2 + 4 (1 - α) r)|
> abs(p1)<1-p2;
      |α r| < -β r + 1
> 1-p2<2
      -β r < 1

```

Analysing the roots $x_1^*=0$ is is locally asymptotically stable if $0 < r < 1$.

```

[
> p1:=D[1](f)(x2,x2);
      p1 := r K α / (K + (r - 1) K)
> p1:=simplify(p1);
      p1 := α
> p2:=D[2](f)(x2,x2);
      p2 := r K β / (K + (r - 1) K) - r K (K α + K β) (r - 1) / (K + (r - 1) K)^2
> p2:=simplify(p2);
      p2 := - (α r - α - β) / r
> p2:=subs(beta=1-alpha,p2);
      p2 := - (α r - 1) / r
> lineq:=y(n+1)=p1*y(n)+p2*y(n-1);
      lineq := y(n + 1) = α y(n) - (α r - 1) y(n - 1) / r

```

```
> chareq:=q^2=p1*q+p2;
```

$$chareq := q^2 = \alpha q - \frac{\alpha r - 1}{r}$$

```
> solve(chareq,q);
```

$$\frac{1}{2} \frac{\alpha r + \sqrt{\alpha^2 r^2 - 4 \alpha r^2 + 4 r}}{r}, -\frac{1}{2} \frac{-\alpha r + \sqrt{\alpha^2 r^2 - 4 \alpha r^2 + 4 r}}{r}$$

```
> abs(p1)<1-p2;
```

$$|\alpha| < 1 + \frac{\alpha r - 1}{r}$$

```
> 1-p2<2
```

$$\frac{\alpha r - 1}{r} < 1$$

Analysing the roots we get that the equilibrium $x_2^* = K$ is locally asymptotically stable if $r > 1$.

Remark. If $r > 1$ then $x_2^* = K$ is globally asymptotically stable

Numerical simulations

```
> K:=100;r:=0.8;alpha:=0.5;beta:=1-alpha;N:=100;
```

$$K := 100$$

$$r := 0.8$$

$$\alpha := 0.5$$

$$\beta := 0.5$$

$$N := 100$$

```
> f(u,v);
```

$$\frac{80.0 (0.5 u + 0.5 v)}{100 - 0.2 v}$$

```
> x[0]:=20;x[1]:=40;
```

$$x_0 := 20$$


$$x_1 := 40$$

```
> for i from 0 to N-2 do
```

```
    x[i+2]:=f(x[i+1],x[i])
```

```
end do;
```

```
> plot([[n,x[n]]$n=0..N],style=point,symbol=circle);
```



```
> K:=100;r:=1.2;alpha:=0.5;beta:=1-alpha;N:=100;
```

```
K:= 100
```

```
r:= 1.2
```

```
 $\alpha$ := 0.5
```

```
 $\beta$ := 0.5
```



```

=
> f(u,v);
=

$$\frac{120.0 (0.5 u + 0.5 v)}{100 + 0.2 v}$$


=
> x[0]:=10;x[1]:=20;
=

$$x_0 := 10$$


$$x_1 := 20$$

=
> for i from 0 to N-2 do
    x[i+2]:=f(x[i+1],x[i])
end do:
=
> plot([[n,x[n]]$n=0..N],[[n,K]$n=0..N],style=[point,point],
symbol=[circle,cross],color=[red,green]);

```



```
> restart;
```

Autonomous Systems of Difference equations

Linear case

$$x_{n+1} = A x_n$$

the solution

$$x_n = A^n x_0$$

Theorem. The following statements hold:


- (i) The zero solution is locally stable if and only if $|\lambda| \leq 1$ and the eigenvalues of unit modulus are simple.
- (ii) The zero solution is asymptotically stable if and only if $|\lambda| < 1$ for all eigenvalues of A.

Example

$$\begin{cases} x_{n+1} = \frac{1}{2}x_n + y_n \\ y_{n+1} = \frac{1}{3}x_n + \frac{1}{8}y_n \end{cases}$$

```
> restart;
> with(linalg):
> A:=matrix([[1/2,1],[1/3,1/8]]);
A :=  $\begin{bmatrix} \frac{1}{2} & 1 \\ \frac{1}{3} & \frac{1}{8} \end{bmatrix}$ 
> eigenvals(A);
 $\frac{5}{16} + \frac{1}{48} \sqrt{849}, \frac{5}{16} - \frac{1}{48} \sqrt{849}$ 
> evalf(%);
0.9195334284, -0.2945334284
both eigenvalues  $|\lambda_{1,2}| < 1$ , so (0,0) is locally asymptotically stable
> N:=100;x[0]:=50;y[0]:=-20;
N:=100
x0:=50
y0:=-20
> for i from 0 to N-1 do
    x[i+1]:=1/2*x[i]+y[i];
    y[i+1]:=1/3*x[i]+1/8*y[i];
end do;
> plot([[n,x[n]]$n=0..N],[[n,y[n]]$n=0..N],style=[point,
```

```
point],symbol=[circle, cross], color=[red,blue]);
```



Nonlinear case

$$x_{n+1} = f(x_n)$$

$x^* = f(x^*)$ equilibrium point

linearized system

$$y_{n+1} = \text{Jacobian}(f)(x^*) y_n$$

Theorem. The following statements hold:

- (i) The x^* is locally asymptotically stable if and only if $|\lambda| < 1$ for all eigenvalues of the $\text{Jacobian}(f)(x^*)$.
- (ii) The x^* is unstable if there exists an eigenvalue of the $\text{Jacobian}(f)(x^*)$ such that $|\lambda| > 1$.

Example

$$\begin{cases} x_{n+1} = y_n \\ y_{n+1} = \frac{\alpha y_n}{1 + \beta x_n} \end{cases}$$

```
> restart;with(linalg):
```

```
> f1:=(x,y)->y;
```

$$f1 := (x, y) \rightarrow y$$

```
> f2:=(x,y)->alpha*y/(1+beta*x);
```

$$f2 := (x, y) \rightarrow \frac{\alpha y}{1 + \beta x}$$

```
> eqp:=solve({f1(x,y)=x,f2(x,y)=y},{x,y});
```

$$eqp := \{x=0, y=0\}, \left\{x = \frac{\alpha - 1}{\beta}, y = \frac{\alpha - 1}{\beta}\right\}$$

```
> J:=jacobian([f1(x,y),f2(x,y)],[x,y]);
```

$$J := \begin{bmatrix} 0 & 1 \\ -\frac{\alpha y \beta}{(\beta x + 1)^2} & \frac{\alpha}{\beta x + 1} \end{bmatrix}$$

```
> A1:=subs(x=0,y=0,evalm(J));
```

$$A1 := \begin{bmatrix} 0 & 1 \\ 0 & \alpha \end{bmatrix}$$

```
> eigenvals(A1);
```

$$0, \alpha$$

```
> A2:=subs(x=(alpha-1)/beta,y=(alpha-1)/beta,evalm(J));
```

$$A2 := \begin{bmatrix} 0 & 1 \\ -\frac{\alpha - 1}{\alpha} & 1 \end{bmatrix}$$

```
> eigenvals(A2);
```


$$\frac{1}{2} \frac{\alpha + \sqrt{-3 \alpha^2 + 4 \alpha}}{\alpha}, -\frac{1}{2} \frac{-\alpha + \sqrt{-3 \alpha^2 + 4 \alpha}}{\alpha}$$

```
> charpoly(A2,lambda);
```


$$\frac{\alpha \lambda^2 - \alpha \lambda + \alpha - 1}{\alpha}$$

Numerical simulation

```
> x[0]:=10;y[0]:=20;
                                      $x_0 := 10$ 
                                      $y_0 := 20$ 
=
> alpha:=0.8;beta:=1;N:=100;
                                      $\alpha := 0.8$ 
                                      $\beta := 1$ 
                                      $N := 100$ 
=
> for i from 0 to N-1 do
    x[i+1]:=f1(x[i],y[i]);
    y[i+1]:=f2(x[i],y[i]);
end do;
=
> plot([[n,x[n]]$n=0..N],style=point,symbol=circle);
```



```
> plot([n,y[n]]$n=0..N,style=point,symbol=circle);
```



```
> x[0]:=10;y[0]:=15;
```

$x_0 := 10$


$y_0 := 15$

```
> alpha:=3;beta:=0.3;N:=100;
```



```

 $\alpha := 3$ 
 $\beta := 0.3$ 
 $N := 100$ 
=
>  $xx := (\alpha - 1) / \beta;$ 
=
 $xx := 6.666666667$ 
> for i from 0 to N-1 do
     $x[i+1] := f1(x[i], y[i]);$ 
     $y[i+1] := f2(x[i], y[i]);$ 
end do:
=
> plot([[n, x[n]]$n=0..N], [[n, xx]$n=0..N], style=[point, point],
    symbol=[circle, cross], color=[red, blue]);
```



```
> plot([[n,y[n]]$n=0..N],[[n,xx]$n=0..N],style=[point,point],  
       symbol=[circle,cross],color=[red,blue]);
```

