

A Matlab[®] toolbox for calculating spring indices from daily meteorological data

Toby R. Ault, Raul Zurita-Milla, and Mark D. Schwartz

Corresponding author email: toby.ault@cornell.edu

May 15, 2014

Contents

1	Introduction	1
1.1	Historical background on spring indices	2
2	Spring Indices Model Formulations	3
2.1	Assumptions	3
2.2	Methods	4
2.3	Model predictors and parameters	5
2.4	Implementation in Matlab [®]	7
3	Spring Index Applications	8
3.1	Data	8
3.2	Example 1 - SI and predictor variables	8
3.3	Example 2 - SI calculated from a US weather station	9
3.4	Example 3 - SI compared with phenological data	9
3.5	Example 4 - SI means and trends in the coterminous US	9
4	Discussion	10
5	Appendix A: Additional resources	11

Abstract

Metrics to track seasonal transitions are needed for a wide variety of ecological and climatological applications. Here a MATLAB toolkit for calculating spring indices (Schwartz et al., 2006) is documented. The spring indices have been widely used in earlier studies to model phenological variability and change through time across a wide range of spatial scales. These indices require only daily minimum and maximum temperature observations (e.g., from meteorological records) as input along with latitude, and produce a day of year (DOY) value corresponding to the simulated average timing of first leaf and first bloom events among three plant cultivars. Core functions to calculate the spring indices require no external dependencies, and data for running several illustrative test cases are included. Instructions and routines for conducting more sophisticated spring index monitoring and modeling are also supplied and documented.

1 Introduction

Understanding the impacts of global climate change on ecosystems requires systematic development and use of metrics to track seasonal transitions clearly and consistently across broad spatial scales and through time. Phenology – the study of seasonally recurrent biological events and their relation to climate – can help guide the development of such metrics because many phenological events are sensitive to variations of weather and climate (Lieth, 1974). Accordingly, worldwide efforts to collect, monitor, and synthesize phenological data offer a unique perspective on how global warming is affecting organisms across a wide range of spatial scales (van Vliet et al., 2003; IPCC, 2007; Janetos et al., 2012; Schwartz et al., 2012). Although historical phenological observations were collected by individual observers at relatively small scales, developments in information and communication technologies are eroding these limitations. Today phenological data collection is “crowd-sourced” to the general public (van Vliet et al., 2003; Schwartz et al., 2012), and observed from space (e.g., White et al., 2009; Zurita-Milla et al., 2013). As these networks of volunteers and remote sensing operations grow in scale and scope, interpretation of the sheer volume of data will be greatly improved if it can be readily compared with clearly defined markers of seasonal evolution.

The spring indices (SI) “suite of metrics” (Schwartz et al., 2006) are a prominent example of phenological indices that have been widely used to characterize the transition from winter into spring in a clear and consistent way across continental scales and through time. These metrics include indices of “first leaf” and “first bloom,” as well as “last freeze” and the “damage index,” which records the anomalous amount of time from the appearance of foliage to the

last freeze (e.g., Schwartz and Marotz, 1986, 1988; Schwartz, 1999; Schwartz and Reiter, 2000; Schwartz et al., 2006, 2013). Here, we present a Matlab[©] toolbox called `ml_si` that calculates the most recent version of SI, which were originally developed by M. D. Schwartz almost 25 year ago. This toolbox consists of six core functions to calculate *spring indices* from daily values of temperature minima (T_{min}) and maxima (T_{max}) from any arbitrary location in the Northern Hemisphere. The toolbox is distributed as a single zipped tarball through the “Matlab[©] Central” file exchange (<http://www.mathworks.com/matlabcentral/fileexchange/>), and also as supplementary material to this document. It will be updated periodically as new versions are released to extend its functionality.

1.1 Historical background on spring indices

In the 1950s Joe Caprio of Montana State University initiated a program to monitor the phenological responses of Lilac (*Syringa vulgaris*) with the aid of a large network of volunteers (e.g., Schwartz and Reiter, 2000; Cayan et al., 2001; Schwartz et al., 2012). At its peak, the network consisted of about a thousand citizen scientists making observations across 12 western states. The project lasted more than 40 years, and even today, certain elements of these early efforts continue to be supported by the USA National Phenological Network (USA-NPN: <http://www.usanpn.org/>), which also coordinates large scale monitoring projects of a much wider diversity of plant and animal species (Schwartz et al., 2012).

Similar observations to those in the Caprio network were also made in the Eastern US, and this continental-scale coverage of phenological data over several decades allowed Schwartz (1985) to develop models of Lilac (*Syringa*) and Honeysuckle (*Lonicera*) phenology using observed meteorological variables (minimum and maximum temperature, in this case). These models were optimized for use at continental-scales, meaning that they reliably indicate the same event (e.g., first leaf) whether they are calculated from meteorological data in Arizona or British Columbia. Such consistency has allowed for the spring indices to be used for a variety of applications, including characterizing the impacts of global change (Schwartz et al., 2006), interpreting satellite observations of land-surface phenology (White et al., 2009), and understanding interannual variations in spring onset and its impacts (Ault et al., 2011; McCabe et al., 2011; Schwartz et al., 2013; Ault et al., 2013).

2 Spring Indices Model Formulations

2.1 Assumptions

Translating daily temperature fluctuations into indices of spring onset is motivated by two features of this seasonal transition. First, variations in the timing of many phenological events vary considerably on interannual timescales, meaning that the day of year when spring foliage first appears on a given plant typically differs from one year to the next. Second, the start of the growing season for many species occurs “in concert” (i.e., coherently) across large spatial scales. Taken together, these two attributes of spring onset motivate a climate-oriented (as opposed to plant-oriented) approach to characterizing spring onset: interannual variations by themselves could be explained by assuming that the timing of plant events is stochastic, while large-scale correlations could arise from deterministic elements of the environment (e.g., the length of the day). From a climatic perspective, however, these two features are interpreted as the hallmark of weather variations on interannual timescales, which are themselves correlated across large spatial scales. This “climate-centric” perspective on spring onset is the starting point for the spring indices.

To translate meteorological data (T_{min} and T_{max}) into indices that track the timing of spring in a consistent way through time and across large spatial scales, the calculation of the SI makes the following assumptions, which are valid for most temperate regions:

1. Phenological observations from a few key indicator species can be used to characterize spring onset in a way that is relevant to a wide range of ecological and climatic transitions at the start of spring.
2. Where moisture availability is not a factor, plant phenological responses reflect the accumulated effects of temperature fluctuations in conjunction with increasing day length.
3. Near surface temperatures in temperate regions do not increase monotonically during springtime, but instead fluctuate as a consequence of large-scale (e.g., ~ 1000 km) physical processes. These phenomena are synoptic-scale weather disturbances that drive multi-day warm episodes by advecting warm air masses in from the south.
4. Synoptic-scale weather events can be detected from meteorological observations of daily T_{min} and T_{max} .

2.2 Methods

The calculation of the SI is underpinned by a multivariate (stepwise) regression that relates climate variables to plant responses. That is, for a given plan (p) and a given event, the following is solved for the m coefficients $A_{1,\dots,m}$:

$$\frac{C}{\text{date}} = A_0 + A_1 \left(\frac{X_1}{\text{date}} \right) + \dots + A_m \left(\frac{X_m}{\text{date}} \right) \quad (1)$$

where C is an arbitrary value (typically 1000), A_0 is a regression constant, and $X_{1,\dots,m}$ are various predictor variables at the time of the phenological event. The term “date” is the day of the phenological event of interest, rendered as “day of year” (i.e., Jan. 1st = 1). It appears in the denominator so that “late” dates (large values) do not unduly influence the value of the regression coefficients.

Once the $A_{1,\dots,m}$ coefficients have been optimized, equation (1) can be used operationally by multiplying through by the actual date and calculating the quantity $C(t)$ as it evolves through time as a function of each time-varying predictor ($X_{1,2,\dots,M}(t)$):

$$\hat{C}(t) = t(A_0) + A_1 X_1(t) + \dots + A_m X_m(t), \quad 0 \leq t \leq T. \quad (2)$$

The date of a given phenological event for a given plant is then predicted if $\hat{C}(t)$ is greater than C . In this way, the SI resemble traditional heat accumulation models because $\hat{C}(t)$ grows with time until the prediction date of a given event is achieved. In order to express multiple equations in the form of (2), we combine the accumulation models in matrix notation:

$$\begin{bmatrix} C_{1,1} & \cdots & C_{1,t} \\ \vdots & \ddots & \vdots \\ C_{p,1} & \cdots & C_{p,t} \end{bmatrix} = \begin{bmatrix} A_{1,1} & \cdots & A_{1,m} \\ \vdots & \ddots & \vdots \\ A_{1,p} & \cdots & A_{p,m} \end{bmatrix} \begin{bmatrix} X_{1,1} & \cdots & X_{1,t} \\ \vdots & \ddots & \vdots \\ X_{1,m} & \cdots & X_{m,t} \end{bmatrix}, \quad (3)$$

which simply defines a matrix C with rows ($1 \leq p \leq P$) that are outputs from each of the accumulation models for each event/plant target, and columns ($t = 1, 2, \dots, T$) that are the values of those accumulation models at each time (days in this case). The next step is to identify the date on which each of the $p = 1, 2, \dots, N$ accumulation models is greater than C , and assign that (raw) date to the vector S'_p :

$$S'_p = \begin{cases} t & \text{if } \hat{C}_{pt} \geq C \\ \text{undefined} & \text{otherwise.} \end{cases}, \text{ for } p = 1, 2, \dots, N; \text{ and } t = 1, 2, \dots, T. \quad (4)$$

The underlying equation of the leaf index can now be written as:

$$\hat{\mathbf{C}} = \mathbf{A}\mathbf{X}, \quad (5)$$

$$S'_p = \begin{cases} t & \text{if } \hat{C}_{pt} \geq C \\ \text{undefined} & \text{otherwise} \end{cases}, \quad (6)$$

where $\hat{\mathbf{C}}$ is the time evolving matrix of accumulations, \mathbf{A} is the matrix of predictor coefficients for each plant for a given event, and \mathbf{X} is the time-evolving matrix of predictors. As in equation (4), $\mathbf{S}'_{\mathbf{p}}$ is a vector of raw dates for each event.

Applying equations (5) through (4) is now trivial for any arbitrary set of years ($n = 1, 2, \dots, N$), points in space ($j = 1, 2, \dots, J$), and events (e.g., the emergence of first leaf or first bloom).

2.3 Model predictors and parameters

Here we define the m predictors comprising \mathbf{X} and the coefficients \mathbf{A} for each plant type and event. Stepwise regression was used to optimize these parameters (\mathbf{X} and \mathbf{A}) for continental-scale phenological monitoring, and as it turned out not all predictors were required to model all phenological events/plants. For clarity, we will use the subscripts “*lf*” and “*bl*” to refer to the equations for the first leaf and first bloom indices, respectively.

The following table describes the predictors used by, e.g., Schwartz et al. (2006) for the leaf index (\mathbf{X}_{lf}):

Predictor abbreviation	Description
MDS0	Days since Jan 1
SYNOP	Number of high-energy synoptic events
DD57	5-7 day degree-hour accumulations
DDE2	2-3 day degree-hour accumulations

The values of SYNOP, DD57 and DDE2 are defined as follows (again, for the *leaf index*):

$$\text{SYNOP}(t) = \begin{cases} 0 & \text{if } t = 1, \\ \text{SYNOP}(t-1) + 1 & \text{if DDE2} \geq 637 \\ \text{SYNOP}(t-1) & \text{otherwise.} \end{cases} \quad (7)$$

$$\text{DDE2}(t) = \sum_{l=0}^2 \text{GDH}(t-l), \quad (8)$$

$$\text{DD57}(t) = \sum_{l=5}^7 \text{GDH}(t-l), \quad (9)$$

where t again is used as a temporal index of days from January 1st onward ($t = 1, 2, \dots, T$). The quantity GDH is a vector of growing degree hours determined by first modeling hourly temperature ($D_{temp}(h)$) from daily T_{min} and T_{max} :

$$D_{temp}(h) = \begin{cases} (T_{max} - T_{min}) \sin(\frac{\pi}{L+4}h) + T_{min} & \text{for } h = 1, 2, \dots, L', \\ D_s - \frac{(D_s - T_{min})}{(\log(24-L)) * \log(h)} & \text{for } h = L' + 1, \dots, 24. \end{cases} \quad (10)$$

where L is the length of the day (in hours) and L' the nearest integer value of L . The term D_s the temperature at sunset, given by:

$$D_s = (T_{max} - T_{min}) \sin(\frac{\pi}{L+4}L) + T_{min}.$$

This formulation assumes a sinusoidal temperature curve during the day that begins at T_{min} just before sunrise and attains a maximum value of T_{max} . Temperatures after sunset are modeled as a logarithmic decay. An illustration of this method for modeling daily temperature evolution is shown in Figure 1.

The growing degree hours are therefore given as:

$$\text{GDH}(t) = \sum_{h=h'}^{h''} D_{temp}(h). \quad (11)$$

Where the quantities h' and h'' are the first and last hours above the base temperature threshold (0.6°C).

We are now prepared to define the matrix \mathbf{A}_{If} of predictor coefficients in equation (3) for the three plants used in the *leaf index*:

$$\mathbf{A}_{\mathbf{lf}} = \begin{matrix} & \text{MDS0} & \text{SYN0P} & \text{DDE2} & \text{DD57} \\ \text{Lilac} & 3.306 & 13.878 & 0.201 & 0.153 \\ \text{Arnold Red} & 4.266 & 20.899 & 0.00 & 0.248 \\ \text{Zabeli} & 2.802 & 21.433 & 0.266 & 0.00 \end{matrix} \quad (12)$$

Development of the *bloom index* proceeds in a similar way, with $\mathbf{X}_{\mathbf{bl}}$ composed of the following predictors:

Predictor abbreviation	Description
MDS0	Days since <i>leaf index</i> was achieved
AGDH	Accumulated Growing Degree Hours

with AGDH is defined as the summation of growing degree hours at some time t from some initial time t_o :

$$\text{AGDH}(t) = \begin{cases} \text{AGDH}(t-1) + \text{GDH}(t) & \text{if } t > t_o \\ 0 & \text{otherwise.} \end{cases} \quad (13)$$

In this case, t_o is taken to be the date of first leaf during any given year.

Finally, $\mathbf{A}_{\mathbf{bl}}$ is defined as:

$$\mathbf{A}_{\mathbf{bl}} = \begin{matrix} & \text{MDS0} & \text{AGDH} \\ \text{Lilac} & -23.934 & 0.116 \\ \text{Arnold Red} & -24.825 & 0.127 \\ \text{Zabeli} & -11.368 & 0.096 \end{matrix} \quad (14)$$

2.4 Implementation in Matlab[©]

The `ml_si` toolbox is distributed as a single zipped tarball. When unzipped and untarred, it will create a main directory called `ML_SI/` with five subdirectories: `data/`, `docs/`, `figs/`, `scripts/` and `si_funcs/`. The actual calculation of SI, however, only requires one function (`calc_si`) and its five dependencies (`calc_daylen`, `calc_soldec`, `growdh`, `synval`, `leaf`), all of which are in the directory `si_funcs/`. One therefore only needs to add the *path* to `si_funcs` to have a fully-functional implementation of `ml_si`. Nonetheless, we supply a number of additional utilities and datasets with the distribution of `ml_si`, all of which are documented in the *User's Guide* in the folder `docs/`. It is worth noting here, however, that in the folder `scripts/` there is a script called `setup_ml_si.m` which will add the appropriate paths and test the functionality of `si_ml`. It is recommended that this script be run first.

The function `calc_si` takes as input three arguments: `Tmin`, `Tmax` and `lat`. The variables `Tmin` and `Tmax` can be 2D matrices or 3D arrays, so long as their *dimensions* are ordered as follows: `years`, `days`, `stations`, where `years` is the total time domain of years for which SI is to be calculated; the *dimension* `days` is the number of days from January 1st onward (366 total, with NaN filling in the last day of all non-leap years); and `stations` is a *dimension* that can be filled by data from one or more stations. It is easiest to think of `calc_si` taking as input a stack of daily `Tmin` and `Tmax` records, with each of the layers in the stack being individual stations or sites. The input values of `Tmin` and `Tmax` should be in units of Fahrenheit, but they need not be derived from station data. The matrices `Alf` and `Abl` are specified in the function `leaf`, but these could be modified in this function, or the function itself could be easily adapted to take these matrices as input.

3 Spring Index Applications

3.1 Data

All data used here to illustrate the functionality of `ml_si` are publicly available. They include “global historical climatology network” (GHCN) daily data from the National Climatic Data Center (<ftp://ftp.ncdc.noaa.gov/pub/data/ghcn/daily/gsn/>), as well as observational phenological datasets archived online by the USA-NPN (<https://www.usanpn.org/>) and provided as part of this toolkit. Examples of how the various metrics are computed from a single GHCN station use daily T_{min} and T_{max} values from station “USC00405187” (Lewisburg, TN, USA).

3.2 Example 1 - SI and predictor variables

Values of T_{min} and T_{max} from a single station for a single year are shown with all derived predictor variables (the time evolving rows of $X_{lf}(t)$) in Figure 2. The day of leaf index output is also shown for that year. A similar example is shown in Figure 3 for the *bloom index*. Both figures illustrate how daily weather fluctuations are translated into a single day-of-year (DOY) value at a given site for a given year for both indices. All data and routines needed to generate these two figures are included in the `ml_si` distribution, and indeed the figures themselves can be reproduced by running the file `ml_si_demo_1.m` in the `scripts/` folder.

3.3 Example 2 - SI calculated from a US weather station

Figure 4 shows simulated leaf (panel **a**) and bloom (panel **b**) events from the three plants comprising the SI models, calculated from a single station for all available years. The values of these time series are average together for each year to produce the leaf and bloom indices shown in Figure 5.

In addition to the leaf and bloom indices, the full suite of *spring indices* include the day of last freeze for each year and the damage index, which records the anomalous amount of time between first leaf and last freeze. Anomalies are used because last freeze typically occurs after the leaf date, but if it happens quite late then presumably the plants have had an unusual amount of time to grow and render themselves vulnerable to frost damage (e.g., Ault et al., 2013). All four SI are shown in Figure 5. Both figures 4 and 5 are generated using the script `ml_si_demo_2.m`.

3.4 Example 3 - SI compared with phenological data

The accuracy of the leaf and bloom indices has been documented elsewhere (see, for example, Schwartz and Reiter, 2000; Schwartz et al., 2006, 2013). Moreover, the `ml_si` code has been validated using the original FORTRAN implementations used in previous studies. Here, nonetheless, we show a qualitative comparison of the leaf and bloom index values (the averages of three plants) with some of the most extensive phenological observations available for the continental United States (Figure 6). To generate these scatter plots, *spring indices* were calculated from a subset GHCND stations that were employed in earlier studies (McCabe et al., 2011; Schwartz et al., 2013; Ault et al., 2013) because of their high data quality and coverage over the 20th century. In Figure 6 the GHCND sites that were within 10km of a lilac observing site are plotted against the phenological events themselves. This figure and all supporting analysis can be generated with the script `ml_si_demo_3.m`, although the user will also need to obtain the raw daily GHCN data (<ftp://ftp.ncdc.noaa.gov/pub/data/ghcn/daily/ghcnd.all.tar.gz>), and store it in the appropriate directory (e.g., `ML_SI/data/ghcnd_all/`).

3.5 Example 4 - SI means and trends in the coterminous US

Regionally-varying trends in global SI are explored extensively by (Schwartz et al., 2006), and interannual variations in North America are documented in Ault et al. (2011) and McCabe et al. (2011), as well as an earlier study documenting variations in observational lilac data (Cayan et al., 2001). In this example, we are interested only in showing how mean dates and trends of

the leaf and bloom indices can be calculated from the subset of GHCN sites used in the previous example (Figure 8). In panel **a**), the overall (1956-2009) mean for each station is shown; trends in these records are shown in panel **b**). These maps differ from those presented in Schwartz et al. (2013) in that no “warming hole” is present in the US southeast: this result is from using fewer records analyzed by Schwartz et al. (2013), and hence continuous coverage in many sites was not available for the full period investigated in the earlier study (1900-2010). Figure 8 can be generated using the script `si_demo_4.m`, although GHCN daily data will first need to be downloaded as with the previous example.

4 Discussion

We have illustrated the underlying conceptual motivation and implementation of the *spring indices* along with their calculation using the `ml_si` toolbox. Several examples using daily GHCN data should give the user a sense for the kinds of analyses that this toolbox makes possible. For instance, in a straightforward way, one could obtain daily station data for a given location where phenological observations have been made, calculate SI, and compare the results with the observed data. Likewise, if daily `Tmin` and `Tmax` values have been recorded at a given plot or phenological monitoring site, they too could be ingested into `ml_si`.

Some of the limitations of `ml_si` are obvious, while others need to be investigated more extensive in future work. For example, the current implementation of SI is only valid in the Northern Hemisphere because of how day lengths are calculated; future developments will include making Southern Hemisphere SI possible. Another drawback is that the models exclude information about precipitation and snow cover; yet, these variables may be critical in governing plant responses at some locations or during some years. By the same token, the detection of synoptic-scale warm weather events is based on T_{min} and T_{max} , but should (ideally) use metrics that record more information about the underlying dynamics of the atmosphere (such as sea level pressure or geopotential height anomalies). Because such variables are often not available when working with international datasets (as in Schwartz et al., 2006), they have been deliberately excluded from the calculation of SI for simplicity and flexibility.

Two more subtle, and perhaps more serious, limitations of SI are worth mentioning. First, the extension of SI into lower latitudes was achieved by removing the plant “chilling” requirements as explained in Schwartz et al. (2013), which had been a key feature of the model before that time. Although this step extended the geographic domain where SI can be calculated, its overall implications remain relatively under-explored. Nonetheless, Schwartz et al. (2013) show good

agreement between the extended SI and existing (non-Lilac and non-Honeysuckle) observational datasets in warmer temperate regions, suggesting that even in these lower latitudes the SI pick up key features of spring onset. Second, the SI are composed of average simulated leaf out and bloom dates from just three indicator species. Although the phenologies of these three species are perhaps the most widely observed in the continental US, it would be desirable at some future time to populate the SI plant models with a richer “ecology” of indicator species. Despite these limitations, the SI have consistently and demonstrably been useful in tracking interannual variations in spring onset and trends across an impressive number of spatial domains and types of variables (Schwartz and Reiter, 2000; White et al., 2009; McCabe et al., 2011; Schwartz et al., 2013). In our view, the SI are therefore best used to characterize interannual variations in temperature-driven phenological and physical responses to spring onset.

Finally, the use and utility of the SI can be thought of in an analogous way to that of the “Palmer Drought Severity Index” (PDSI) and its many variants (e.g., Dai, 2011). The PDSI was developed for a very narrow purpose in a limited geographic domain nearly 50 years ago. Nonetheless, it manages to capture something essential about the nature of moisture balances and deficits across a wide range of spatial and temporal scales, making it one of the most widely used drought indices even today, and in spite of its many limitations (see Alley, 1984; McKee et al., 1993; Guttman, 1999; Redmond, 2002; Dai, 2011). In our view, the SI are conceptually similar to drought indices because they attempt to translate noisy local weather fluctuations into a consistent indicator of climatic conditions across space and through time. By producing and distributing `ml_si`, it is hoped that the wider research community will continue to explore its usefulness and limitations, as well as develop new, alternative indices to characterize spring onset in our changing climate.

5 Appendix A: Additional resources

Complete documentation of the `si_ml` toolbox may be found in the *User’s Guide* (`ML_SI/docs/users_guide.pdf`). Here, we note a few additional functions and datasets that are distributed in the package. First, the script `setup_ml_si.m` will add the appropriate path to the SI functions and test various options, including the functionality of `wget` to retrieve station data as well as the routines needed to import GHCN metadata, GHCN daily data, and the North American lilac data. None of these options are needed, however, for the core functions comprising `ml_si` to work properly.

Second, the function `read_ghcnd_dly_file` (in `si_funcs/`) will read a GHCN file with daily

station data and return a *structure* with appropriate metadata and data attached to it. This function is called by `si_demo_3.m`, `si_demo_4.m`.

Lastly, the data found in `ML_SI/data/mds/mds_verification_data/` includes `SI-x_1981_2010_norms25_noWBAN.xls` and `SI-x_select6_optimized_output.xls`. The first Excel[©] file contains the list of the 733 quality-controlled GHCND sites (first column) as well as some metadata and output from the original FORTRAN code written by M. D. Schwartz. This file is used by `si_demo_3.m` to calculate spring indices for each of the 733 sites; `si_demo_4.m` uses the output from `si_demo_3.m` and maps out the trends and means for these sites. A second file (`SI-x_select6_optimized_output.xls`) contains the output from Schwartz’s original FORTRAN code to calculate SI from a “Select 6” subset of the GHCN database. The output has also been formatted for Excel[©]. If modifications are made to any of the core functions comprising `ml_si`, it is recommended that the new results be verified against this dataset (an example of this verification process is found in `setup_ml_si`). It is not necessary, in our view, that new output be exactly 1-to-1 with the original FORTRAN output because truncation differences between Matlab and FORTRAN can introduce up to 1 day of random error during some years. However, any errors should be uncorrelated and not systematically biased in any one direction.

Acknowledgments

Coordination and effort to develop this project was made possible in part by the USA National Phenology Network, including J.L. Betancourt and J.F. Weltzin who provided helpful discussions during the development of this toolkit. We are especially grateful to Alyssa Rosemartin and the National Coordinating Office of the USA-NPN for supplying quality-controlled versions of the historical Lilac phenology datasets from the Eastern and Western US. The project described in this publication was supported by Grant/Cooperative Agreement Number G13AC00248 from the United States Geological Survey. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the USGS.

References

- Alley, W.M., 1984. The Palmer Drought Severity Index - Limitations and Assumptions. *Journal of Climate and Applied Meteorology* 23, 1100–1109.
- Ault, T.R., Henebry, G.M., de Beurs, K.M., Schwartz, M.D., Betancourt, J.L., Moore, D.,

2013. The false spring of 2012, earliest in north american record. *Eos, Transactions American Geophysical Union* 94, 181–182. URL: <http://dx.doi.org/10.1002/2013EO200001>, doi:10.1002/2013EO200001.
- Ault, T.R., Macalady, A.K., Pederson, G.T., Betancourt, J.L., Schwartz, M.D., 2011. Northern hemisphere modes of variability and the timing of spring in western north america. *Journal of Climate* 24, 4003–4014. URL: <http://dx.doi.org/10.1175/2011JCLI4069.1>, doi:10.1175/2011JCLI4069.1.
- Cayan, D.R., Kammerdiener, S.A., Dettinger, M.D., Caprio, J.M., Peterson, D.H., 2001. Changes in the onset of spring in the western United States. *Bulletin of the American Meteorological Society* 82, 399–415.
- Dai, A., 2011. Characteristics and trends in various forms of the Palmer Drought Severity Index during 1900–2008. *Journal of Geophysical Research-Atmospheres* 116. doi:10.1029/2010JD015541.
- Guttman, N.B., 1999. Accepting the standardized precipitation index: A calculation algorithm. *Journal of the American Water Resources Association* 35, 311–322.
- IPCC, 2007. Chapter 1: Assessment of observed changes and responses in natural and manage systems., in: Working Group II: Impacts, Adpatation, and Vulnerability. Cambridge University Press, United Kingdom and New York, NY, USA.
- Janetos, A.C., Chen, R.S., Arndt, D., Kenney, M.A., 2012. National Climate Assessment Indicators: Background, Development, and Examples. Technical Report. National Climate Assessment, U.S. Global Change Research Program, Washington DC.
- Lieth, H., 1974. Phenology and seasonality modeling. *Ecological Studies* 8.
- McCabe, G.J., Ault, T.R., Cook, B.I., Betancourt, J.L., Schwartz, M.D., 2011. Influences of the El Niño southern oscillation and the pacific decadal oscillation on the timing of the north american spring. *International Journal of Climatology* , n/a–n/aURL: <http://dx.doi.org/10.1002/joc.3400>, doi:10.1002/joc.3400.
- McKee, T.B., Doeskin, N.J., Kleist, J., 1993. The Relationship of Drought Frequency and Duration to Time Scales. *Proc. 8th Conf. on Aplied Climatology* , 179–184.
- Redmond, K.T., 2002. The depiction of drought - A commentary. *Bulletin of the American Meteorological Society* 83, 1143–1147.

- Schwartz, M., 1985. The advance of phenological spring across Eastern and Central North America. Ph.D. thesis. University of Kansas.
- Schwartz, M., Marotz, G., 1986. An approach to examining regional atmosphere plant interactions with phenological data. *Journal of Biogeography* 13, 551–560.
- Schwartz, M.D., 1999. Advancing to full bloom: planning phenological research for the 21st century. *International Journal of Biometeorology* 42, 113–118.
- Schwartz, M.D., Ahas, R., Aasa, A., 2006. Onset of spring starting earlier across the Northern Hemisphere. *Global Change Biology* 12, 343–351.
- Schwartz, M.D., Ault, T.R., Betancourt, J.L., 2013. Spring onset variations and trends in the continental united states: past and regional assessment using temperature-based indices. *International Journal of Climatology* URL: <http://dx.doi.org/10.1002/joc.3625>, doi:10.1002/joc.3625.
- Schwartz, M.D., Betancourt, J.L., Weltzin, J.F., 2012. From caprio’s lilacs to the usa national phenology network. *Frontiers in Ecology and the Environment* 10, 324–327. URL: <http://dx.doi.org/10.1890/110281>, doi:10.1890/110281.
- Schwartz, M.D., Marotz, G.A., 1988. Synoptic Events and Spring Phenology. *Physical Geography* 9, 151–161.
- Schwartz, M.D., Reiter, B.E., 2000. Changes in North American spring. *International Journal of Climatology* 20, 929–932.
- van Vliet, A., Groot, R., Bellens, Y., Braun, P., Bruegger, R., Bruns, E., Clevers, J., Estreguil, C., Flechsig, M., Jeanneret, F., Maggi, M., Martens, P., Menne, B., Menzel, A., Sparks, T., 2003. The european phenology network. *International Journal of Biometeorology* 47, 202–212. URL: <http://dx.doi.org/10.1007/s00484-003-0174-2>, doi:10.1007/s00484-003-0174-2.
- White, M.A., de Beurs, K.M., Didan, K., Inouye, D.W., Richardson, A.D., Jensen, O.P., O’Keefe, J., Zhang, G., Nemani, R.R., van Leeuwen, W.J.D., Brown, J.F., de Wit, A., Schaepman, M., Lin, X.M., Dettinger, M., Bailey, A.S., Kimball, J., Schwartz, M.D., Baldocchi, D.D., Lee, J.T., Lauenroth, W.K., 2009. Intercomparison, interpretation, and assessment of spring phenology in North America estimated from remote sensing for 1982–2006. *Global Change Biology* 15, 2335–2359.

Zurita-Milla, R., van Gijsel, J., Hamm, N., Augustijn, P., Vrieling, A., 2013. Exploring spatiotemporal phenological patterns and trajectories using self-organizing maps. *Geoscience and Remote Sensing, IEEE Transactions on* 51, 1914–1921. doi:10.1109/TGRS.2012.2223218.




Figure 1: Illustration of how the GDH model is employed using T_{min} and T_{max} values of 20°F and 60°F, respectively. The red shading shows the area under the curve above the base temperature (31°F). These values would be typical of many temperate regions during spring.




Figure 2: Illustration of the terms (SYNOP, DDE2, and DD57) comprising the predictor variables of X that change as function of time. This example is from a single year (1902) at a single weather station in the GHCN (USC00405187). The top panel shows T_{\min} and T_{\max} , and the next one shows GDH calculated from those two variables. GDH is then used to identify synoptic events (SYNOP), as well as the amount of heat accumulation on days 0-2 and 5-7 prior. In all panels, the first leaf date is denoted by the dashed green vertical line.




Figure 3: Same as figure 2, but for the bloom index.




Figure 4: Time series of simulated phenology of three plants (Lilac, Arnold Red, and Zabeli) whose simulated leaf and bloom dates are modeled to compute the spring indices from a single station (USC00405187). **a)** “First Leaf” dates for each of the three plants; **b)** “First Bloom” dates. These dates from the individual plants are averaged together to create the leaf and bloom indices (see Figure 5).




Figure 5: Time series of the four indices comprising SI, again calculated for a single station (USC00405187). The top shows the three indices with values that are days of the calendar year (DOY), while the bottom panel shows the Damage Index, which records the anomalous amount of time between first leaf and last freeze. This index is expressed as an anomaly because its raw values are almost always negative (first leaf typically occurs before last freeze); when expressed as departures from the long term mean, the most negative values correspond to the most anomalously long periods of time between first leaf and last freeze, and hence the most potentially deleterious impacts on plant growth.





Figure 6: **a)** Leaf Index and **b)** Bloom Index values calculated from GHCN data across North America and plotted against the nearest (within 10km) first leaf and first bloom observations. The observational data are archived by the National Phenology Network (<https://www.usanpn.org>), and are distributed with this toolkit.

a) Leaf Index Means



b) Bloom Index Means





Figure 7: GHCN-derived mean values of the **a)** *leaf index*; and **b)** *bloom index*. GHCN stations are the same as those used to produce figure 6, and were selected because they were the closest stations with maximum overlap in time with the observational lilac and honeysuckle data. However, all available data were used to produce this figure. Means were computed for all available data over the period 1956-2009

a) Leaf Index Trends



b) Bloom Index Trends




Figure 8: GHCN-derived trends (days/decade) of the **a)** *leaf index*; and **b)** *bloom index*. GHCN stations are the same as those used to produce figure 6, and were selected because they were the closest stations with maximum overlap in time with the observational lilac and honeysuckle data. As in Figure 7, all available data were used to produce this figure. Trends were computed for all available data over the period 1956-2003.