

PROGRAMMING AND PROBLEM SOLVING (SE 1105) MIDTERM	A	Grading				
		Q1	Q2	Q3	Q4	Σ
Instructors	ID #	Name-Surname	Time	Date/Room #		
Dr. Dindar ÖZ Dr. Kazım ERDOĞDU Dr. Mete EMİNAĞAOĞLU			80 mins.	November 19, 2024 (09:40-..)		

Notes: If you believe that necessary data or assumptions are missing from the problem statement, make your own assumption(s) and write them clearly.

QUESTIONS

1. (30 pts.) Write the outputs of the following C programs.

a) (15pts)

```
#include <stdio.h>

int g(int m, int M[], int n)
{
    int i = 0;
    while (i < n)
    {
        if (m == M[i])
        {
            return i;
        }
        i++;
    }
    return -1;
}

void f(int A[], int B[], int n)
{
    int i = 0, p = A[0];
    while (B[i] != A[0])
    {
        i = g(B[i], A, n);
        printf("%d --> %d\n", p, A[i]);
        p = A[i];
    }
    printf("%d --> %d\n", p, B[i]);
}

void main()
{
    int A[] = {6, 0, 3, 1, 5, 2, 4};
    int B[] = {1, 2, 4, 5, 3, 0, 6};
    f(A, B, 7);
}
```

b) (15pts)

```
#include <stdio.h>

void main()
{
    int i = 1, j, par1, par2;
    int lastr = 5;

    par1 = lastr + 1;

    while (i <= lastr)
    {
        par2 = par1;
        while (par2 >= 1)
        {
            printf(" ");
            par2--;
        }

        j = 1;
        while (j <= i)
        {
            printf("%d ", i);
            j++;
        }

        printf("\n");
        par1--;
        i++;
    }
}
```


PROGRAMMING AND PROBLEM SOLVING (SE 1105) MIDTERM	A	Grading				
		Q1	Q2	Q3	Q4	Σ
Instructors	ID #	Name-Surname		Time	Date/Room #	
Dr. Dindar ÖZ Dr. Kazım ERDOĞDU Dr. Mete EMİNAĞAOĞLU				80 mins.	November 19, 2024 (09:40-..)	

3. (25 pts.) Write a C function that reads two positive integers (a and b) from the user and prints all perfect squares between a and b. Assume that the user always enters positive integers and a smaller number first. (Hint: Perfect squares are 1,4,9,16,25...)

Attention: You cannot use `<math.h>` and `sqrt()` function in your solution.

Example:

```
Enter the first number: 2
Enter the second number: 18
Perfect squares between 2 and 18 are: 4 9 16

Enter the first number: 5
Enter the second number: 90
Perfect squares between 5 and 90 are: 9 16 25 36 49 64 81
```

PROGRAMMING AND PROBLEM SOLVING (SE 1105) MIDTERM	A	Grading				
		Q1	Q2	Q3	Q4	Σ
Instructors	ID #	Name-Surname	Time	Date/Room #		
Dr. Dindar ÖZ Dr. Kazım ERDOĞDU Dr. Mete EMİNAĞAOĞLU			80 mins.	November 19, 2024 (09:40-..)		

4. (25 pts.) Write a C function that takes the measured average daily temperature of a city for a full month (30 days) as an integer array of length 30. The function returns the biggest temperature change after the hottest day of the month that occurs between two consecutive days. Assume that there is only one hottest day in the array

Example: If the array: { 10, 15, 12, 14, 13, 12 } then it must return 15-12 = 3
 If the array: { 10, 12, 18, 14, 11, 10, 15, 12 } then it must return 15-10=5
 If the array: { 12, 11, 12, 13, 13, 14, 10, 15 } then it must return 0 as there is no temp change after 15.

Good luck...