TARGET STANDARD: NEET UG (MEDICAL ENTRANCE)

Biotechnology: Principles and Processes

Complete Conceptual Module, High-Yield Theory, Diagrammatic Models, and Exam Practice

1. Learning Objectives and Exam Blueprint

In recent NEET UG examinations, the unit on **Biotechnology** accounts for roughly 5 to 6 questions (~20 to 24 marks), with **Chapter 9: Principles and Processes** forming the mechanistic bedrock. Candidates are expected to master:

- **Definition & Scope:** Traditional vs. modern molecular biotechnology and the EFB definition.
- **Core Principles:** Genetic engineering (rDNA construction) and bioprocess engineering (sterile scaling).
- **Historical Milestones:** Boyer and Cohen's foundational 1972 experiment.
- **Tools of rDNA Technology:** Restriction endonucleases (nomenclature, recognition sequences, palindromes, staggered cleavage, sticky ends), DNA ligase, alkaline phosphatase, gel electrophoresis (agarose matrix, ethidium bromide, UV detection, elution), cloning vectors (ori, selectable markers, unique cloning sites, insertional inactivation, pBR322 architecture, Ti plasmid of *Agrobacterium tumefaciens*, disarmed retroviruses), and host competency techniques.
- **Recombinant Processes:** Isolation of pure DNA (enzymatic lysis, spooling), Polymerase Chain Reaction (denaturation, annealing, extension, Taq polymerase kinetics), bioreactors (simple stirred-tank vs. sparged stirred-tank), and downstream processing.

2. Foundations and Core Principles

2.1 Definition of Biotechnology

Biotechnology deals with techniques of using live organisms or enzymes from organisms to produce products and processes useful to humans. Traditional processes like making curd, bread, or wine are microbially mediated, but the term is applied today in a restricted sense: processes that utilize **genetically modified organisms (GMOs)** to achieve large-scale production. Modern biotechnology encompasses techniques such as *in vitro* fertilisation (test-tube baby programme), chemical gene synthesis, DNA vaccine synthesis, and gene therapy.

> **EFB DEFINITION OF BIOTECHNOLOGY**
>
> The **European Federation of Biotechnology (EFB)** formulated an integrated definition encompassing both traditional views and modern molecular biotechnology:
> *"The integration of natural science and organisms, cells, parts thereof, and molecular analogues for products and services."*

2.2 The Two Pillars of Modern Biotechnology

1. **Genetic Engineering:** Techniques to alter the chemistry of genetic material (DNA and RNA), introduce this modified material into host organisms, and thereby change the phenotype of the recipient host.
2. **Bioprocess Engineering:** Maintenance of a sterile (contamination-free) ambience in chemical engineering processes to allow the selective growth of only the desired microbe or eukaryotic cell in large quantities for the manufacture of antibiotics, vaccines, enzymes, and therapeutics.

2.3 Sexual vs. Asexual Reproduction & The Power of Genetic Engineering

Asexual reproduction preserves genetic information strictly without change, whereas sexual reproduction permits variations and novel genetic combinations. Traditional hybridisation programmes in plant and animal breeding invariably result in the co-inheritance and multiplication of undesirable genes alongside desirable ones.

BOSONANCE

Recombinant DNA technology bypasses this bottleneck via three fundamental steps:

1. Identification of DNA with desirable genes.
2. Introduction of the identified foreign DNA into a suitable host.
3. Maintenance of introduced DNA in the host and faithful transfer to its progeny.

2.4 Conceptual Breakthrough: The Cohen & Boyer Experiment (1972)

An alien piece of DNA introduced into an organism cannot replicate autonomously unless it becomes an integrated part of a chromosome having a specific DNA sequence called the **origin of replication (ori)**. The *ori* initiates replication and determines the copy number of the propagating DNA.

In **1972**, **Stanley Cohen** and **Herbert Boyer** synthesized the first artificial recombinant DNA (rDNA) molecule:

- Boyer had observed that specific restriction enzymes isolated from *Escherichia coli* cleave DNA leaving single-stranded overhanging ends ("sticky ends").
- Cohen had pioneered techniques to isolate autonomous, circular extrachromosomal DNA rings called **plasmids** and reintroduce them into bacterial cells.
- They isolated an antibiotic resistance gene by cutting a native plasmid of *Salmonella typhimurium* using restriction endonucleases.
- This gene was spliced into a plasmid vector using **DNA ligase**, creating an autonomous circular rDNA molecule *in vitro*.
- The recombinant plasmid was transferred into *Escherichia coli*, where it replicated using host DNA polymerase, cloning the antibiotic resistance gene.

3. Tools of Recombinant DNA Technology

3.1 Restriction Endonucleases ("Molecular Scissors")

In **1963**, scientists isolated two enzymes from *E. coli* that restricted bacteriophage propagation:

- One added methyl groups ($-\mathrm{CH}_3$) to bacterial DNA to protect host DNA from self-cleavage.
- The other cleaved foreign viral DNA and was termed **restriction endonuclease**.

The first sequence-specific restriction endonuclease characterized was ***Hind* II** (isolated five years later from *Haemophilus influenzae* Rd). It consistently cleaves DNA double helices by recognizing a specific sequence of **6 base pairs**. Today, over **900 restriction enzymes** have been isolated from more than **230 bacterial strains**, each recognizing distinct sequences.

Nomenclature Rules:

- First letter: Genus of the prokaryote (capitalized; e.g., *E* for *Escherichia*).
- Next two letters: Species of the prokaryote (lowercase; e.g., *co* for *coli*).
- Fourth letter: Strain designation (e.g., *R* for strain RY 13).
- Roman numeral: Order in which the enzyme was isolated from that particular strain (e.g., I for first isolated).

Exonucleases vs. Endonucleases:

Feature	Exonucleases	Restriction Endonucleases
Site of Cleavage	Remove nucleotides sequentially from terminal ends (5' or 3') of DNA.	Make internal cuts at precise, specific phosphodiester positions within the DNA.
Sequence Specificity	Generally non-specific; degrades terminal ends.	Strictly sequence-specific; recognizes distinct palindromic motifs.
Role in Genetic Engineering	Limited; used for trimming DNA fragments.	Central tool; creates defined fragments and cohesive sticky ends.

BOSONANCE

Palindromes and Staggered Cleavage

A DNA palindrome is a sequence of base pairs that reads identical on both anti-parallel strands when reading in the same orientation (either $5' \rightarrow 3'$ or $3' \rightarrow 5'$).

For example, the recognition sequence for ***Eco*RI**:

$$5'\text{---G}\downarrow\text{A---A---T---T---C---}3'$$
$$3'\text{---C---T---T---A---A}\uparrow\text{G---}5'$$

*Eco*RI binds the hexanucleotide palindrome and cuts both strands between the same two bases (G and A) slightly away from the center of symmetry. This staggered cut leaves short, single-stranded unpaired extensions at each terminus called **sticky ends**. These overhanging stretches readily form hydrogen bonds with complementary counterparts cut by the identical restriction enzyme. The stickiness greatly accelerates the final covalent sealing of the sugar-phosphate backbone by **DNA ligase**.




Figure 1: Cleavage of foreign DNA and vector plasmid by EcoRI producing matching cohesive sticky ends, followed by covalent annealing using DNA ligase.

3.2 Separation and Isolation of DNA Fragments: Agarose Gel Electrophoresis

Restriction digestion produces a heterogeneous mixture of DNA fragments. These fragments are separated using **agarose gel electrophoresis** based on the following physical principles:

- **Charge Property:** DNA fragments possess a constant negative charge across their length due to negatively charged phosphate groups (PO_4^{3-}). Hence, under an applied electrical potential gradient, DNA fragments migrate toward the **anode (positive electrode)** away from the cathode (wells).
- **Matrix:** Agarose is a natural polysaccharide polymer extracted from **seaweeds**. It polymerizes into a cross-linked network that provides a **sieving effect**.
- **Resolution by Size:** Smaller DNA fragments navigate through the matrix pores with less friction and migrate faster and farther from the loading wells, whereas larger fragments migrate shorter distances. Migration distance is inversely proportional to the log of fragment molecular weight.
- **Visualisation:** Pure DNA fragments cannot be seen under visible white light. The gel must be stained with **ethidium bromide (EtBr)** and exposed to **ultraviolet (UV) radiation**, revealing vibrant **bright orange-colored bands**.
- **Elution:** The isolated band containing the desired gene is physically cut out from the agarose gel slab, and the DNA is extracted from the gel slice. This crucial purification step is termed **elution**.

BOSONANCE




Figure 2: Agarose gel electrophoresis showing resolution of negatively charged DNA fragments from cathode to anode according to molecular size through sieving action.

3.3 Cloning Vectors

Vectors are DNA vehicles used to deliver foreign target DNA into a host cell for autonomous replication. Commonly utilized vectors include bacterial **plasmids** (circular, extrachromosomal dsDNA; typically 1–100 copies per cell) and **bacteriophages** (viruses infecting bacteria; high copy number per cell).

Essential Characteristics of an Engineered Cloning Vector:

1. **Origin of Replication (ori):** A specific nucleotide sequence where replication commences. Any alien DNA linked to *ori* replicates in synchrony with the vector. It also strictly **controls the copy number** of the cloned DNA.
2. **Selectable Marker:** A gene that permits the identification and selection of transformants while selectively eliminating non-transformants. Transformation is the procedure by which exogenous DNA is introduced into a recipient bacterium. Genes conferring resistance to antibiotics like **ampicillin** ($\mathrm{amp}^{\mathrm{R}}$), **tetracycline** ($\mathrm{tet}^{\mathrm{R}}$), **chloramphenicol**, or **kanamycin** serve as standard selectable markers for *E. coli*, as normal wild-type *E. coli* strains are sensitive to all these antibiotics.
3. **Cloning (Restriction) Sites:** To ligate alien DNA, the vector requires very few, preferably a **single (unique) recognition site** for commonly used restriction enzymes. If more than one recognition site exists for a given enzyme within the vector, digestion will cut the vector into multiple fragments, severely complicating gene cloning.
4. **Small Size / Low Molecular Weight:** Facilitates easy isolation, manipulation, and high-efficiency transformation into host cells.

3.4 Anatomy of Plasmid pBR322 and Insertional Inactivation

The artificial cloning vector **pBR322** constructed in *E. coli* exhibits a precisely engineered distribution of restriction sites, marker genes, and regulatory regions:

- **amp^{R} Gene (Ampicillin resistance):** Possesses unique recognition sites for ***Pst* I** and ***Pvu* I**.
- **tet^{R} Gene (Tetracycline resistance):** Contains unique recognition sites for ***Bam*H I** and ***Sal* I**.
- **Additional Unique Restriction Sites: *Eco*R I**, ***Cla* I**, and ***Hind* III** (located close together outside the antibiotic genes).
- **ori:** Origin of replication sequence controlling replication initiation and copy number.
- **rop:** Codes for proteins involved in the replication of the plasmid; contains a unique cleavage site for ***Pvu* II**.

BOSONANCE




Figure 3: Restriction map of plasmid vector pBR322 indicating selectable markers (amp^R and tet^R), rop, ori, and unique restriction cleavage sites.

Selection via Dual Antibiotic Plating (Replica Plating):

When a foreign gene is ligated into the ***Bam*H I** site of pBR322, it disrupts the coding frame of the tetracycline resistance gene. This loss of phenotypic function caused by foreign DNA insertion is called **insertional inactivation**.

- **Non-transformants:** Cells that fail to take up any plasmid remain sensitive to both ampicillin and tetracycline and die on ampicillin-containing medium.
- **Transformants:** Cells that absorb a plasmid grow on ampicillin plates.
- **Recombinants vs. Non-recombinants:** Transformant colonies from the ampicillin master plate are replicated onto tetracycline agar plates:
  - **Recombinants:** Plasmids carry the insert in tet^R. Hence, they grow on ampicillin medium but *fail to grow* on tetracycline medium.
  - **Non-recombinants (self-ligated vector):** Carry intact tet^R. They grow on *both* ampicillin and tetracycline media.

3.5 Alternative Selectable Marker: Chromogenic Selection (Blue-White Screening)

Selection by dual-antibiotic replica plating is labor-intensive and slow because it mandates simultaneous plating across two sets of agar plates. To overcome this, modern vectors utilize **chromogenic selection** based

on the insertional inactivation of the enzyme β-**galactosidase**:

- The cloning site resides within the coding sequence of the β-galactosidase gene (e.g., *lacZ*).
- **Non-recombinant Plasmids:** The uninterrupted β-galactosidase gene produces functional active enzyme. In the presence of a chromogenic substrate (e.g., X-gal), the colonies turn **blue**.
- **Recombinant Plasmids:** Insertion of the foreign DNA fragment causes **insertional inactivation** of the β-galactosidase gene. No active enzyme is produced, and the recombinant colonies remain **white / colorless**.

3.6 Vectors for Cloning in Plants and Animals

1. **Plant Vectors (*Agrobacterium tumefaciens*):** A soil bacterium pathogenic to many dicotyledonous plants. It naturally transfers a defined DNA segment called **T-DNA (transfer DNA)** from its large **Tumor-inducing (Ti) plasmid** into host plant chromosomes, transforming normal plant cells into crown gall tumors that synthesize specialized chemicals required by the pathogen. In biotechnology, the Ti plasmid has been **disarmed** by deleting oncogenic tumor-forming genes while retaining its natural transfer machinery, making it an efficient vehicle for delivering genes of interest into crops.
2. **Animal Vectors (Retroviruses):** Retroviruses naturally transform normal animal cells into cancerous cells. By disarming retroviruses (stripping their oncogenes and pathogenicity), they are converted into vectors that stably transduce desired therapeutic genes into mammalian host genomes.

3.7 Competent Host and Methods of Gene Transfer

Because DNA is a negatively charged, highly **hydrophilic** molecule, it cannot passively traverse the hydrophobic core of phospholipid cell membranes. Specific physical and chemical treatments make host cells **competent** to take up exogenous DNA:

Gene Transfer Method	Principle & Mechanism	Target Host System
Divalent Cation + Heat Shock	Cells are treated with a divalent cation (e.g., Ca^{2+}), which neutralizes repulsive negative charges and increases cell wall pore permeability. The cells and rDNA are incubated on ice $(0^{\circ}\mathrm{C})$, exposed briefly to $42^{\circ}\mathrm{C}$ **(heat shock)**, and immediately returned to ice.	Bacterial hosts (e.g., *E. coli*).
Micro-injection	Recombinant DNA is directly injected into the **nucleus** of a target cell using an ultra-fine glass micropipette under a high-resolution inverted microscope.	Animal cells (oocytes/zygotes).
Biolistics / Gene Gun	Microscopic heavy-metal particles of **gold or tungsten** are coated with recombinant DNA and accelerated at high velocities into target cells using helium pressure.	Plant cells (penetrates thick plant cell walls).
Disarmed Pathogen Vectors	Genetically engineered, non-pathogenic bacteria or viruses infect host cells naturally and transfer rDNA into the host genome.	Plants (*Agrobacterium*) and animals (retroviruses).

4. Processes of Recombinant DNA Technology

Constructing and expressing recombinant DNA follows an orderly sequence of distinct molecular and biochemical phases:

4.1 Step 1: Isolation of Genetic Material (DNA)

Nucleic acids are enclosed within cellular membranes associated with proteins, lipids, and polysaccharides. Purified DNA free of macromolecules is isolated through sequential steps:

1. **Enzymatic Cell Lysis:** Digesting cell envelopes using specific enzymes:
   - **Bacterial cells: Lysozyme** (hydrolyzes peptidoglycan cell walls).
   - **Plant cells: Cellulase** and pectinase (degrades cellulose).
   - **Fungal cells: Chitinase** (digests chitinous walls).
2. **Macromolecular Removal:** Cellular RNA is degraded using **ribonucleases (RNase)**, while proteins (including chromatin histones) are digested using **proteases**.

BOSONANCE

3. **Precipitation with Chilled Ethanol:** DNA is insoluble in cold alcohol. Adding **chilled ethanol** forces pure DNA to precipitate out of aqueous solution, appearing as a suspension of fine, white, thread-like fibers.
4. **Spooling:** The precipitated filamentous DNA threads are collected onto a glass rod in a physical extraction process called **spooling**.

4.2 Step 2: Cutting of DNA at Specific Locations

Purified target genomic DNA and chosen vector DNA are separately incubated with the selected restriction endonuclease under optimal temperature, salt, and pH conditions. Digestion progress is monitored by running reaction aliquots on an agarose gel. The excised target gene and opened vector are then mixed together with **DNA ligase** to produce circular recombinant DNA.

4.3 Step 3: Amplification of Gene of Interest by PCR

The **Polymerase Chain Reaction (PCR)**, devised by Kary Mullis, is an *in vitro* enzymatic amplification technique used to synthesize billions of copies of a target DNA sequence from trace templates.

Essential Reaction Components:

- **DNA Template:** Double-stranded DNA segment containing the region to be amplified.
- **Primers:** Two sets of chemically synthesized, short oligonucleotide sequences (~18–25 nucleotides) complementary to opposite ends of the target region.
- **Deoxyribonucleoside triphosphates (dNTPs):** dATP, dCTP, dGTP, dTTP as building blocks.
- **Thermostable DNA Polymerase: *Taq* polymerase**, isolated from the thermophilic bacterium ***Thermus aquaticus***. It withstands high temperatures without denaturing.
- **Divalent Cations:** Mg^{2+} acting as an essential cofactor for polymerase activity.

The Three Cyclic Steps of PCR:

1. **Denaturation ($\approx 94^{\circ}\mathrm{C}$ to $96^{\circ}\mathrm{C}$):** Thermal heating breaks the hydrogen bonds holding the antiparallel strands together, separating double-stranded DNA into single strands.
2. **Primer Annealing ($\approx 50^{\circ}\mathrm{C}$ to $60^{\circ}\mathrm{C}$):** The temperature is reduced to permit the two oligonucleotide primers to hybridize to their complementary sequences at the 3' ends of the single-stranded DNA templates.
3. **Extension ($\approx 72^{\circ}\mathrm{C}$):** At the optimal temperature for *Taq* polymerase, the enzyme extends the primers by synthesizing new complementary strands in the $5' \rightarrow 3'$ direction using the free dNTPs in the reaction mixture.

AMPLIFICATION FORMULA & 30-CYCLE YIELD

For n cycles of PCR, theoretical amplification follows an exponential factor of 2^n.
After **30 cycles** of PCR:

$$\text{Amplified copies} = 2^{30} = 1{,}073{,}741{,}824 \approx 10^9 \text{ copies } (1 \text{ billion-fold amplification})$$

BOSONANCE




Figure 4: Schematic of one PCR amplification cycle showing thermal denaturation, primer hybridization, and Taq polymerase extension.

4.4 Step 4: Insertion of Recombinant DNA into Host

The ligated plasmid is transferred into chemically competent recipient cells (e.g., *E. coli*). When the plasmid carries a selectable marker such as $\mathrm{amp}^{\mathrm{R}}$, spreading the cell suspension on ampicillin-containing agar isolates transformants: only cells that acquired the resistance gene survive, while untransformed host cells are eliminated.

4.5 Step 5: Large-Scale Production and Bioreactors

When a protein-encoding gene is expressed in a heterologous host, the resulting product is known as a **recombinant protein**. Small-scale laboratory shake flasks cannot yield commercially viable quantities. Industrial mass manufacturing requires large closed vessels called **bioreactors** (processing working volumes of **100 to 1,000 liters**).

Continuous Culture System:

In a continuous culture system, depleted used medium is drained from one side while fresh nutrient medium is continuously supplied from the other. This dynamic steady-state maintains growing cells in their physiologically most active **logarithmic (exponential) phase**, yielding a significantly higher biomass and maximum recombinant protein output compared to stationary batch cultures.

Bioreactor Designs:

A bioreactor provides optimum biological growth conditions by precisely regulating temperature, pH, substrate levels, oxygen, vitamins, and salts.

- **Simple Stirred-Tank Bioreactor:** Typically features a cylindrical vessel with a **curved base** to eliminate dead zones and ensure thorough mixing. It is equipped with:
  - An **agitator system** with flat-bladed impellers for continuous mixing.
  - An **oxygen delivery system** with a sparger for air aeration.
  - A **foam-breaker system** to control surface froth.
  - Temperature and pH control sensors with acid/base addition ports.
  - Steam injection ports for *in situ* thermal sterilization.
  - **Sampling ports** to periodically withdraw small aliquots for testing.

BOSONANCE

- **Sparged Stirred-Tank Bioreactor:** Sterile air bubbles are actively sparged (forced under pressure) through the liquid broth. These millions of tiny bubbles dramatically increase the surface area available for oxygen mass transfer into the culture medium.




Figure 5: Structural comparison between a simple stirred-tank bioreactor and a sparged bioreactor showing gas entrainment and mass aeration mechanisms.

4.6 Step 6: Downstream Processing

Once the biosynthetic phase inside the bioreactor concludes, the target product must undergo a series of post-fermentation recovery steps collectively termed **downstream processing**:

- **Separation:** Filtering cell debris, harvesting cells, or precipitating crude broth proteins.
- **Purification:** Fractional purification through chromatography (ion-exchange, affinity, gel-filtration).
- **Formulation:** Stabilizing the product with suitable non-toxic preservatives.
- **Clinical Trials:** Compulsory for therapeutic biopharmaceuticals to establish human safety, efficacy, and dosage pharmacokinetics.
- **Quality Control (QC) Testing:** Strict batch-to-batch analytical assays to evaluate product purity, specific activity, and stability.

> **NEET TRAP ALERT**
>
> Downstream processing encompasses only **separation**, **purification**, **formulation**, and **quality control**. It does **NOT** include upstream operations such as gene cloning, transformation, expression induction, or biomass growth inside the bioreactor.

BOSONANCE

5. NEET UG Conceptual Worked Examples

Worked Example 1

CALCULATIONS & PCR KINETICS

A molecular biologist begins a polymerase chain reaction with 6 double-stranded DNA template molecules. How many double-stranded DNA molecules will be present in the reaction tube following the completion of 6 cycles of PCR?

STEP-BY-STEP SOLUTION

Each cycle of PCR duplicates the number of template molecules present at the start of that cycle. The general equation for DNA amplification after n cycles is:

$$N = N_0 \times 2^n$$

Where:
$N_0 = 6$ (initial double-stranded DNA templates)
$n = 6$ (number of cycles completed)

$$N = 6 \times 2^6 = 6 \times 64 = 384 \text{ dsDNA molecules}$$

Answer: 384 double-stranded DNA molecules are generated.

Worked Example 2

ENZYME DIGESTION & GEL BAND COUNT

A linear double-stranded DNA molecule contains 3 recognition sites for the restriction endonuclease *Eco*RI. A circular plasmid DNA molecule also contains 3 recognition sites for *Eco*RI. If both molecules are completely digested with *Eco*RI, how many fragments will each produce?

STEP-BY-STEP SOLUTION

1. **Linear DNA Molecule:**
A linear molecule with k recognition sites cut completely by an endonuclease yields $(k + 1)$ fragments.
Number of fragments $= 3 + 1 = 4$.

2. **Circular Plasmid DNA Molecule:**
A closed circular molecule with k recognition sites cut completely yields exactly k fragments because each cleavage opens a segment of the continuous ring.
Number of fragments $= 3$.

Answer: The linear DNA produces 4 fragments; the circular plasmid produces 3 fragments.

BOSONANCE

6. NEET UG Exam-Style Practice Questions

Question 1

SINGLE-CORRECT MCQ

Stanley Cohen and Herbert Boyer constructed the first artificial recombinant DNA molecule in 1972 by isolating an antibiotic resistance gene from a native plasmid of which organism?

(1) *Escherichia coli*

(2) *Salmonella typhimurium*

(3) *Thermus aquaticus*

(4) *Agrobacterium tumefaciens*

CORRECT ANSWER: (2)

Explanation: Cohen and Boyer isolated the antibiotic resistance gene by cutting DNA from a native plasmid of *Salmonella typhimurium*. This piece was subsequently linked to a plasmid vector and transferred into *E. coli* for cloning and expression.

Question 2

SINGLE-CORRECT MCQ

In the nomenclature of the restriction endonuclease *Eco*RI, what do the letter 'R' and the Roman numeral 'I' designate?

(1) 'R' stands for restriction, and 'I' indicates the first discovered enzyme in bacteria.

(2) 'R' is derived from the strain RY 13, and 'I' indicates it was the first enzyme isolated from that strain.

(3) 'R' designates the species *rectale*, and 'I' indicates one single active catalytic site.

(4) 'R' denotes round colony morphology, and 'I' represents its position in gel electrophoresis.

CORRECT ANSWER: (2)

Explanation: In *Eco*RI:
• **E** = Genus *Escherichia*
• **co** = Species *coli*
• **R** = Strain RY 13
• **I** = Roman numeral indicating the order in which the enzyme was isolated from that particular strain of bacteria.

BOSONANCE

Question 3

MATCHING TYPE

Match the enzymes listed in Column I with their specific functional roles in rDNA technology in Column II:

Column I (Enzyme)	Column II (Function / Source)
A. Lysozyme	p. Degrades fungal cell walls
B. Chitinase	q. Thermostable enzyme for primer extension in PCR
C. *Taq* polymerase	r. Degrades bacterial peptidoglycan cell walls
D. DNA ligase	s. Catalyzes phosphodiester sealing of DNA ends

(1) A-r, B-p, C-q, D-s

(2) A-p, B-r, C-s, D-q

(3) A-r, B-q, C-p, D-s

(4) A-s, B-p, C-q, D-r

CORRECT ANSWER: (1)

Explanation:

- Lysozyme hydrolyzes bacterial cell walls (A-r).
- Chitinase breaks down the fungal chitinous wall (B-p).
- *Taq* polymerase, isolated from *Thermus aquaticus*, extends primers at $72^{\circ}\mathrm{C}$ during PCR (C-q).
- DNA ligase forms covalent phosphodiester bonds to join matching DNA ends (D-s).

Question 4

ASSERTION - REASON

Assertion (A): In agarose gel electrophoresis, smaller DNA fragments migrate farther toward the anode than larger fragments.
Reason (R): The agarose matrix provides a sieving effect where movement is inversely proportional to fragment size, while all DNA fragments carry a uniform negative charge.

(1) Both (A) and (R) are true, and (R) is the correct explanation of (A).

(2) Both (A) and (R) are true, but (R) is NOT the correct explanation of (A).

(3) (A) is true, but (R) is false.

(4) (A) is false, but (R) is true.

CORRECT ANSWER: (1)

Explanation: DNA fragments have a uniform net negative charge across their length due to the phosphate backbone; under an electric field, they are driven toward the positive electrode (anode). The agarose gel acts as a molecular sieve, allowing smaller fragments to travel through its pores with greater ease and distance than larger ones.

BOSONANCE

Question 5

TWO-STATEMENT QUESTION

Evaluate the following statements regarding cloning vectors:
Statement I: If a vector contains multiple recognition sites for a single restriction endonuclease, it complicates gene cloning by generating numerous fragments.
Statement II: The *rop* gene in plasmid pBR322 codes for proteins that confer resistance to the antibiotic ampicillin.

(1) Both Statement I and Statement II are correct.

(2) Both Statement I and Statement II are incorrect.

(3) Statement I is correct, but Statement II is incorrect.

(4) Statement I is incorrect, but Statement II is correct.

CORRECT ANSWER: (3)

Explanation: Statement I is correct: a vector should ideally have only one recognition site for a chosen enzyme to avoid fragmentation. Statement II is incorrect: the *rop* gene codes for proteins involved in the replication of the plasmid, whereas ampicillin resistance is coded by the amp^R gene.

Question 6

SINGLE-CORRECT MCQ

A researcher ligates a foreign DNA fragment into the *Bam*HI site of plasmid pBR322 and transforms recipient *E. coli* cells. What phenotype will distinguish the recombinant transformants from non-recombinant transformants?

(1) Growth on ampicillin medium, but no growth on tetracycline medium.

(2) Growth on tetracycline medium, but no growth on ampicillin medium.

(3) Inability to grow on both ampicillin and tetracycline media.

(4) Ability to grow robustly on both ampicillin and tetracycline media.

CORRECT ANSWER: (1)

Explanation: The *Bam*HI recognition site is located within the tet^R gene of pBR322. Insertion of foreign DNA inactivates the tet^R gene (insertional inactivation) while leaving the amp^R gene intact. Consequently, recombinant transformants grow on ampicillin plates but cannot grow on tetracycline plates. Non-recombinants grow on both.

BOSONANCE

Question 7

SINGLE-CORRECT MCQ

In blue-white screening, what visual observation identifies recombinant bacterial colonies?

(1) Bright blue colonies resulting from functional β-galactosidase synthesis.

(2) Colorless/white colonies due to insertional inactivation of the β-galactosidase gene.

(3) Fluorescent orange colonies when exposed to UV light without staining.

(4) Deep red colonies due to lysis of host cell membranes.

CORRECT ANSWER: (2)

Explanation: In the presence of a chromogenic substrate (X-gal), non-recombinants produce active β-galactosidase and form blue colonies. Recombinants carry a foreign insert within the β-galactosidase coding sequence, leading to insertional inactivation and white/colorless colonies.

Question 8

SINGLE-CORRECT MCQ

Which of the following methods is specifically suited for introducing recombinant DNA into plant cells by bombarding them with micro-particles of gold or tungsten?

(1) Micro-injection

(2) Biolistics (Gene Gun)

(3) Divalent calcium heat shock

(4) Disarmed retroviral transduction

CORRECT ANSWER: (2)

Explanation: Biolistics or the gene gun method bombards plant cells with high-velocity micro-particles of gold or tungsten coated with DNA, effectively penetrating the rigid cellulosic plant cell wall. Micro-injection is primarily used for animal cells.

BOSONANCE

Question 9

SINGLE-CORRECT MCQ

During the isolation of pure DNA from a biological sample, what is the role of adding chilled ethanol?

(1) To denature double-stranded DNA into single strands.

(2) To selectively digest cellular ribonucleic acids and proteins.

(3) To precipitate out purified DNA as fine threads for spooling.

(4) To provide Mg^{2+} cofactor ions for restriction enzymes.

CORRECT ANSWER: (3)

Explanation: DNA is insoluble in ice-cold ethanol. Adding chilled ethanol causes pure DNA to precipitate out of aqueous solution, forming fine white threads that can be wound onto a glass rod (spooling).

Question 10

SINGLE-CORRECT MCQ

Why is a continuous culture system preferred over a closed batch system for large-scale production of recombinant proteins in bioreactors?

(1) It allows cells to reach the stationary phase rapidly to conserve nutrients.

(2) It drains used medium and continuously adds fresh medium to maintain cells in their physiologically most active exponential (log) phase.

(3) It eliminates the need for aeration and foam control.

(4) It completely avoids the need for downstream processing.

CORRECT ANSWER: (2)

Explanation: In a continuous culture, spent medium is removed from one side while fresh medium is supplied from the other. This dynamic equilibrium keeps cells in their active exponential (log) growth phase, maximizing biomass accumulation and recombinant product yield.

BOSONANCE

7. Common Pitfalls and High-Yield Revision Checklist

NEET TRAP WARNINGS & HIGH-FREQUENCY DISTINCTIONS

- **First Restriction Endonuclease Discovered vs. Characterized:** The enzymes restricting bacteriophage growth were discovered in 1963 in *E. coli*. However, the first sequence-specific endonuclease characterized was ***Hind* II** (six base-pair specificity), isolated five years later.
- **Agarose Gel Loading & Direction:** DNA is loaded into wells positioned near the **negative cathode** and migrates toward the **positive anode**. Smaller fragments migrate farther; the largest fragments remain closest to the wells.
- **pBR322 Restriction Sites:** Memorize the locations!
  - amp^R **contains:** *Pst* I and *Pvu* I.
  - tet^R **contains:** *Bam*H I and *Sal* I.

- rop **contains:** *Pvu* II.
- **Outside markers:** *Eco*R I, *Cla* I, *Hind* III.

- **Downstream Processing Limits:** Downstream processing consists strictly of **separation**, **purification**, **preservative formulation**, and **quality control**. Gene cloning, transformation, and biomass fermentation belong to upstream processing.

Mastery Checklist

- ✓ **EFB Definition:** Integration of natural science and organisms, cells, parts thereof, and molecular analogues.
- ✓ **1972 Cohen & Boyer:** First recombinant DNA constructed using *Salmonella typhimurium* plasmid and expressed in *E. coli*.
- ✓ **EcoRI Palindrome:** $5'$---GAATTC---$3'$ cut between G and A, leaving $5'$ overhanging sticky ends.
- ✓ **Gel Visualization:** Ethidium bromide + UV radiation yields bright orange bands; bands are recovered via elution.
- ✓ **pBR322:** Insertional inactivation at *Bam*HI produces resistance to ampicillin but sensitivity to tetracycline.
- ✓ **PCR Steps:** Denaturation (94°C) $\rightarrow$ Annealing (50–60°C) $\rightarrow$ Extension (72°C) with *Taq* polymerase ($2^{30} \approx 10^9$ copies).
- ✓ **Gene Delivery:** *Agrobacterium tumefaciens* (Ti plasmid / T-DNA) for plants, disarmed retroviruses for animals, gene gun (gold/tungsten) for plant bombardment.
- ✓ **Cell Wall Lysis:** Lysozyme (bacteria), Cellulase (plants), Chitinase (fungi); precipitation achieved with chilled ethanol.

8. Concise Chapter Summary

Modern biotechnology rests upon two foundational disciplines: **genetic engineering** (altering nucleic acid sequences to generate novel host phenotypes) and **bioprocess engineering** (maintaining sterile conditions for industrial cultivation).

Constructing recombinant DNA requires key molecular tools: **restriction endonucleases** that cleave at specific palindromic sequences to yield sticky ends, **DNA ligase** to covalently seal phosphodiester bonds, **vectors** (such as pBR322, Ti plasmids, or retroviruses) equipped with an *ori*, selectable markers, and unique cloning sites, and a **competent host** prepared by divalent cation treatment, heat shock, micro-injection, or biolistics.

The complete recombinant workflow begins with cell lysis (using lysozyme, cellulase, or chitinase) and chilled ethanol spooling, followed by sequence-specific restriction digestion, gel electrophoretic resolution and elution, enzymatic amplification of the target gene via **PCR**, ligation into a cloning vector, host transformation, high-density cultivation in **stirred-tank bioreactors**, and final recovery via **downstream processing**.

BOSONANCE