NEET UG — BIOLOGY TUTORIAL

Animal Kingdom

Syllabus Focus: Fundamental basis of animal classification (levels of organisation, symmetry, germinal layers, coelom, segmentation, notochord) and salient features with examples of Non-Chordate phyla (Porifera to Hemichordata) and Chordate groups (Urochordata, Cephalochordata, and Vertebrata classes from Cyclostomata to Mammalia).

---

1. Learning Objectives & Overview

Over a million species of animals have been described till now. Because animals exhibit vast differences in structure and form, classification is essential both to study them systematically and to assign a proper systematic position to newly described species.

By completing this tutorial, you will be able to:

- Identify and apply the six fundamental criteria used to classify members of Kingdom Animalia: levels of organisation, body symmetry, embryonic germ layers, nature of coelom, metameric segmentation, and notochord.
- Distinguish between incomplete and complete digestive tracts, open and closed circulatory systems, and direct and indirect development.
- Compare all eleven major animal phyla (Porifera through Chordata) on the basis of diagnostic features, organ systems, and reproduction.
- Differentiate between protochordates (Urochordata and Cephalochordata) and vertebrates, and compare the seven classes of Vertebrata (Cyclostomata, Chondrichthyes, Osteichthyes, Amphibia, Reptilia, Aves, and Mammalia).
- Recall exact scientific names and common names of representative organisms for high-yield NEET UG questions.

2. Basis of Classification

Despite differences in external form and structure, animals share fundamental features in relation to the arrangement of cells, body symmetry, nature of coelom, and patterns of digestive, circulatory, or reproductive systems. These features form the basis of animal classification.

2.1 Levels of Organisation

Although all members of Kingdom Animalia are multicellular, they do not exhibit the same pattern of cell organisation:

- **Cellular Level of Organisation:** Cells are arranged as loose cell aggregates with some division of labour (activities) among them. *Example:* Sponges (Phylum Porifera).
- **Tissue Level of Organisation:** Cells performing the same function are arranged into tissues. *Examples:* Coelenterates (Cnidaria) and Ctenophores.
- **Organ Level of Organisation:** Tissues are grouped together to form organs, each specialised for a particular function. *Example:* Platyhelminthes (flatworms) and higher phyla.
- **Organ-System Level of Organisation:** Organs associate to form functional systems, each concerned with a specific physiological function. *Examples:* Aschelminthes, Annelids, Arthropods, Molluscs, Echinoderms, Hemichordates, and Chordates.

Complexities in Organ Systems


- **Digestive System:**
  - *Incomplete digestive system:* Has only a single opening to the outside of the body that serves as both mouth and anus (e.g., Coelenterata, Ctenophora, Platyhelminthes).
  - *Complete digestive system:* Has two openings: a mouth and an anus (from Aschelminthes to Chordates).
- **Circulatory System:**
  - *Open type:* Blood is pumped out of the heart and cells and tissues are directly bathed in it (e.g., Arthropoda, Mollusca, Hemichordata, Urochordata).
  - *Closed type:* Blood is circulated through a series of vessels of varying diameters—arteries, veins, and capillaries (e.g., Annelida, Chordata).

2.2 Body Symmetry

Animals can be categorised based on how their bodies can be divided by planes passing through their centre:

- **Asymmetrical:** Any plane that passes through the centre does not divide the organism into equal halves. *Example:* Sponges (mostly asymmetrical).
- **Radial Symmetry:** Any plane passing through the central axis of the body divides the organism into two identical halves. *Examples:* Coelenterates, Ctenophores, and adult Echinoderms.

- **Bilateral Symmetry:** The body can be divided into identical left and right halves in only one plane. *Examples:* Platyhelminthes, Aschelminthes, Annelids, Arthropods, Molluscs, Echinoderm larvae, Hemichordates, and Chordates.



Tutorial Diagram 1: Comparison of Asymmetrical, Radial, and Bilateral Body Plans


2.3 Diploblastic and Triploblastic Organisation

- **Diploblastic Animals:** Cells are arranged in two embryonic layers—an external **ectoderm** and an internal **endoderm**. An undifferentiated, non-cellular layer called **mesoglea** is present between the ectoderm and endoderm. *Examples:* Coelenterates (Cnidaria) and Ctenophores.
- **Triploblastic Animals:** The developing embryo has a third cellular germinal layer, the **mesoderm**, located between the ectoderm and endoderm. *Examples:* Platyhelminthes to Chordates.

2.4 Coelom (Body Cavity)

The presence or absence of a cavity between the body wall and the gut wall is a crucial classification criterion. A true **coelom** is a body cavity lined by mesoderm.

- **Coelomates (Eucoelomates):** Animals possessing a true body cavity lined on all sides by mesoderm. *Examples:* Annelids, Molluscs, Arthropods, Echinoderms, Hemichordates, and Chordates.
- **Pseudocoelomates:** The body cavity is not lined by mesoderm; instead, the mesoderm is present as **scattered pouches** between the ectoderm and endoderm. Such a body cavity is called a **pseudocoelom**. *Example:* Aschelminthes (roundworms).
- **Acoelomates:** Animals in which the body cavity is completely absent. *Examples:* Porifera, Coelenterata, Ctenophora, and Platyhelminthes (flatworms are triploblastic acoelomates).



Tutorial Diagram 2: Germinal Layers and Sectional Views of Coelomate, Pseudocoelomate, and Acoelomate Body Plans


2.5 Segmentation

In some animals, the body is externally and internally divided into segments with a serial repetition of at least some organs. For example, in the earthworm, the body shows this pattern called **metameric segmentation**, and the phenomenon is known as **metamerism**. True segmentation is observed in Annelida, Arthropoda, and Chordata.

2.6 Notochord

The **notochord** is a mesodermally derived rod-like structure formed on the dorsal side during embryonic development in some animals:

- **Chordates:** Animals that possess a notochord at least during some stage of their embryonic development.
- **Non-chordates:** Animals that do not form a notochord (Phylum Porifera to Phylum Hemichordata).



Tutorial Diagram 3: Broad Classification of Kingdom Animalia Based on Common Fundamental Features

3. Classification of Animals: Non-Chordate Phyla

3.1 Phylum — Porifera (Sponges)

- **Habitat & Symmetry:** Generally marine (except *Spongilla*, which is a freshwater sponge) and mostly asymmetrical.
- **Level of Organisation:** Primitive multicellular animals with **cellular level** of organisation.
- **Water Transport or Canal System:** Water enters through minute pores called **ostia** in the body wall into a central cavity called the **spongocoel**, from where it goes out through a single large opening, the **osculum**:

$$\text{Exterior Water} \xrightarrow{\text{Ostia}} \text{Spongocoel} \xrightarrow{\text{Osculum}} \text{Exterior}$$

  This pathway is helpful in food gathering, respiratory exchange, and removal of waste.
- **Diagnostic Cells & Digestion: Choanocytes** (collar cells) line the spongocoel and the canals. Digestion is strictly **intracellular**.
- **Skeleton:** Body is supported by a skeleton made up of **spicules** or **spongin fibres**.
- **Reproduction & Development:** Sexes are not separate (**hermaphrodite** / monoecious), meaning eggs and sperms are produced by the same individual. Asexual reproduction occurs by fragmentation, and sexual reproduction by gamete formation. Fertilisation is **internal**, and development is **indirect** (having a larval stage morphologically distinct from the adult).

Examples: *Sycon* (Scypha), *Spongilla* (Freshwater sponge), and *Euspongia* (Bath sponge).


3.2 Phylum — Coelenterata (Cnidaria)

- **Habitat & Body Plan:** Aquatic, mostly marine, sessile or free-swimming, **radially symmetrical**, **diploblastic** animals with **tissue level** of organisation.
- **Cnidoblasts (Cnidocytes):** The phylum name is derived from cnidoblasts or cnidocytes (which contain stinging capsules called **nematocysts**) present on the tentacles and the body. They are used for **anchorage, defense, and capture of prey**.
- **Digestive Cavity:** They have a central **gastro-vascular cavity** with a single opening—the mouth located on an elevated **hypostome**. Digestion is both **extracellular and intracellular**.
- **Skeleton:** Some cnidarians, such as **corals**, have a skeleton composed of calcium carbonate ($\mathrm{CaCO_3}$).
- **Body Forms & Metagenesis:** Cnidarians exhibit two basic body forms:
  - **Polyp:** Sessile and cylindrical form (e.g., *Hydra*, *Adamsia*).
  - **Medusa:** Umbrella-shaped and free-swimming form (e.g., *Aurelia* or jellyfish).

  Cnidarians that exist in both body forms exhibit alternation of generations (**metagenesis**), where polyps produce medusae **asexually** and medusae form polyps **sexually** (e.g., *Obelia*):

$$\text{Polyp (Sessile, Cylindrical)} \underset{\text{Sexual (Gametes)}}{\overset{\text{Asexual}}{\rightleftharpoons}} \text{Medusa (Free-swimming, Umbrella-shaped)}$$

Examples: *Physalia* (Portuguese man-of-war), *Adamsia* (Sea anemone), *Pennatula* (Sea-pen), *Gorgonia* (Sea-fan), *Meandrina* (Brain coral), *Aurelia* (Jellyfish), *Obelia*, and *Hydra*.



(a) Medusa Form (e.g., Aurelia) **(b) Polyp Form (e.g., Adamsia)** **(c) Cnidoblast with Nematocyst**

Tutorial Diagram 4: Cnidarian Body Forms (Medusa and Polyp) and Structure of a Cnidoblast

3.3 Phylum — Ctenophora (Sea Walnuts or Comb Jellies)

- **Habitat & Body Plan: Exclusively marine**, radially symmetrical, diploblastic organisms with tissue level of organisation.
- **Locomotion:** The body bears **eight external rows of ciliated comb plates**, which help in locomotion.
- **Digestion:** Both extracellular and intracellular.
- **Bioluminescence:** The property of a living organism to emit light is well-marked in ctenophores.
- **Reproduction & Development:** Sexes are not separate (hermaphrodite). Reproduction takes place **only by sexual means**. Fertilisation is **external** with **indirect development**.

Examples: *Pleurobrachia* and *Ctenoplana*.

3.4 Phylum — Platyhelminthes (Flatworms)

- **Body Shape & Habitat:** Dorso-ventrally flattened body, hence called **flatworms**. Mostly endoparasites found in animals including human beings.
- **Body Plan:** Bilaterally symmetrical, **triploblastic**, and **acoelomate** animals with **organ level** of organisation (and incomplete digestive tract where present).
- **Parasitic Adaptations: Hooks and suckers** are present in parasitic forms. Some of them (e.g., tapeworms) absorb nutrients from the host directly through their body surface.
- **Excretion & Osmoregulation:** Specialised cells called **flame cells** help in osmoregulation and excretion.
- **Reproduction & Regeneration:** Sexes are not separate (hermaphrodite). Fertilisation is **internal** and development is indirect through **many larval stages**. Some members like *Planaria* possess **high regeneration capacity**.

Examples: *Taenia* (Tapeworm), *Fasciola* (Liver fluke), and *Planaria*.

3.5 Phylum — Aschelminthes (Roundworms)

- **Body Shape & Habitat:** Circular in cross-section, hence named **roundworms**. They may be free-living, aquatic and terrestrial, or parasitic in plants and animals.
- **Body Plan: Organ-system level** of body organisation, bilaterally symmetrical, triploblastic, and **pseudocoelomate** animals.
- **Digestive System:** Alimentary canal is **complete** with a well-developed **muscular pharynx**.
- **Excretory System:** An excretory tube removes body wastes from the body cavity through the **excretory pore**.
- **Reproduction & Sexual Dimorphism:** Sexes are separate (**dioecious**), meaning males and females are morphologically distinct. **Often females are longer than males** (and the male posterior end is curved). Fertilisation is **internal**, and development may be **direct** (the young ones resemble the adult) or **indirect**.

Examples: *Ascaris* (Roundworm), *Wuchereria* (Filaria worm), and *Ancylostoma* (Hookworm).

3.6 Phylum — Annelida

- **Habitat & Body Plan:** Aquatic (marine and freshwater) or terrestrial; free-living, and sometimes parasitic. They exhibit organ-system level of organisation, bilateral symmetry, and are triploblastic, **metamerically segmented**, and **coelomate** animals.
- **Segmentation:** Body surface is distinctly marked out into segments or **metameres** (Latin, *annulus*: little ring).
- **Locomotion:** Possess longitudinal and circular muscles that help in locomotion. Aquatic annelids like *Nereis* possess lateral appendages called **parapodia**, which help in swimming.
- **Circulatory & Excretory Systems:** A **closed circulatory system** is present. **Nephridia** (singular: nephridium) help in osmoregulation and excretion.
- **Neural System:** Consists of paired ganglia (singular: ganglion) connected by lateral nerves to a **double ventral nerve cord**.
- **Reproduction:** *Nereis* (an aquatic form) is **dioecious**, whereas earthworms and leeches are **monoecious**. Reproduction is sexual.

Examples: *Nereis*, *Pheretima* (Earthworm), and *Hirudinaria* (Blood-sucking leech).

3.7 Phylum — Arthropoda

- **Abundance:** The **largest phylum** of Animalia, which includes insects. Over **two-thirds of all named species** on earth are arthropods.
- **Body Plan & Exoskeleton:** Organ-system level of organisation; bilaterally symmetrical, triploblastic, segmented, and coelomate animals. The body is covered by a **chitinous exoskeleton** and consists of **head, thorax, and abdomen**. They possess **jointed appendages** (*arthros* = joint, *poda* = appendages).
- **Respiration & Circulation:** Respiratory organs are **gills, book gills, book lungs, or tracheal system**. Circulatory system is of the **open type**.
- **Sensory & Excretory Organs:** Sensory organs like antennae, eyes (compound and simple), and **statocysts** (balancing organs) are present. Excretion takes place through **Malpighian tubules**.
- **Reproduction:** Mostly dioecious. Fertilisation is usually internal. They are mostly oviparous, and development may be direct or indirect.

Examples:
• *Economically important insects: Apis* (Honey bee), *Bombyx* (Silkworm), *Laccifer* (Lac insect)
• *Vectors: Anopheles*, *Culex*, and *Aedes* (Mosquitoes)
• *Gregarious pest: Locusta* (Locust)
• *Living fossil: Limulus* (King crab)

3.8 Phylum — Mollusca

- **Abundance & Habitat:** The **second largest** animal phylum. Terrestrial or aquatic (marine or freshwater).
- **Body Plan:** Organ-system level of organisation; bilaterally symmetrical, triploblastic, and coelomate. The body is covered by a **calcareous shell** and is **unsegmented** with a distinct **head, muscular foot, and visceral hump**.
- **Mantle & Gills:** A soft and spongy layer of skin forms a **mantle** over the visceral hump. The space between the hump and the mantle is called the **mantle cavity**, in which **feather-like gills** are present. These gills perform both **respiratory and excretory functions**.
- **Sensory & Feeding Organs:** The anterior head region has sensory tentacles. The mouth contains a file-like rasping organ for feeding, called the **radula**.
- **Reproduction:** Usually dioecious and oviparous with indirect development.

Examples: *Pila* (Apple snail), *Pinctada* (Pearl oyster), *Sepia* (Cuttlefish), *Loligo* (Squid), *Octopus* (Devil fish), *Aplysia* (Sea-hare), *Dentalium* (Tusk shell), and *Chaetopleura* (Chiton).

3.9 Phylum — Echinodermata

- **Skeleton & Habitat:** Have an **endoskeleton of calcareous ossicles** and hence the name Echinodermata (spiny-bodied). **All are marine** with organ-system level of organisation.
- **Symmetry & Coelom:** Adult echinoderms are **radially symmetrical**, whereas their **larvae are bilaterally symmetrical**. They are triploblastic and coelomate animals.
- **Digestive System:** Complete with mouth on the lower (**ventral**) side and anus on the upper (**dorsal**) side.
- **Water Vascular System:** The most distinctive feature of echinoderms; it helps in **locomotion, capture and transport of food, and respiration**.
- **Excretion & Reproduction:** An **excretory system is absent**. Sexes are separate. Reproduction is sexual, fertilisation is usually **external**, and development is **indirect** with a free-swimming larva.

Examples: *Asterias* (Starfish), *Echinus* (Sea urchin), *Antedon* (Sea lily), *Cucumaria* (Sea cucumber), and *Ophiura* (Brittle star).

3.10 Phylum — Hemichordata

- **Systematic Position & Stomochord:** Earlier considered a subphylum under Phylum Chordata, now placed as a separate phylum under Non-chordata. Hemichordates possess a rudimentary structure in the collar region called the **stomochord**, a structure similar to the notochord.
- **Body Plan:** Small group of **worm-like marine** animals with organ-system level of organisation; bilaterally symmetrical, triploblastic, and coelomate.
- **Body Divisions:** Cylindrical body composed of an anterior **proboscis**, a **collar**, and a long **trunk**.
- **Organ Systems & Reproduction:** Circulatory system is of the **open type**. Respiration takes place through **gills**. The excretory organ is the **proboscis gland**. Sexes are separate, fertilisation is **external**, and development is **indirect**.


Examples: *Balanoglossus* and *Saccoglossus*.

4. Phylum — Chordata

Animals belonging to Phylum Chordata are fundamentally characterised by the presence of:

1. A **notochord**
2. A **dorsal hollow nerve cord**
3. **Paired pharyngeal gill slits**
4. A **post-anal tail**

Chordates are bilaterally symmetrical, triploblastic, and coelomate with an organ-system level of organisation and a **closed circulatory system**.



Tutorial Diagram 5: Body Divisions of Balanoglossus (Hemichordata) vs. Fundamental Diagnostic Features of Chordata

Comparison of Chordates and Non-Chordates


S.No.	Chordates	Non-chordates
1.	Notochord present.	Notochord absent.
2.	Central nervous system is **dorsal, hollow, and single**.	Central nervous system is **ventral, solid, and double**.
3.	Pharynx perforated by **gill slits**.	Gill slits are absent.
4.	Heart is **ventral**.	Heart is **dorsal** (if present).
5.	A **post-anal part (tail)** is present.	Post-anal tail is absent.

4.1 Subphyla of Phylum Chordata

Phylum Chordata is divided into three subphyla:

1. **Subphylum Urochordata (Tunicata):** Exclusively marine. The notochord is present **only in the larval tail**. *Examples: Ascidia*, *Salpa*, *Doliolum*.
2. **Subphylum Cephalochordata:** Exclusively marine. The notochord extends from **head to tail region** and is **persistent throughout their life**. *Example: Branchiostoma* (Amphioxus or Lancelet).
   Note: Subphyla Urochordata and Cephalochordata are together referred to as **protochordates** and are exclusively marine.
3. **Subphylum Vertebrata:** Possess a notochord during the **embryonic period**, which is **replaced by a cartilaginous or bony vertebral column** in the adult. Therefore, *all vertebrates are chordates, but all chordates are not vertebrates* (since protochordates

are chordates that lack a vertebral column). Besides basic chordate characters, vertebrates have a **ventral muscular heart** with two, three, or four chambers, **kidneys** for excretion and osmoregulation, and **paired appendages** (which may be fins or limbs).



Tutorial Diagram 6: Division of Subphylum Vertebrata into Agnatha and Gnathostomata

4.2 Classes of Subphylum Vertebrata

4.2.1 Class — Cyclostomata

- **Habit & Mouth:** All living members are **ectoparasites on some fishes**. They have a **sucking and circular mouth without jaws**.
- **Respiration & Integument:** Elongated body bearing **6–15 pairs of gill slits** for respiration. Their body is **devoid of scales and paired fins**.
- **Skeleton & Circulation:** Cranium and vertebral column are **cartilaginous**. Circulation is of the **closed type**.
- **Spawning Migration:** Cyclostomes are **marine** but migrate for spawning to **freshwater**. After spawning, within a few days, they die. Their larvae, after metamorphosis, return to the ocean.

Examples: *Petromyzon* (Lamprey) and *Myxine* (Hagfish).

4.2.2 Class — Chondrichthyes (Cartilaginous Fishes)

- **Habitat, Skeleton & Mouth:** Marine animals with streamlined body and **cartilaginous endoskeleton**. Mouth is located **ventrally**. **Notochord is persistent throughout life**.
- **Gills & Buoyancy:** Gill slits are separate and **without operculum** (gill cover). Due to the **absence of an air bladder**, they have to swim constantly to avoid sinking.
- **Skin, Teeth & Defense:** Skin is tough, containing minute **placoid scales**. Teeth are **modified placoid scales** which are backwardly directed; jaws are very powerful (predaceous animals). Some have **electric organs** (e.g., *Torpedo*) and some possess a **poison sting** (e.g., *Trygon*).
- **Heart & Thermoregulation:** Heart is **two-chambered** (one auricle and one ventricle). They are cold-blooded (**poikilothermous**) animals—they lack the capacity to regulate their body temperature.
- **Reproduction:** Sexes are separate. In males, **pelvic fins bear claspers**. They have **internal fertilisation** and many of them are **viviparous**.

Examples: *Scoliodon* (Dogfish), *Pristis* (Sawfish), *Carcharodon* (Great white shark), *Trygon* (Stingray), and *Torpedo* (Electric ray).

4.2.3 Class — Osteichthyes (Bony Fishes)

- **Habitat, Skeleton & Mouth:** Includes both marine and freshwater fishes with a **bony endoskeleton**. Streamlined body with a mostly **terminal mouth**.
- **Gills, Scales & Buoyancy:** Possess **four pairs of gills** covered by an **operculum** on each side. Skin is covered with **cycloid/ctenoid scales**. An **air bladder is present** which regulates buoyancy.
- **Heart & Thermoregulation:** Heart is **two-chambered** (one auricle and one ventricle). Cold-blooded (poikilothermous).
- **Reproduction:** Sexes are separate. Fertilisation is usually **external**. Mostly **oviparous** and development is **direct**.

Examples:
• *Marine: Exocoetus* (Flying fish), *Hippocampus* (Sea horse)
• *Freshwater: Labeo* (Rohu), *Catla* (Katla), *Clarias* (Magur)
• *Aquarium: Betta* (Fighting fish), *Pterophyllum* (Angel fish)

4.2.4 Class — Amphibia

- **Habitat & Body Form:** Can live in aquatic as well as terrestrial habitats (Greek, *Amphi* = dual, *bios* = life). Most have two pairs of limbs. Body is divisible into **head and trunk** (tail may be present in some, such as *Salamandra*).
- **Skin, Eyes & Ear:** Skin is **moist (without scales)**. Eyes have eyelids. A **tympanum** represents the ear.
- **Cloaca:** Alimentary canal, urinary, and reproductive tracts open into a common chamber called the **cloaca**, which opens to the exterior.
- **Respiration, Heart & Thermoregulation:** Respiration is by **gills, lungs, and through skin**. Heart is **three-chambered** (two auricles and one ventricle). Cold-blooded (poikilothermous).
- **Reproduction:** Sexes are separate. Fertilisation is **external**. They are **oviparous** and development is **indirect**.

Examples: *Bufo* (Toad), *Rana* (Frog), *Hyla* (Tree frog), *Salamandra* (Salamander), and *Ichthyophis* (Limbless amphibia).

4.2.5 Class — Reptilia

- **Locomotion & Skin:** Named for their creeping or crawling mode of locomotion (Latin, *repere* or *reptum* = to creep or crawl). Mostly terrestrial; body is covered by **dry and cornified skin, epidermal scales or scutes**. Snakes and lizards shed their scales as **skin cast**.
- **Ear & Limbs:** Do not have external ear openings; **tympanum** represents the ear. Limbs, when present, are two pairs (absent in snakes).
- **Heart & Thermoregulation:** Heart is usually **three-chambered**, but **four-chambered in crocodiles**. Reptiles are **poikilotherms**.
- **Reproduction:** Sexes are separate. Fertilisation is **internal**. They are **oviparous** and development is **direct**.

Examples: *Chelone* (Turtle), *Testudo* (Tortoise), *Chameleon* (Tree lizard), *Calotes* (Garden lizard), *Crocodilus* (Crocodile), *Alligator* (Alligator), *Hemidactylus* (Wall lizard); *Poisonous snakes: Naja* (Cobra), *Bangarus* (Krait), *Vipera* (Viper).

4.2.6 Class — Aves (Birds)

- **Diagnostic Features & Limbs:** Presence of **feathers** and a **beak**. Most can fly except flightless birds (e.g., Ostrich). **Forelimbs are modified into wings**. Hind limbs generally have **scales** and are modified for walking, swimming, or clasping tree branches.
- **Skin & Endoskeleton:** Skin is **dry without glands** except the **oil gland (preen gland) at the base of the tail**. Endoskeleton is fully ossified (bony) and the long bones are hollow with air cavities (**pneumatic bones**).
- **Digestion & Respiration:** Digestive tract has additional chambers—the **crop and gizzard**. Respiration is by **lungs**, and **air sacs connected to lungs supplement respiration**.
- **Heart & Thermoregulation:** Heart is **completely four-chambered**. They are warm-blooded (**homoiothermous**) animals—able to maintain a constant body temperature.
- **Reproduction:** Sexes are separate. Fertilisation is **internal**. They are **oviparous** and development is **direct**.

Examples: *Corvus* (Crow), *Columba* (Pigeon), *Psittacula* (Parrot), *Struthio* (Ostrich), *Pavo* (Peacock), *Aptenodytes* (Penguin), and *Neophron* (Vulture).

4.2.7 Class — Mammalia

- **Habitats:** Found in polar ice caps, deserts, mountains, forests, grasslands, and dark caves; some have adapted to fly (*Pteropus*) or live in water (*Delphinus*, *Balaenoptera*).
- **Unique Mammalian Features:** Presence of milk-producing glands (**mammary glands**) by which young ones are nourished, **hair** on the skin, and external ears or **pinnae**.
- **Limbs & Dentition:** Two pairs of limbs adapted for walking, running, climbing, burrowing, swimming, or flying. **Different types of teeth** (heterodont dentition) are present in the jaw.
- **Heart, Respiration & Thermoregulation:** Heart is **four-chambered**. Respiration is by **lungs**. They are **homoiothermous** (warm-blooded).
- **Reproduction:** Sexes are separate, fertilisation is **internal**. They are **viviparous with few exceptions** (Platypus is oviparous) and development is **direct**.

Examples:
• *Oviparous: Ornithorhynchus* (Platypus)
• *Viviparous: Macropus* (Kangaroo), *Pteropus* (Flying fox), *Camelus* (Camel), *Macaca* (Monkey), *Rattus* (Rat), *Canis* (Dog), *Felis* (Cat), *Elephas* (Elephant), *Equus* (Horse), *Delphinus* (Common dolphin), *Balaenoptera* (Blue whale), *Panthera tigris* (Tiger), *Panthera leo* (Lion).

5. Master Comparison Tables

The following table summarises the salient features of all eleven phyla in the Animal Kingdom for rapid comparative revision:

Phylum	Level of Organisation	Symmetry	Coelom	Segmentation	Digestive System	Circulatory System	Respiratory System	Distinctive Features
Porifera	Cellular	Various (Mostly asymmetrical)	Absent	Absent	Absent	Absent	Absent	Body with pores and canals in walls; choanocytes.
Coelenterata (Cnidaria)	Tissue	Radial	Absent	Absent	Incomplete	Absent	Absent	Cnidoblasts present; polyp and medusa forms.
Ctenophora	Tissue	Radial	Absent	Absent	Incomplete	Absent	Absent	Comb plates for locomotion; bioluminescence.
Platyhelminthes	Organ & Organ-system	Bilateral	Absent	Absent	Incomplete	Absent	Absent	Flat body, hooks and suckers, flame cells.
Aschelminthes	Organ-system	Bilateral	Pseudocoelomate	Absent	Complete	Absent	Absent	Often worm-shaped, elongated; muscular pharynx.
Annelida	Organ-system	Bilateral	Coelomate	Present	Complete	Present (Closed)	Absent	Body segmentation like rings; nephridia, parapodia.
Arthropoda	Organ-system	Bilateral	Coelomate	Present	Complete	Present (Open)	Present	Exoskeleton of cuticle/chitin, jointed appendages.
Mollusca	Organ-system	Bilateral	Coelomate	Absent	Complete	Present	Present	External calcareous shell usually present; radula, mantle.
Echinodermata	Organ-system	Radial (Adult), Bilateral (Larva)	Coelomate	Absent	Complete	Present	Present	Water vascular system, calcareous endoskeleton.
Hemichordata	Organ-system	Bilateral	Coelomate	Absent	Complete	Present (Open)	Present (Gills)	Worm-like with proboscis, collar, and trunk; stomochord.
Chordata	Organ-system	Bilateral	Coelomate	Present	Complete	Present (Closed)	Present	Notochord, dorsal hollow nerve cord, gill slits, limbs or fins.

6. Conceptual Connections & Worked Examples

Understanding how classification principles connect to physiology and morphology is essential for solving analytical questions. Study the following worked examples carefully.

Worked Example 1: Systematic Procedure to Classify an Unknown Specimen

Problem: Why are common fundamental features necessary for animal classification, and what step-by-step procedure would you follow if given an unknown animal specimen to classify?

Solution:

Without common fundamental features, it would be impossible to group over a million diverse animal species logically, study their evolutionary relationships, or assign a systematic position to newly discovered species. Every organism would have to be studied in isolation.

Step-by-Step Classification Procedure:

1. **Determine the Level of Organisation:** Check whether cells form loose aggregates (Cellular $\rightarrow$ Porifera), tissues (Tissue $\rightarrow$ Coelenterata/Ctenophora), organs (Organ $\rightarrow$ Platyhelminthes), or functional organ systems (Aschelminthes to Chordata).
2. **Determine Body Symmetry:** Observe if the body is asymmetrical, radially symmetrical, or bilaterally symmetrical.

3. **Examine Germinal Layers & Coelom:** Identify if the specimen is diploblastic or triploblastic, and whether the body cavity is absent (acoelomate), a pseudocoelom (scattered mesodermal pouches), or a true mesoderm-lined coelom (coelomate).
4. **Check for Segmentation & Notochord:** Determine if true metameric segmentation is present (Annelida, Arthropoda, Chordata) and whether a dorsal rod-like notochord is formed during development (Chordata vs. Non-chordata).
5. **Identify Phylum-Specific Diagnostic Features:** Look for unique traits such as jointed appendages (Arthropoda), calcareous shell and radula (Mollusca), water vascular system (Echinodermata), or vertebral column (Vertebrata).

Worked Example 2: Key Physiological & Developmental Distinctions

Problem: Distinguish clearly between: (a) Intracellular and extracellular digestion, (b) Direct and indirect development, and (c) Oviparous and viviparous reproduction in terms of the number of young ones produced.

Solution:

- **(a) Intracellular vs. Extracellular Digestion:** In *intracellular digestion*, food particles are engulfed by cells and digested inside food vacuoles within the cell (characteristic of Porifera). In *extracellular digestion*, digestive enzymes are secreted into a cavity (such as the gastro-vascular cavity or alimentary canal lumen) where food is broken down outside the cells. Coelenterates and Ctenophores exhibit both extracellular and intracellular digestion, whereas higher phyla have predominantly extracellular digestion in a digestive tract.
- **(b) Direct vs. Indirect Development:** In *direct development*, there is no larval stage and the newly hatched or born young ones morphologically resemble the adult (e.g., Earthworm, Reptiles, Birds, Mammals). In *indirect development*, embryonic development includes one or more morphologically distinct larval stages that undergo metamorphosis to become adults (e.g., Porifera, Cnidaria, Ctenophora, Platyhelminthes, Mollusca, Echinodermata, Hemichordata, Amphibia).
- **(c) Number of Eggs/Young Ones in Oviparous vs. Viviparous Mothers:** The number of eggs produced by an oviparous mother and the number of young ones produced by a viviparous mother are **not equal**. An oviparous mother lays eggs in the external environment where developing embryos face high risks of predation and harsh environmental conditions; hence, a large number of eggs are produced to ensure survival of the species (especially in fishes and amphibians). In contrast, a viviparous mother retains and nourishes the embryo inside the maternal body, providing high parental protection and care, so fewer young ones are produced at a time.

Worked Example 3: Structural & Functional Adaptations Across Animal Groups

Problem: Explain the biological basis for the following observations:
(1) Arthropods constitute the largest group of the animal kingdom.
(2) Presence of an air bladder is important in Osteichthyes compared to Chondrichthyes.
(3) Birds show specific structural modifications that enable flight.
(4) Certain animals are adapted to live as endoparasites in human beings.

Solution:

1. **Success of Arthropoda:** Arthropods constitute over two-thirds of all named species because of three major adaptations: (i) a tough **chitinous exoskeleton** that prevents desiccation and provides protection on land, (ii) **jointed appendages** that allow diverse modes of locomotion, feeding, and defense, and (iii) diverse respiratory structures (gills, book gills, book lungs, tracheal system) and well-developed sensory organs (compound/simple eyes, antennae, statocysts) enabling survival in almost every habitat.
2. **Role of Air Bladder in Pisces:** In bony fishes (Osteichthyes), the **air bladder regulates buoyancy**, allowing the fish to stay at a particular water depth without expending energy on swimming. Cartilaginous fishes (Chondrichthyes) lack an air bladder and therefore must **swim constantly to avoid sinking**.
3. **Flight Modifications in Birds (Aves):** (i) Forelimbs are modified into **wings**, (ii) Body is covered with **feathers** and streamlined, (iii) Long bones are hollow with air cavities (**pneumatic bones**), reducing body weight, and (iv) **Air sacs** connected to the lungs supplement respiration to meet the high oxygen demand of flight.
4. **Human Parasites & Adaptations:** Common human endoparasites include *Taenia* (Tapeworm) and *Fasciola* (Liver fluke) from Platyhelminthes, and *Ascaris* (Roundworm), *Wuchereria* (Filaria worm), and *Ancylostoma* (Hookworm) from Aschelminthes. Parasitic flatworms possess **hooks and suckers** for attachment to the host and can absorb pre-digested nutrients directly through their body surface.

7. NEET UG Practice Questions with Worked Solutions

Note: The following are original exam-aligned practice questions designed according to the NEET UG standard to test conceptual clarity, factual accuracy, and analytical reasoning.

Practice Question 1 (Single-Choice MCQ)

Which of the following animal groups is correctly paired with its level of organisation, body symmetry, and nature of coelom?

(1) Platyhelminthes — Organ level, Bilateral symmetry, Pseudocoelomate

(2) Aschelminthes — Organ-system level, Bilateral symmetry, Pseudocoelomate

(3) Ctenophora — Tissue level, Bilateral symmetry, Acoelomate

(4) Echinodermata (Adult) — Organ-system level, Bilateral symmetry, Coelomate

Correct Answer: (2)
Worked Solution: Aschelminthes (roundworms) exhibit an organ-system level of body organisation, bilateral symmetry, triploblastic germ layers, and a pseudocoelom (body cavity with scattered pouches of mesoderm). Platyhelminthes are acoelomates (not pseudocoelomates). Ctenophores exhibit radial symmetry. Adult echinoderms are radially symmetrical (only their larvae are bilaterally symmetrical).

Practice Question 2 (Single-Choice MCQ)

Water vascular system is the most distinctive characteristic feature of which of the following phyla, and what are its primary functions?

(1) Porifera — Food gathering, respiratory exchange, and removal of waste

(2) Ctenophora — Locomotion and bioluminescence

(3) Echinodermata — Locomotion, capture and transport of food, and respiration

(4) Mollusca — Filter feeding and osmoregulation

Correct Answer: (3)
Worked Solution: The **water vascular system** is the distinctive feature of Phylum **Echinodermata** (e.g., *Asterias*, *Echinus*) and functions in locomotion, capture and transport of food, and respiration. Do not confuse this with the **water transport or canal system** of Phylum **Porifera** (ostia $\rightarrow$ spongocoel $\rightarrow$ osculum), which functions in food gathering, respiratory exchange, and removal of waste.

Practice Question 3 (Single-Choice MCQ)

True metameric segmentation (metamerism) in the body is first observed in the evolutionary hierarchy in which of the following phyla?

(1) Platyhelminthes

(2) Aschelminthes

(3) Annelida

(4) Mollusca

Correct Answer: (3)
Worked Solution: True metameric segmentation—where the body is externally and internally divided into segments (metameres) with a serial repetition of at least some organs—is first observed in Phylum **Annelida** (e.g., *Pheretima* / earthworm). It is also present in Arthropoda and Chordata, but absent in Platyhelminthes, Aschelminthes, and Mollusca.

Practice Question 4 (Matching Type MCQ)

Match the diagnostic structures in Column I with their corresponding animal groups in Column II and select the correct option:

Column I (Structure)	Column II (Taxon)
(a) Operculum	(i) Ctenophora
(b) Parapodia	(ii) Mollusca
(c) Epidermal scales / scutes	(iii) Porifera
(d) Comb plates	(iv) Reptilia
(e) Radula	(v) Annelida
(f) Hairs and pinnae	(vi) Cyclostomata and Chondrichthyes
(g) Choanocytes	(vii) Mammalia
(h) Uncovered gill slits	(viii) Osteichthyes

(1) (a)-(viii), (b)-(v), (c)-(iv), (d)-(i), (e)-(ii), (f)-(vii), (g)-(iii), (h)-(vi)

(2) (a)-(vi), (b)-(v), (c)-(iv), (d)-(i), (e)-(ii), (f)-(vii), (g)-(iii), (h)-(viii)

(3) (a)-(viii), (b)-(ii), (c)-(iv), (d)-(iii), (e)-(v), (f)-(vii), (g)-(i), (h)-(vi)

(4) (a)-(viii), (b)-(v), (c)-(vii), (d)-(i), (e)-(ii), (f)-(iv), (g)-(iii), (h)-(vi)

Correct Answer: (1)
Worked Solution:
• (a) **Operculum** covers the four pairs of gills in bony fishes (**Osteichthyes**, viii).
• (b) **Parapodia** are lateral swimming appendages in aquatic **Annelida** like *Nereis* (v).
• (c) **Scales/scutes** cover the dry, cornified skin of **Reptilia** (iv).
• (d) Eight external rows of ciliated **comb plates** aid locomotion in **Ctenophora** (i).
• (e) **Radula** is the file-like rasping feeding organ in **Mollusca** (ii).
• (f) **Hairs** on skin and external ears (pinnae) are unique to **Mammalia** (vii).
• (g) **Choanocytes** (collar cells) line the spongocoel and canals in **Porifera** (iii).
• (h) Separate **gill slits** without an operculum occur in **Cyclostomata and Chondrichthyes** (vi).

Practice Question 5 (Multiple-Statement Evaluation)

Read the following statements regarding Phylum Coelenterata (Cnidaria) and Phylum Ctenophora:

I. In *Obelia*, polyps produce medusae sexually and medusae form polyps asexually.
II. Both Cnidarians and Ctenophores are radially symmetrical, diploblastic animals with tissue level of organisation and both extracellular and intracellular digestion.
III. Ctenophores reproduce both asexually by fragmentation and sexually by gamete formation.
IV. Corals belong to Phylum Cnidaria and possess a skeleton composed of calcium carbonate.

Which of the above statements are **correct**?

(1) I and III only

(2) II and IV only

(3) I, II, and IV only

(4) II, III, and IV only

Correct Answer: (2)
Worked Solution:
• Statement I is **incorrect**: In metagenesis (e.g., *Obelia*), polyps produce medusae **asexually** and medusae form the polyps **sexually**.
• Statement II is **correct**: Both Cnidaria and Ctenophora exhibit tissue level of organisation, radial symmetry, diploblastic germ layers, an incomplete digestive tract, and both extracellular and intracellular digestion.
• Statement III is **incorrect**: Reproduction in Ctenophora takes place **only by sexual means**.
• Statement IV is **correct**: Corals (such as *Meandrina*, brain coral) are cnidarians with a calcium carbonate skeleton.

Practice Question 6 (Assertion–Reason Type)

Assertion (A): All vertebrates are chordates, but all chordates are not vertebrates.
Reason (R): All chordates possess a notochord at some stage of development, Technology/Evolutionary replacement of the embryonic notochord by a cartilaginous or bony vertebral column occurs only in subphylum Vertebrata and not in protochordates.

(1) Both (A) and (R) are true and (R) is the correct explanation of (A).

(2) Both (A) and (R) are true, but (R) is not the correct explanation of (A).

(3) (A) is true, but (R) is false.

(4) (A) is false, but (R) is true.

Correct Answer: (1)
Worked Solution: Phylum Chordata includes three subphyla: Urochordata, Cephalochordata, and Vertebrata. All members possess a notochord (defining them as chordates). However, in Urochordata (notochord only in larval tail) and Cephalochordata (notochord persistent from head to tail throughout life), a vertebral column never forms. Only in subphylum Vertebrata is the embryonic notochord replaced by a cartilaginous or bony vertebral column in the adult. Hence, both A and R are true and R correctly explains A.

Practice Question 7 (Single-Choice MCQ)

Compare the three worm phyla—Platyhelminthes, Aschelminthes, and Annelida—and select the statement that is **completely accurate**:

(1) *Taenia*, *Ascaris*, and *Pheretima* are all monoecious (hermaphrodite) animals.

(2) Flame cells in *Fasciola* and nephridia in *Hirudinaria* both perform osmoregulation and excretion.

(3) All three phyla possess a complete alimentary canal with a separate mouth and anus.

(4) *Nereis* is monoecious, whereas earthworms and leeches are dioecious.

Correct Answer: (2)
Worked Solution: Specialised flame cells in Platyhelminthes (e.g., *Fasciola*) and nephridia in Annelida (e.g., *Hirudinaria*) both help in osmoregulation and excretion. Option (1) is incorrect because *Ascaris* (Aschelminthes) is dioecious (sexes are separate). Option (3) is incorrect because Platyhelminthes have an incomplete digestive system. Option (4) is reversed: *Nereis* is dioecious, whereas earthworms and leeches are monoecious.

Practice Question 8 (Single-Choice MCQ)

Which of the following pairs of animals is correctly matched with its biological category and phylum?

(1) *Limulus* (King crab) — Living fossil belonging to Phylum Mollusca

(2) *Laccifer* and *Bombyx* — Gregarious pests belonging to Phylum Arthropoda

(3) *Pinctada* (Pearl oyster) and *Loligo* (Squid) — Unsegmented coelomates with a radula belonging to Phylum Mollusca

(4) *Antedon* (Sea lily) and *Cucumaria* (Sea cucumber) — Bilaterally symmetrical adults with a proboscis gland belonging to Phylum Echinodermata

Correct Answer: (3)
Worked Solution: *Pinctada* (Pearl oyster) and *Loligo* (Squid) belong to Phylum Mollusca, which are unsegmented, bilaterally symmetrical coelomates possessing a file-like rasping organ (radula) and mantle cavity. Option (1) is wrong because *Limulus* (King crab) is a living fossil in Phylum **Arthropoda**. Option (2) is wrong because *Laccifer* (Lac insect) and *Bombyx* (Silkworm) are **economically important insects**, whereas *Locusta* (Locust) is a gregarious pest. Option (4) is wrong because adult echinoderms are radially symmetrical and lack an excretory system (proboscis gland belongs to Hemichordata).

Practice Question 9 (Single-Choice MCQ)

A marine biologist collects an animal specimen with a cylindrical body divisible into an anterior proboscis, a collar, and a long trunk. Which of the following sets of features will also be present in this organism?

(1) Stomochord in the collar region, open circulatory system, respiration by gills, and proboscis gland for excretion

(2) Persistent notochord from head to tail, closed circulatory system, and flame cells for excretion

(3) Notochord restricted to the larval tail, water vascular system, and internal fertilisation

(4) Dorsal solid nerve cord, ventral heart, and direct development

Correct Answer: (1)
Worked Solution: A worm-like marine animal with a cylindrical body divided into proboscis, collar, and trunk belongs to Phylum **Hemichordata** (e.g., *Balanoglossus*, *Saccoglossus*). Hemichordates have a rudimentary structure in the collar region called the **stomochord**, an **open circulatory system**, **gill respiration**, a **proboscis gland** for excretion, separate sexes, external fertilisation, and indirect development.

Practice Question 10 (Comparative Fish Biology MCQ)

Which of the following features correctly distinguishes *Scoliodon* (Dogfish) from *Hippocampus* (Sea horse)?

(1) *Scoliodon* has a terminal mouth and cycloid scales, whereas *Hippocampus* has a ventral mouth and placoid scales.

(2) *Scoliodon* possesses a persistent notochord, ventral mouth, placoid scales, and pelvic claspers in males, whereas *Hippocampus* has a bony endoskeleton, operculum, and air bladder.

(3) *Scoliodon* has a three-chambered heart and is homoiothermous, whereas *Hippocampus* has a two-chambered heart and is poikilothermous.

(4) *Scoliodon* exhibits external fertilisation and oviparity, whereas *Hippocampus* exhibits internal fertilisation and viviparity.

Correct Answer: (2)
Worked Solution: *Scoliodon* belongs to Class **Chondrichthyes** (cartilaginous fishes), which have a persistent notochord, ventral mouth, placoid scales, gill slits without operculum, no air bladder, pelvic claspers in males, and internal fertilisation. *Hippocampus* (Sea horse) belongs to Class **Osteichthyes** (bony fishes), which have a bony endoskeleton, terminal mouth, four pairs of gills covered by an operculum, cycloid/ctenoid scales, and an air bladder. Both classes have a two-chambered heart and are poikilothermous.

Practice Question 11 (Assertion–Reason Type)

Assertion (A): Adult *Petromyzon* (Lamprey) migrates from its marine habitat to freshwater rivers and streams during its life cycle.
Reason (R): Cyclostomes are marine animals that migrate to freshwater for spawning, after which the adults die within a few days and their larvae return to the ocean after metamorphosis.

(1) Both (A) and (R) are true and (R) is the correct explanation of (A).

(2) Both (A) and (R) are true, but (R) is not the correct explanation of (A).

(3) (A) is true, but (R) is false.

(4) (A) is false, but (R) is true.

Correct Answer: (1)
Worked Solution: Members of Class Cyclostomata (*Petromyzon* and *Myxine*) are jawless ectoparasites on fishes. They live in the sea (marine) but migrate to freshwater for spawning. Within a few days of spawning, the adult cyclostomes die, and their larvae undergo metamorphosis in freshwater before returning to the ocean. Thus both Assertion and Reason are true, and Reason is the exact explanation of Assertion.

Practice Question 12 (Tetrapoda Comparison MCQ)

Select the correct statement regarding the classes of Superclass Tetrapoda:

(1) All members of Class Reptilia possess a three-chambered heart and shed their scales as skin cast.

(2) Both Amphibia and Reptilia have a tympanum representing the ear, are poikilothermous, and exhibit indirect development.

(3) *Crocodilus* (Reptilia), *Pavo* (Aves), and *Ornithorhynchus* (Mammalia) all possess a four-chambered heart and are oviparous.

(4) In Class Aves, the skin is moist and glandular throughout the body, and forelimbs have scales for clasping tree branches.

Correct Answer: (3)
Worked Solution:
• *Crocodilus* (Crocodile) is an exception among reptiles in having a **four-chambered heart**, and like other reptiles, it is **oviparous**.
• *Pavo* (Peacock, Class Aves) has a completely **four-chambered heart** and is **oviparous**.
• *Ornithorhynchus* (Platypus, Class Mammalia) has a **four-chambered heart** and is an exceptional **oviparous mammal**.
Option (1) is incorrect because crocodiles have a 4-chambered heart and only snakes and lizards shed scales as skin cast. Option (2) is incorrect because Reptilia shows **direct development** (with internal fertilisation), whereas Amphibia shows indirect development (with external fertilisation). Option (4) is incorrect because bird skin is dry without glands except the oil gland at the base of the tail, and scales are on the **hind limbs**.

8. Common Exam Pitfalls & Revision Checklist

Pitfall 1: Confusing "Water Canal System" with "Water Vascular System"

Remember: The **water transport or canal system** (ostia → spongocoel → osculum, lined by choanocytes) is found in **Porifera**. The **water vascular system** is the distinctive feature of **Echinodermata**.

Pitfall 2: Misleading Common Names ("False Fishes")

In NEET UG, common names containing the word "fish" frequently belong to non-chordate phyla:

- **Jellyfish** (*Aurelia*) → Phylum Coelenterata (Cnidaria)
- **Cuttlefish** (*Sepia*), **Devil fish** (*Octopus*) → Phylum Mollusca
- **Starfish** (*Asterias*) → Phylum Echinodermata
- **Hagfish** (*Myxine*) → Class Cyclostomata (Jawless vertebrate, not a true jawed fish)
- **True Fishes (Superclass Pisces):** *Scoliodon* (Dogfish), *Pristis* (Sawfish), *Exocoetus* (Flying fish), *Hippocampus* (Sea horse), *Betta* (Fighting fish), *Pterophyllum* (Angel fish).

Pitfall 3: Exceptions in Symmetry, Monoecy/Dioecy, and Heart Chambers

- **Echinodermata Symmetry:** Adults are **radially symmetrical**, but larvae are **bilaterally symmetrical**.
- **Annelida Reproduction:** Earthworm (*Pheretima*) and Leech (*Hirudinaria*) are **monoecious**, whereas *Nereis* is **dioecious**.
- **Reptilian Heart:** Usually 3-chambered, but **4-chambered in crocodiles** (*Crocodilus*, *Alligator*).
- **Mammalian Reproduction:** Viviparous, except ***Ornithorhynchus*** **(Platypus)** which is **oviparous**.

Pitfall 4: Exclusively Marine Taxa

Questions often ask which groups are **exclusively marine**: **Ctenophora**, **Echinodermata**, **Hemichordata**, **Protochordates** (Urochordata and Cephalochordata), and Class **Chondrichthyes** are exclusively marine. (Note: Porifera and Cnidaria are *mostly* marine, as *Spongilla* and *Hydra* are freshwater forms).

9. Concise Final Recap

- **Basis of Classification:** Cellular level (Porifera); Tissue level (Coelenterata, Ctenophora); Organ level (Platyhelminthes); Organ-system level (Aschelminthes to Chordata). Diploblastic (Coelenterata, Ctenophora); Triploblastic (Platyhelminthes to Chordata). Acoelomate (up to Platyhelminthes); Pseudocoelomate (Aschelminthes); Coelomate (Annelida to Chordata).
- **Diagnostic Non-Chordate Structures:** Choanocytes & canal system (Porifera); Cnidoblasts & metagenesis (Cnidaria); Eight ciliated comb plates & bioluminescence (Ctenophora); Hooks, suckers & flame cells (Platyhelminthes); Muscular pharynx & sexual dimorphism (Aschelminthes); Metameric segmentation, nephridia & parapodia (Annelida); Chitinous exoskeleton, jointed appendages & Malpighian tubules (Arthropoda); Calcareous shell, mantle, feather-like gills & radula (Mollusca); Spiny skin & water vascular system (Echinodermata); Proboscis-collar-trunk, stomochord & proboscis gland (Hemichordata).
- **Phylum Chordata:** Characterised by a notochord, dorsal hollow single nerve cord, paired pharyngeal gill slits, ventral heart, and post-anal tail. Divided into Urochordata (notochord in larval tail), Cephalochordata (notochord head-to-tail throughout life), and Vertebrata (notochord replaced by vertebral column in adult).
- **Vertebrate Classes:** Agnatha includes jawless Cyclostomata. Gnathostomata includes Pisces (Chondrichthyes and Osteichthyes, bearing fins, 2-chambered heart, poikilothermous) and Tetrapoda (Amphibia, Reptilia, Aves, and Mammalia, bearing two pairs of limbs). Fishes, amphibians, and reptiles are poikilotherms; birds and mammals are homoiotherms with completely 4-chambered hearts.