Lecture 11: Sept 16

Last time

- Continuous Random Variables

Today

- Presentations
- Review part 2

Review part 2

Definition Given a sample space S and an associated sigma algebra $\mathcal{B}$, a *probability function* is a function Pr with domain $\mathcal{B}$ that satisfies

1. $\Pr(A) \geqslant 0$ for all $A \in \mathcal{B}$.
2. $\Pr(S) = 1$.
3. If $A_1, A_2, \dots \in \mathcal{B}$ are pairwise disjoint, then $\Pr(\cup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} \Pr(A_i)$.

The above three properties are usually referred to as the Axioms of Probability (or the Kolmogorov Axioms, after A. Kolmogorov, one of the fathers of probability theory). Any function that satisfies the Axioms of Probability is called a probability function.

Theorem If Pr is a probability function and A is any set in $\mathcal{B}$, then

1. $\Pr(\varnothing) = 0$, where $\varnothing$ is the empty set;
2. $\Pr(A) \leqslant 1$;
3. $\Pr(A^c) = 1 - \Pr(A)$.

Theorem If Pr is a probability function and A and B are any sets in $\mathcal{B}$, then

1. $\Pr(B \cap A^c) = \Pr(B) - \Pr(A \cap B)$;
2. $\Pr(A \cup B) = \Pr(A) + \Pr(B) - \Pr(A \cap B)$;
3. If $A \subset B$, then $\Pr(A) \leqslant \Pr(B)$.

Theorem If Pr is a probability function, then

1. $\Pr(A) = \sum_{i=1}^{\infty} \Pr(A \cap C_i)$ for any partition $C_1, C_2, \dots$;
2. $\Pr(\cup_{i=1}^{\infty} A_i) \leqslant \sum_{i=1}^{\infty} \Pr(A_i)$ for any sets $A_1, A_2, \dots$.

where (1) is also referred to as "Total probability" and (2) is Boole's inequality.

Definition If A and B are events in S, and $\Pr(B) > 0$, then the *conditional probability* of A given B, written $\Pr(A|B)$, is

$$\Pr(A|B) = \frac{\Pr(A \cap B)}{\Pr(B)}.$$

Note that B becomes the sample space now: $\Pr(B|B) = 1$.

Theorem (Bayes' Rule) Let $A_1, A_2, \ldots$ be a partition of the sample space, and let B be any set. Then, for each $i = 1, 2, \ldots$,

$$\Pr(A_i|B) = \frac{\Pr(B|A_i)\Pr(A_i)}{\sum_{j=1}^{\infty}\Pr(B|A_j)\Pr(A_j)}.$$