

Lecture 16: Sept 25

Last time

- Transformations of Random Variables

Today

- One-page one-sided letter-size cheat sheet for midterm 1
- Expected Values
- Moments

Expected Values

Definition The *expected value* or *mean* of a random variable $g(X)$, denoted by $Eg(X)$, is

$$Eg(X) = \begin{cases} \int_{-\infty}^{\infty} g(x)f(x)dx & \text{if } X \text{ is continuous} \\ \sum_{x \in \mathcal{X}} g(x) \Pr(X = x) & \text{if } X \text{ is discrete} \end{cases}$$

Provided the integral or summation exists.

If we let $g(X) = X$, then we get

$$EX = \begin{cases} \int_{-\infty}^{\infty} xf(x)dx & \text{if } X \text{ is continuous} \\ \sum_{x \in \mathcal{X}} x \Pr(X = x) & \text{if } X \text{ is discrete} \end{cases}$$

Example (Exponential mean) Suppose X has an *exponential distribution* with parameter λ , $X \sim \text{Exp}(\lambda)$, that is, it has pdf given by

$$f_X(x) = \frac{1}{\lambda}e^{-x/\lambda}, \quad 0 \leq x < \infty, \lambda > 0.$$

Find out EX .

Solution:

Example (Binomial mean) if X has a *binomial distribution*, $X \sim \text{Binomial}(n, p)$, its pmf is given by

$$\Pr(X = x) = \binom{n}{x} p^x (1-p)^{n-x}, \quad x = 0, 1, \dots, n,$$

where n is a positive integer, $0 \leq p \leq 1$, and for every fixed pair n and p the pmf sums to 1. Find out EX .

Solution:

The process of taking expectations is a linear operation, which means that the expectation of a linear function of X can be easily evaluated by noting that for any constants a and b , such that

$$E(aX + b) = aEX + b$$

Theorem Let X be a random variable and let a , b , and c be constants. Then for any functions $g_1(x)$ and $g_2(x)$ whose expectations exist,

1. $E(ag_1(X) + bg_2(X) + c) = aEg_1(X) + bEg_2(X) + c$.
2. If $g_1(x) \geq 0$ for all x , then $Eg_1(X) \geq 0$.
3. If $g_1(x) \geq g_2(x)$ for all x , then $Eg_1(X) \geq Eg_2(X)$.
4. If $a \leq g_1(x) \leq b$ for all x , then $a \leq Eg_1(X) \leq b$.

Proof: