

HW1 Due: Feb 12th

1 GitHub setup

On [GitHub](#)

- Obtain [student developer pack](#).
- Create a private repository `math-6040-2021-spring` (please substitute 6040 by 7260 if you are taking the graduate level). Add `xji3` and `louisnass` as your collaborators with write permission ([instruction](#)).

On your local machine:

- clone the repository: please refer to [this webpage](#) with instructions for your operating system.
- enter the folder: `cd math-6040-2021-spring`.
- after finishing the rest of the questions, save your file inside your git repository folder `math-6040-2021-spring` with name `hw1.pdf`. Please make it human-readable.
- now using git commands to stage this change: `git add hw1.pdf`
- commit: `git commit -m "hw1 submission"` (remember to replace the quotation mark)
- push to remote server: `git push`
- tag version hw1: `git tag hw1` and push: `git push --tags`.

Take a look at the tags on GitHub ([instructions](#)).

2 Show that for matrix $\mathbf{A} \in \mathbb{R}^{m \times n}$:

$$\text{rank}(\mathbf{A}) = \text{rank}(\mathbf{A}^T) = \text{rank}(\mathbf{A}^T \mathbf{A}) = \text{rank}(\mathbf{A} \mathbf{A}^T)$$

3 Show that for simple linear regression

$$y_i = \beta_0 + \beta_1 x_i + \epsilon_i,$$

the line ($\hat{y}_i = \hat{\beta}_0 + \hat{\beta}_1 x_i$) through the means of the variables (the point (\bar{x}, \bar{y})) has $\sum \epsilon_i = 0$

4 JF exercise 5.1

Prove that the least-squares fit in simple regression analysis has the following properties:

- (a) $\sum \hat{y}_i \epsilon_i = 0$.
- (b) $\sum (y_i - \hat{y}_i)(\hat{y}_i - \bar{y}) = \sum \epsilon_i(\hat{y}_i - \bar{y}) = 0$.

5 JF exercise 5.6

Why is it the case that the multiple-correlation coefficient R^2 can never get smaller when an explanatory variable is added to the regression equation? [*Hint:* Recall that the regression equation is fit by minimizing the residual sum of squares, which is equivalent as maximizing R^2 (why?).]