

Math 6040/7260 Linear Models

Mon/Wed/Fri 9:00am - 9:50am

Instructor: Dr. Xiang Ji, xji4@tulane.edu

1 Lecture 1: Jan 18

Today

- Introduction
- Introduce yourself
- Course logistics

What is this course about?

The term “linear models” describes a wide class of methods for the statistical analysis of multivariate data. The underlying theory is grounded in linear algebra and multivariate statistics, but applications range from biological research to public policy. The objective of this course is to provide a solid introduction to both the theory and practice of linear models, combining mathematical concepts with realistic examples.

Prerequisite

- **Must:** Introduction to Probability (Math 3070/6070), Mathematical Statistics (Math 3080/6080)
- **Good to have:** Scientific Computation II

A hierarchy of linear models

- The linear mean model:

$$\underset{n \times 1}{\mathbf{y}} = \underset{n \times p}{\mathbf{X}} \underset{p \times 1}{\boldsymbol{\beta}} + \underset{n \times 1}{\boldsymbol{\epsilon}}$$

where $\mathbf{E}(\boldsymbol{\epsilon}) = \mathbf{0}$. Only assumption is that errors have mean 0.

- Gauss-Markov model:

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}$$

where $\mathbf{E}(\boldsymbol{\epsilon}) = \mathbf{0}$ and $\mathbf{Var}(\boldsymbol{\epsilon}) = \sigma^2 \mathbf{I}$. Uncorrelated errors with constant variance.

- Aitken model or general linear model:

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}$$

where $\mathbf{E}(\boldsymbol{\epsilon}) = \mathbf{0}$ and $\mathbf{Var}(\boldsymbol{\epsilon}) = \sigma^2 \mathbf{V}$. \mathbf{V} is fixed and known.

- Variance components models: $\mathbf{y} \sim N(\mathbf{X}\beta, \sigma_1^2\mathbf{V}_1 + \sigma_2^2\mathbf{V}_2 + \dots + \sigma_r^2\mathbf{V}_r)$ with $\mathbf{V}_1, \mathbf{V}_2, \dots, \mathbf{V}_r$ known.
- General mixed linear Model:

$$\mathbf{y} = \mathbf{X}\beta + \epsilon$$

where $\mathbf{E}(\epsilon) = \mathbf{0}$ and $\mathbf{Var}(\epsilon) = \Sigma(\theta)$.

- Generalized linear models (GLMs). Logistic regression, probit regression, log-linear model (Poisson regression), ... Note the difference from the general linear model. GLMs are generalization of the *concept* of linear models. They are covered in Math 7360 - Data Analysis class (<https://tulane-math-7360-2021.github.io/>).

Syllabus

Check course website frequently for updates and announcements.

<https://tulane-math-7260-2023.github.io/>

HW submission

Through Github with demo on Friday class.

Presentations

Let me know your pick by the end of Friday (01/20/2023).

Last year (spring 2022) comments

1. Experience in this course

- I really enjoyed this class.
- The lecture is dry. Expect Dr. Ji to read through the lecture notes until someone asks a question. Also, having a background in statistics gives context to the lecture notes.

Response: I did emphasize on pre-requisite last year, but it still seemed not enough.

- I think this course was well laid out. Even though I did take it as a graduation requirement, I ended up enjoying the course. I also think the work load is manageable and professor Xi does provide all the tools necessary to succeed in the course. It can be overwhelming at first but with a little time and effort you can get the hang of the material.

Response: Mind the typos, students.

2. Strong aspects of this course

- Professor Ji is a great professor.

- Dr. Ji provides an inhuman amount of course material to help supplement learning, it was extremely helpful to have labs (answered and unanswered) as well as homework keys posted.

Response: There will be labs again.

- Notes are very structured and the professor is nice.
- The strongest aspect of this course is the homework. They are a great way to interact with and learn the material. The problems can seem challenging, but are doable with a little effort. Also, he publishes the keys afterword so you can check your work and see what the reasoning behind the answer is. This coupled with the lab sessions are a great way to prepare for the exams.

3. There will be an internal mid-term-ish evaluation for this course. Will remember to go over them.

Last last year (spring 2021) comments

1. Experience in this course

- Overall, I had a pretty good experience in this course. It moved quickly, but that is expected from this level of course. Sometimes it was hard to stay engaged with the lectures and to really absorb the course material. Because the lectures moved so fast, I really appreciated how Professor made the full notes available at the time of the lecture. I would have liked if there were a few more examples with the notes, as sometimes the homework felt disjoint from the notes.

Response: I will try to move slower this semester. I will start lab sessions earlier too.

- The professor is an extremely intelligent, kind, and understanding professor. He prioritizes in making sure that we understand the material and seeing how the material can be applied. His lecture notes were a godsend because the texts could be a bit ambiguous at times but he elucidated the material in such a comprehensible manner.

Response: I will try to fix the left-over typos.

- Mentioned in class from other students/internal evaluation, conveying the mathematical concepts through the presentation is not a good idea to follow the class in real-time. Prepared presentation can give rise to a distraction on what we have been going over.

Response: I am still delivering this class in hybrid-mode. I found the presentations fit online teaching better. I think the difficulty might be caused by (1) fast moving lecture (2) I only realized the need of reviewing basic concepts of probability almost a quarter into the semester...

- I found the setup of the course not very engaging. Additionally, many of the class notes came directly from the additional sources with no additional information or

explanation, which I found to be not very helpful.

Response: I actually like them. I was the guinea pig to test them.

- Easily help us to understand the main course, and the notes and details are great.

Response: There will be notes.

- Moves very quickly and can be hard to keep up with. Sometimes instructions are unclear.

Response: I will try to slow down.

- Both the instructor and the TA were helpful. It was hard to follow along in class though.

Response: We don't have TA this time. Make use of the office hour. And I have to say, it needs effort to ace in this class.

2. Strong aspects of this course

- Having the lecture notes and labs available was very helpful. Professor was also always very nice and accommodating, and willing to meet with me when I needed help. He also always responded to student feedback, if we asked for an extra day or two on the homework or something like that.

Response: Here is an example of correctly using the office hours.

- His lecture notes and the lab sessions.

Response: They will be there again.

- Lab session is necessarily required to this class. A lot of computations in the class would be done by computer due to the complexity, and students are expected to handle with the computer programming properly at a desired level. The course can be an introduction to the statistical computation, which does not exist in the mathematics department.

Response: Hmm, there is a course Math 7360 Data Analysis that focuses more on the computational side.

- I appreciated the homework reviews in class and felt these helped clarify the material.

Response: Of course, the reviews will be there again. The purpose of the course is for you to learn.

- Grading was easy which made up for the rigor.

Response: Don't rely on this...

- Really appreciate that Professor Xiang made such a neat and tidy notes for us. It is really helpful for me to review. And notes have a great interaction with us, Professor Xiang also leaves some questions to help us think about the logic behind.

Response: Well, Xiang is my first name. Please call me Prof. X.

- Prof. Xiang was highly organized and wanted his students to understand the course content more than he made them worry about grades. I learned a lot

about Linear Models and feel confident applying the course content professionally and academically. I wish most of the Math department had his teaching style and implemented his course documents and organization structure. Prof. Xiang made the course content in class digestible and if I needed to review the material I could easily find it through his course notes and textbook. I wish I could say the same about my other courses.

Response: Hmm, I like Prof. X. better.

- I really appreciated the emphasis on learning. It allowed for most students to take it at the pace that was good for them.

Response: Please don't let your score rely on this comment.

3. There will be an internal mid-term-ish evaluation for this course. Will remember to go over them.

2 Lecture 2:Jan 20

Last time

- Introduction
- Course logistics

Today

- Reply to the “Presentation Dates” thread on Canvas by the end of today.
- Git

What is git?

Git is currently the most popular system for version control according to [Google Trend](#). Git was initially designed and developed by [Linus Torvalds](#) in 2005 for Linux kernel development. Git is the British English slang for unpleasant person.

Why using git?

- [GitHub](#) is becoming a de facto central repository for open source development.
- **Advertise** yourself through GitHub (e.g., host a free personal webpage on GitHub).
- a skill that employers look for (according to [this AmStat article](#)).

Git workflow

Figure 2.1 shows its basic workflow.

What do I need to use Git?

- A **Git server** enabling multi-person collaboration through a centralized repository.
- A **Git client** on your own machine.
 - Linux: Git client program is shipped with many Linux distributions, e.g., Ubuntu and CentOS. If not, install using a package manager, e.g., `yum install git` on CentOS.
 - Mac: follow instructions at <https://www.atlassian.com/git/tutorials/install-git>.
 - Windows: Git for Windows at <https://gitforwindows.org> (GUI) aka Git Bash.
- Do **not** totally rely on GUI or IDE. Learn to use Git on command line, which is needed for cluster and cloud computing.




Figure 2.1

Git survival commands

- `git pull` synchronize local Git directory with remote repository.
- Modify files in local working directory.
- `git add FILES` add snapshots to staging area
- `git commit -m "message"` store snapshots permanently to (**local**) Git repository
- `git push` push commits to remote repository.

Git basic usage

Working with your local copy.

- `git pull` : update local Git repository with remote repository (fetch + merge).
- `git log FILENAME` : display the current status of working directory.
- `git diff` : show differences (by default difference from the most recent commit).
- `git add file1 file2 ...` : add file(s) to the staging area.

- `git commit` : commit changes in staging area to Git directory.
- `git push` : publish commits in local Git repository to remote repository.
- `git reset --soft HEAD 1` : undo the last commit.
- `git checkout FILENAME` : go back to the last commit, discarding all changes made.
- `git rm FILENAME` : remove files from git control.

Git demonstration

Show how to create a private git repository for HW and Exam submissions.

On [GitHub](#)

- Obtain [student developer pack](#).
- Create a private repository `math-6040-2023-spring` (please substitute 6040 by 7260 if you are taking the graduate level). Add `xji3` as your collaborators with write permission ([instruction](#)).

On your local machine:

- clone the repository: please refer to [this webpage](#) with instructions for your operating system.
- enter the folder: `cd math-6040-2023-spring`.
- after finishing the rest of the questions, save your file inside your git repository folder `math-6040-2023-spring` with name `hw1.pdf` (for example). Please make it human-readable.
- now using git commands to stage this change: `git add hw1.pdf`
- commit: `git commit -m "hw1 submission"` (remember to replace the quotation mark)
- push to remote server: `git push`
- tag version hw1: `git tag hw1` and push: `git push --tags`.

Take a look at the tags on GitHub ([instructions](#)).

When submitting your hw, please email your instructor (xji4@tulane.edu) a link to your tag ([instructions](#)).

3 Lecture 3:Jan 23

Last time

- Git

Today

- HW1 posted
- Linear algebra: vector and vector space, rank of a matrix
- Column space and Nullspace (JM Appendix A)

Notations

$$\underset{n \times 1}{\mathbf{y}} = \underset{n \times p}{\mathbf{X}} \underset{p \times 1}{\boldsymbol{\beta}} + \underset{n \times 1}{\boldsymbol{\epsilon}}$$

$$\begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix} = \begin{pmatrix} x_{11} & x_{12} & \dots & x_{1p} \\ x_{21} & x_{22} & \dots & x_{2p} \\ \vdots & \vdots & & \vdots \\ x_{n1} & x_{n2} & \dots & x_{np} \end{pmatrix} \begin{pmatrix} \beta_1 \\ \beta_2 \\ \vdots \\ \beta_p \end{pmatrix} + \begin{pmatrix} \epsilon_1 \\ \epsilon_2 \\ \vdots \\ \epsilon_n \end{pmatrix}$$

- All vectors are column vector
- Write dimensions underneath as in $\underset{n \times p}{\mathbf{X}}$ or as $\mathbf{X} \in \mathbb{R}^{n \times p}$
- Bold upper-case letters for Matrices. Bold lower-case letters for Vectors.

Vector and vector space

(from JM Appendix A)

- A set of vectors $\mathbf{x}_1, \dots, \mathbf{x}_n$ are *linearly dependent* if there exist coefficients c_j for $j = 1, 2, \dots, n$ such that $\sum_{j=1}^n c_j \mathbf{x}_j = \mathbf{0}$ and $\|\mathbf{c}\|_2 = \sum_{j=1}^n c_j^2 > 0$. They are *linearly independent* if $\sum_{j=1}^n c_j \mathbf{x}_j = \mathbf{0}$ implies (i.e. \implies) $c_j = 0$ for all j .
- Two vectors are *orthogonal* to each other, written $\mathbf{x} \perp \mathbf{y}$, if their inner product is 0, that is $\mathbf{x}^T \mathbf{y} = \mathbf{y}^T \mathbf{x} = \sum_j x_j y_j = 0$.
- A set of vectors $\mathbf{x}^{(1)}, \mathbf{x}^{(2)}, \dots, \mathbf{x}^{(n)}$ are mutually orthogonal iff (i.e. \iff) $\mathbf{x}^{(i)T} \mathbf{x}^{(j)} = 0$ for $\forall i \neq j$.
- The most common set of vectors that are mutually orthogonal are the *elementary* vectors $\mathbf{e}^{(1)}, \mathbf{e}^{(2)}, \dots, \mathbf{e}^{(n)}$, which are all zero, except for one element equal to 1, so that $\mathbf{e}_i^{(i)} = 1$ and $\mathbf{e}_j^{(i)} = 0, \forall j \neq i$.

- A *vector space* \mathcal{S} is a set of vectors that are closed under addition and scalar multiplication, that is
 - if $\mathbf{x}^{(1)}$ and $\mathbf{x}^{(2)}$ are in \mathcal{S} , then $c_1\mathbf{x}^{(1)} + c_2\mathbf{x}^{(2)}$ is in \mathcal{S} .
- A vector space \mathcal{S} is *generated* or *spanned* by a set of vectors $\mathbf{x}^{(1)}, \mathbf{x}^{(2)}, \dots, \mathbf{x}^{(n)}$, written as $\mathcal{S} = \text{span}\{\mathbf{x}^{(1)}, \mathbf{x}^{(2)}, \dots, \mathbf{x}^{(n)}\}$, if any vector \mathbf{x} in the vector space is a linear combination of $\mathbf{x}_i, i = 1, 2, \dots, n$.
- A set of linearly independent vectors that generate or span a space \mathcal{S} is called a *basis* of \mathcal{S} .

Example A.1

Let

$$\mathbf{x}^{(1)} = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \mathbf{x}^{(2)} = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \end{bmatrix}, \text{ and } \mathbf{x}^{(3)} = \begin{bmatrix} -3 \\ -1 \\ 1 \\ 3 \end{bmatrix}.$$

Then $\mathbf{x}^{(1)}$ and $\mathbf{x}^{(2)}$ are linearly independent, but $\mathbf{x}^{(1)}, \mathbf{x}^{(2)}$, and $\mathbf{x}^{(3)}$ are linearly dependent since $5\mathbf{x}^{(1)} - 2\mathbf{x}^{(2)} + \mathbf{x}^{(3)} = \mathbf{0}$

Rank

Some matrix concepts arise from viewing columns or rows of the matrix as vectors. Assume $\mathbf{A} \in \mathbb{R}^{m \times n}$.

- $\text{rank}(\mathbf{A})$ is the maximum number of linearly independent rows or columns of a matrix.
- $\text{rank}(\mathbf{A}) \leq \min\{m, n\}$.
- A matrix is *full rank* if $\text{rank}(\mathbf{A}) = \min\{m, n\}$. It is *full row rank* if $\text{rank}(\mathbf{A}) = m$. It is *full column rank* if $\text{rank}(\mathbf{A}) = n$.
- a square matrix $\mathbf{A} \in \mathbb{R}^{n \times n}$ is *singular* if $\text{rank}(\mathbf{A}) < n$ and *non-singular* if $\text{rank}(\mathbf{A}) = n$.
- $\text{rank}(\mathbf{A}) = \text{rank}(\mathbf{A}^T) = \text{rank}(\mathbf{A}^T \mathbf{A}) = \text{rank}(\mathbf{A} \mathbf{A}^T)$. (Show this in HW.)
- $\text{rank}(\mathbf{A} \mathbf{B}) \leq \min\{\text{rank}(\mathbf{A}), \text{rank}(\mathbf{B})\}$. (Hint: Columns of $\mathbf{A} \mathbf{B}$ are spanned by columns of \mathbf{A} and rows of $\mathbf{A} \mathbf{B}$ are spanned by rows of \mathbf{B} .)
- if $\mathbf{A} \mathbf{x} = \mathbf{0}_m$ for some $\mathbf{x} \neq \mathbf{0}_n$, then $\text{rank}(\mathbf{A}) \leq n - 1$.

Column space

Definition: The column space of a matrix, denoted by $\mathcal{C}(\mathbf{A})$ is the vector space spanned by the columns of the matrix, that is,

$$\mathcal{C}(\mathbf{A}) = \{\mathbf{x} : \text{there exists a vector } \mathbf{c} \text{ such that } \mathbf{x} = \mathbf{A} \mathbf{c}\}.$$

This means that if $\mathbf{x} \in \mathcal{C}(\mathbf{A})$, we can find coefficients c_j such that

$$\mathbf{x} = \sum_j c_j \mathbf{a}^{(j)}$$

where $\mathbf{a}^{(j)} = \mathbf{A}_{\cdot j}$ denotes the j^{th} column of matrix \mathbf{A} .

- The column space of a matrix consists of all vectors formed by multiplying that matrix by any vector.
- The number of basis vectors for $\mathcal{C}(\mathbf{A})$ is then the number of linearly independent columns of the matrix \mathbf{A} , and so, $\dim(\mathcal{C}(\mathbf{A})) = \text{rank}(\mathbf{A})$.
- The dimension of a space is the number of vectors in its basis.

Example A.2

Let $\mathbf{A} = \begin{bmatrix} 1 & 1 & -3 \\ 1 & 2 & -1 \\ 1 & 3 & 1 \\ 1 & 4 & 3 \end{bmatrix}$ and $\mathbf{c} = \begin{bmatrix} 5 \\ 4 \\ 3 \end{bmatrix}$. Show that \mathbf{Ac} is a linear combination of columns in \mathbf{A} .

solution:

$$\mathbf{Ac} = \begin{bmatrix} 1 \times 5 + 1 \times 4 + (-3) \times 3 \\ 1 \times 5 + 2 \times 4 + (-1) \times 3 \\ 1 \times 5 + 3 \times 4 + 1 \times 3 \\ 1 \times 5 + 4 \times 4 + 3 \times 3 \end{bmatrix} = \begin{bmatrix} 0 \\ 10 \\ 20 \\ 30 \end{bmatrix}.$$

You could recognize that

$$\mathbf{Ac} = 5 \times \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + 4 \times \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \end{bmatrix} + 3 \times \begin{bmatrix} -3 \\ -1 \\ 1 \\ 3 \end{bmatrix} = 5\mathbf{a}^{(1)} + 4\mathbf{a}^{(2)} + 3\mathbf{a}^{(3)} = \begin{bmatrix} 0 \\ 10 \\ 20 \\ 30 \end{bmatrix}.$$

Result A.1

$$\text{rank}(\mathbf{AB}) \leq \min(\text{rank}(\mathbf{A}), \text{rank}(\mathbf{B})).$$

proof: Each column of \mathbf{AB} is a linear combination of columns of \mathbf{A} (i.e. $(\mathbf{AB})_{\cdot j} = \mathbf{A}\mathbf{b}^{(j)}$), so the number of linearly independent columns of \mathbf{AB} cannot be greater than that of \mathbf{A} . Similarly, $\text{rank}(\mathbf{AB}) = \text{rank}(\mathbf{B}^T \mathbf{A}^T)$, the same argument gives $\text{rank}(\mathbf{B}^T)$ as an upper bound.

Result A.2

- (a) If $\mathbf{A} = \mathbf{BC}$, then $\mathcal{C}(\mathbf{A}) \subseteq \mathcal{C}(\mathbf{B})$.
- (b) If $\mathcal{C}(\mathbf{A}) \subseteq \mathcal{C}(\mathbf{B})$, then there exists a matrix \mathbf{C} such that $\mathbf{A} = \mathbf{BC}$.

proof: For (a), any vector $\mathbf{x} \in \mathcal{C}(\mathbf{A})$ can be written as $\mathbf{x} = \mathbf{Ad} = \mathbf{B}(\mathbf{Cd})$.

For (b), $\mathbf{A}_{\cdot j} \in \mathcal{C}(\mathbf{B})$, so that there exists a vector $\mathbf{c}^{(j)}$ such that $\mathbf{A}_{\cdot j} = \mathbf{B}\mathbf{c}^{(j)}$. The matrix $\mathbf{C} = (\mathbf{c}^{(1)}, \mathbf{c}^{(2)}, \dots, \mathbf{c}^{(n)})$ satisfies that $\mathbf{A} = \mathbf{BC}$.

4 Lecture 4: Jan 25

Last time

- Linear algebra: vector and vector space, rank of a matrix
- Column space (JM Appendix A)

Today

- Nullspace
- Probability review

Null space

Definition: The null space of a matrix, denoted by $\mathcal{N}(\mathbf{A})$, is $\mathcal{N}(\mathbf{A}) = \{\mathbf{y} : \mathbf{A}\mathbf{y} = \mathbf{0}\}$.

Result A.3

If \mathbf{A} has full-column rank, then $\mathcal{N}(\mathbf{A}) = \{\mathbf{0}\}$.

proof: Matrix \mathbf{A} has full-column rank means its columns are linearly independent, which means that $\mathbf{A}\mathbf{c} = \mathbf{0}$ implies $\mathbf{c} = \mathbf{0}$.

Theorem A.1

Assume $\mathbf{A} \in \mathbb{R}^{m \times n}$, then $\dim(\mathcal{C}(\mathbf{A})) = r$ and $\dim(\mathcal{N}(\mathbf{A})) = n - r$, where $r = \text{rank}(\mathbf{A})$.

See JM Appendix Theorem A.1 for the proof.

proof: Denote $\dim(\mathcal{N}(\mathbf{A}))$ by k , to be determined, and construct a set of basis vectors for $\mathcal{N}(\mathbf{A}) : \{\mathbf{u}^{(1)}, \mathbf{u}^{(2)}, \dots, \mathbf{u}^{(k)}\}$, so that $\mathbf{A}\mathbf{u}^{(i)} = \mathbf{0}$, for $i = 1, 2, \dots, k$. Now, construct a basis for \mathbb{R}^n by adding the vectors $\{\mathbf{u}^{(k+1)}, \dots, \mathbf{u}^{(n)}\}$, which are not in $\mathcal{N}(\mathbf{A})$. Clearly, $\mathbf{A}\mathbf{u}^{(i)} \in \mathcal{C}(\mathbf{A})$ for $i = k+1, \dots, n$, and so the span of these vectors form a subspace of $\mathcal{C}(\mathbf{A})$. These vectors $\{\mathbf{A}\mathbf{u}^{(i)}, i = k+1, \dots, n\}$ are also linearly independent from the following argument: suppose $\sum_{i=k+1}^n c_i \mathbf{A}\mathbf{u}^{(i)} = \mathbf{0}$; then $\sum_{i=k+1}^n c_i \mathbf{A}\mathbf{u}^{(i)} = \mathbf{A} [\sum_{i=k+1}^n c_i \mathbf{u}^{(i)}] = \mathbf{0}$, and hence $\sum_{i=k+1}^n c_i \mathbf{u}^{(i)}$ is a vector in $\mathcal{N}(\mathbf{A})$. Therefore, there exist b_i such that $\sum_{i=k+1}^n c_i \mathbf{u}^{(i)} = \sum_{i=1}^k b_i \mathbf{u}^{(i)}$, or $\sum_{i=1}^k b_i \mathbf{u}^{(i)} - \sum_{i=k+1}^n c_i \mathbf{u}^{(i)} = \mathbf{0}$. Since $\{\mathbf{u}^{(i)}\}$ form a basis for \mathbb{R}^n , c_i must all be zero. Therefore $\mathbf{A}\mathbf{u}^{(i)}, i = k+1, \dots, n$ are linearly independent. At this point, since $\text{span}\{\mathbf{A}\mathbf{u}^{(k+1)}, \dots, \mathbf{A}\mathbf{u}^{(n)}\} \subseteq \mathcal{C}(\mathbf{A})$, $\dim(\mathcal{C}(\mathbf{A}))$ is at least $n - k$. Suppose there is a vector \mathbf{y} that is in $\mathcal{C}(\mathbf{A})$, but not in the span; then there exists $\mathbf{u}^{(n+1)}$ so that $\mathbf{y} = \mathbf{A}\mathbf{u}^{(n+1)}$ and $\mathbf{u}^{(n+1)}$ is linearly independent of $\{\mathbf{u}^{(k+1)}, \dots, \mathbf{u}^{(n)}\}$ (and clearly not in $\mathcal{N}(\mathbf{A})$), making $n+1$ linearly independent vectors in \mathbb{R}^n . Since that is not possible, the span is equal to $\mathcal{C}(\mathbf{A})$ and $\dim(\mathcal{C}(\mathbf{A})) = n - k = r = \text{rank}(\mathbf{A})$, so that $k = \dim(\mathcal{N}(\mathbf{A})) = n - r$.

Interpretation: “dimension of column space + dimension of null space = # columns”

Mis-Interpretation: Columns space and null space are orthogonal complement to each other.

They are of different orders in general! Next result gives the correct statement.

Reference:

- Statistical Inference, 2nd Edition, by George Casella & Roger L. Berger
- [Review of Probability Theory](#) by Arian Maleki and Tom Do

Probability theory review

A few basic elements to define a probability on a set:

- **Sample space** S is the set that contains all possible outcomes of a particular experiment.
- An **event** is any collection of possible outcomes of an experiment, that is, any subset of S (including S itself).
- Event operations
 1. Union: The union of A and B , written $A \cup B$, is the set of elements that belong to either A or B or both:

$$A \cup B = \{x : x \in A \text{ or } x \in B\}$$

2. Intersection: The intersection of A and B , written $A \cap B$, is the set of elements that belong to both A and B :

$$A \cap B = \{x : x \in A \text{ and } x \in B\}$$

3. Complementation: The complement of A , written as A^c , is the set of all elements that are not in A :

$$A^c = \{x : x \notin A\}.$$

- **Sigma algebra (or Borel field):** A collection of subsets of S is called a sigma algebra (or Borel field), denoted by \mathcal{B} , if it satisfies the following three properties:
 1. $\emptyset \in \mathcal{B}$ (the empty set is an element of \mathcal{B})
 2. If $A \in \mathcal{B}$, then $A^c \in \mathcal{B}$ (\mathcal{B} is closed under complementation).
 3. If $A_1, A_2, \dots \in \mathcal{B}$, then $\cup_{i=1}^{\infty} A_i \in \mathcal{B}$ (\mathcal{B} is closed under countable unions).
- **Axioms of probability:** Given a sample space S and an associated sigma algebra \mathcal{B} , a *probability function* is a function $\Pr(\cdot)$ with domain \mathcal{B} that satisfies
 1. $\Pr(A) \geq 0$ for all $A \in \mathcal{B}$
 2. $\Pr(S) = 1$.
 3. If $A_1, A_2, \dots \in \mathcal{B}$ are pairwise disjoint, then $\Pr(\cup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} \Pr(A_i)$.

Properties:

If $\Pr()$ is a *probability function* and A and B are any sets in \mathcal{B} , then

- $\Pr(\emptyset) = 0$, where \emptyset is the empty set
Proof: $1 = \Pr(S) = \Pr(S \cup \emptyset)$
- $\Pr(A) \leq 1$
Proof: see below and remember $\Pr(A^c) \geq 0$
- $\Pr(A^c) = 1 - \Pr(A)$
Proof: $1 = \Pr(S) = \Pr(A \cup A^c) = \Pr(A) + \Pr(A^c)$
- $\Pr(B \cap A^c) = \Pr(B) - \Pr(A \cap B)$
Proof: $B = \{B \cap A\} \cup \{B \cap A^c\}$
- $\Pr(A \cup B) = \Pr(A) + \Pr(B) - \Pr(A \cap B)$
Proof: $A \cup B = A \cup \{B \cap A^c\}$ and use the above property.
- $\Pr(A \cup B) = \Pr(A) + \Pr(B \cap A^c) = \Pr(A) + \Pr(B) - \Pr(A \cap B)$
- If $A \subset B$, then $\Pr(A) \leq \Pr(B)$.
Proof: If $A \subset B$, then $A \cap B = A$ and use $\Pr(B \cap A^c) = \Pr(B) - \Pr(A \cap B)$.

Conditional probability

Definition: If A and B are events in S , and $\Pr(B) > 0$, then the conditional probability of A given B , written $\Pr(A|B)$, is

$$\Pr(A|B) = \frac{\Pr(A \cap B)}{\Pr(B)}$$

Note that what happens in the conditional probability calculation is that B becomes the sample space: $\Pr(B|B) = 1$, in other words, $\Pr(A|B)$ is the probability measure of the event A after observing the occurrence of event B .

Definition: Two events A and B are statistically independent if $\Pr(A \cap B) = \Pr(A) \Pr(B)$. When A and B are independent events, then $\Pr(A|B) = \Pr(A)$ and the following pairs are also independent

- A and B^c

proof:

$$\begin{aligned}\Pr(A \cap B^c) &= \Pr(A) - \Pr(A \cap B) \\ &= \Pr(A) - \Pr(A) \Pr(B) \\ &= \Pr(A)(1 - \Pr(B)) \\ &= \Pr(A) \Pr(B^c)\end{aligned}$$

- A^c and B
- A^c and B^c

Random variables

Definition: A random variable is a function from a sample space S into the real numbers.

Experiment	Random variable
Toss two dice	$X = \text{sum of the numbers}$
Toss a coin 25 times	$X = \text{number of heads in 25 tosses}$
Apply different amounts of fertilizer to corn plants	$X = \text{yield/acre}$

Suppose we have a sample space

$$S = \{s_1, \dots, s_n\}$$

with a probability function \Pr and we define a random variable X with range $\mathcal{X} = \{x_1, \dots, x_m\}$. We can define a probability function \Pr_X on \mathcal{X} in the following way. We will observe $X = x_i$ if and only if the outcome of the random experiment is an $s_j \in S$ such that $X(s_j) = x_i$. Thus,

$$\Pr_X(X = x_i) = \Pr(\{s_j \in S : X(s_j) = x_i\}).$$

We will simply write $\Pr(X = x_i)$ rather than $\Pr_X(X = x_i)$.

A note on notation: Random variables are often denoted with uppercase letters and the realized values of the variables (or its range) are denoted by corresponding lowercase letters.

Distribution functions

Definition: The cumulative distribution function or cdf of a random variable (r.v.) X , denoted by $F_X(x)$ is defined by

$$F_X(x) = \Pr(X \leq x), \text{ for all } x.$$

The function $F(x)$ is a cdf if and only if the following three conditions hold:

1. $\lim_{x \rightarrow \infty} F(x) = 1$.
2. $F(x)$ is a nondecreasing function of x .
3. $F(x)$ is right-continuous; that is, for every number x_0 , $\lim_{x \downarrow x_0} F(x) = F(x_0)$.

Definition: A random variable X is continuous if $F(x)$ is a continuous function of x . A random variable X is discrete if $F(x)$ is a step function of x .

The following two statements are equivalent:

1. The random variables X and Y are identically distributed.
2. $F_X(x) = F_Y(x)$ for every x .

Density and mass functions

Definition: The probability mass function (pmf) of a discrete random variable X is given by

$$f_X(x) = \Pr(X = x) \text{ for all } x.$$

Example (Geometric probabilities) For the geometric distribution, we have the pmf

$$f_X(x) = \Pr(X = x) = \begin{cases} p(1-p)^{x-1} & \text{for } x = 1, 2, \dots \\ 0 & \text{otherwise.} \end{cases}$$

Definition: The probability density function or pdf, $f_X(x)$, of a continuous random variable X is the function that satisfies

$$F_X(x) = \int_{-\infty}^x f_X(t)dt \quad \text{for all } x.$$

A note on notation: The expression “ X has a distribution given by $F_X(x)$ ” is abbreviated symbolically by “ $X \sim F_X(x)$ ”, where we read the symbol “ \sim ” as “is distributed as”.

Example (Logistic distribution) For the logistic distribution, we have

$$F_X(x) = \frac{1}{1 + e^{-x}}$$

and, hence,

$$f_X(x) = \frac{d}{dx}F_X(x) = \frac{e^{-x}}{(1 + e^{-x})^2}.$$

A function $f_X(x)$ is a pdf (or pmf) of a random variable X if and only if

1. $f_X(x) \geq 0$ for all x
2. $\sum_x f_X(x) = 1$ (pmf) or $\int_{-\infty}^{\infty} f_X(x)dx = 1$ (pdf).

Expectations

The expected value, or expectation, of a random variable is merely its average value, where we speak of “average” value as one that is weighted according to the probability distribution.

Definition: The expected value or mean of a random variable $g(X)$, denoted by $\mathbf{E}(g(X))$, is

$$\mathbf{E}(g(X)) = \begin{cases} \int_{-\infty}^{\infty} g(x)f_X(x)dx & \text{if } X \text{ is continuous} \\ \sum_{x \in \mathcal{X}} g(x)f_X(x) = \sum_{x \in \mathcal{X}} g(x)\Pr(X = x) & \text{if } X \text{ is discrete,} \end{cases}$$

Exponential mean

Suppose $X \sim \text{Exp}(\lambda)$ distribution, that is, it has pdf given by

$$f_X(x) = \frac{1}{\lambda}e^{-x/\lambda}, \quad 0 \leq x < \infty, \quad \lambda > 0$$

Then $\mathbf{E}(X)$ is:

$$\begin{aligned}\mathbf{E}(X) &= \int_0^\infty \frac{1}{\lambda} x e^{-x/\lambda} dx \\ &= -x e^{-x/\lambda} \Big|_0^\infty + \int_0^\infty e^{-x/\lambda} dx \\ &= \int_0^\infty e^{-x/\lambda} dx = \lambda\end{aligned}$$

Binomial mean

IF X has binomial distribution, i.e. $X \sim \text{binomial}(n, p)$, its pmf is given by

$$\Pr(X = x) = \binom{n}{x} p^x (1-p)^{n-x}, \quad x = 0, 1, \dots, n,$$

where n is a positive integer, $0 \leq p \leq 1$, and for every fixed pair n and p the pmf sums to 1. The expected value of a binomial random variable is then given by

$$\mathbf{E}(X) = \sum_{x=0}^n x \binom{n}{x} p^x (1-p)^{n-x}$$

Now, use the identity $x \binom{n}{x} = n \binom{n-1}{x-1}$ to derive the Expected value.

$$\begin{aligned}\mathbf{E}(X) &= \sum_{x=1}^n x \binom{n}{x} p^x (1-p)^{n-x} \\ &= \sum_{x=1}^n n \binom{n-1}{x-1} p^x (1-p)^{n-x} \\ &= \sum_{y=0}^{n-1} n \binom{n-1}{y} p^{y+1} (1-p)^{n-(y+1)} \\ &= np \sum_{y=0}^{n-1} \binom{n-1}{y} p^y (1-p)^{n-1-y} \\ &= np,\end{aligned}$$

since the last summation must be 1, being the sum over all possible values of a $\text{binomial}(n-1, p)$ pmf.

properties:

Let X be a random variable and let a, b and c be constants. Then for any functions $g_1(x)$ and $g_2(x)$ whose expectations exist,

$$1. \quad \mathbf{E}(a \cdot g_1(X) + b \cdot g_2(X) + c) = a\mathbf{E}(g_1(X)) + b\mathbf{E}(g_2(X)) + c.$$

2. If $g_1(x) \geq 0$ for all x , then $\mathbf{E}(g_1(X)) \geq 0$.
3. If $g_1(x) \geq g_2(x)$ for all x , then $\mathbf{E}(g_1(X)) \geq \mathbf{E}(g_2(X))$.
4. If $a \leq g_1(x) \leq b$ for all x , then $a \leq \mathbf{E}(g_1(X)) \leq b$.

Moments

The various moments of a distribution are an important class of expectations.

Definition: For each integer n , the n^{th} moment of X (or $F_X(x)$), μ'_n , is

$$\mu'_n = \mathbf{E}(X^n).$$

The n^{th} central moment of X , μ_n , is

$$\mu_n = \mathbf{E}((X - \mu)^n),$$

where $\mu = \mu'_1 = \mathbf{E}(X)$.

Variance

Definition: The variance of a random variable X is its second central moment, $\mathbf{Var}(X) = \mathbf{E}((X - EX)^2)$. The positive square root of $\mathbf{Var}(X)$ is the standard deviation of X .

Exponential variance

Let X have the exponential(λ) distribution, $X \sim \text{Exp}(\lambda)$. Then the variance of X is

$$\begin{aligned} \mathbf{Var}(X) &= \mathbf{E}((X - EX)^2) = \mathbf{E}((X - \lambda)^2) \\ &= \int_0^\infty (x - \lambda)^2 \frac{1}{\lambda} e^{-x/\lambda} dx \\ &= \int_0^\infty (x^2 - 2x\lambda + \lambda^2) \frac{1}{\lambda} e^{-x/\lambda} dx \\ &= \lambda^2. \end{aligned}$$

properties

1. $\mathbf{Var}(aX + b) = a^2 \mathbf{Var}(X)$.

proof:

$$\begin{aligned} \mathbf{Var}(aX + b) &= \mathbf{E}(((aX + b) - \mathbf{E}(aX + b))^2) \\ &= \mathbf{E}((aX - aEX)^2) \\ &= a^2 \mathbf{E}((X - EX)^2) \\ &= a^2 \mathbf{Var}(X) \end{aligned}$$

$$2. \text{Var}(X) = \mathbf{E}(X^2) - (\mathbf{E}(X))^2.$$

proof:

$$\begin{aligned} \text{Var}(X) &= \mathbf{E}(X - EX)^2 \\ &= \mathbf{E}(X^2 - 2X\mathbf{E}(X) + (\mathbf{E}(X))^2) \\ &= \mathbf{E}(X^2) - 2\mathbf{E}(X)\mathbf{E}(X) + (\mathbf{E}(X))^2 \\ &= \mathbf{E}(X^2) - (\mathbf{E}(X))^2 \end{aligned}$$

Moment generating function

Definition: Let X be a random variable with cdf F_X . The moment generating function or mgf of X (or F_X), denoted by $M_X(t)$, is

$$M_X(t) = \mathbf{E}(e^{tX}),$$

provided that the expectation exists for t in some neighborhood of 0. That is, there exists an $h > 0$ such that for all t in $-h < t < h$, $\mathbf{E}(e^{tX})$ exists. If the expectation does not exist in a neighborhood of 0, we say that the moment generating function does not exist.

Property: If X has mgf $M_X(t)$, then

$$\mathbf{E}(X^n) = M_X^{(n)}(0),$$

where we define

$$M_X^{(n)}(0) = \left. \frac{d^n}{dt^n} M_X(t) \right|_{t=0}.$$