NCK Paper 2 Batch 8

Revision Questions with Correct Answers and Detailed Rationales

Use this document for focused revision. Each question is followed by the correct answer and a clear rationale explaining why that option is correct and why others are less appropriate.

Question 1

The nurse employed in the emergency room is responsible for triage for 4 clients injured in a Motor vehicle accident. Which of the following clients should receive priority care?

A. A 10-year-old with lacerations of the face
B. A 15-year-old with sternal bruises
C. A 34-year-old with a fractured femur
D. A 50-year-old with dislocation of the elbow

✓ Correct Answer: B

Rationale:
Sternal bruises after blunt chest trauma raise concern for myocardial contusion, cardiac tamponade, or major vessel injury. These are life-threatening and take priority over facial lacerations, long-bone fracture, or elbow dislocation in triage (ABC approach).

Question 2

Which of the following roommates would be most suitable for the client with myasthenia gravis?

A. A client with Hypothyroidism
B. A client with Crohn's disease
C. A client with pyelonephritis
D. A client with Bronchitis

✓ Correct Answer: A

Rationale:
Clients with myasthenia gravis are highly susceptible to respiratory infections that can precipitate myasthenic crisis. Hypothyroidism is non-infectious and does not pose an infection risk, making it the safest roommate choice.

Question 3

The invasive stage of Entamoeba histolytica which is responsible for tissue amoebiasis is:

A. Cyst
B. Trophozoite
C. Larvae
D. Egg

✓ Correct Answer: B

Rationale:
The trophozoite (motile feeding form) invades the intestinal mucosa and can cause tissue destruction (amoebic colitis, liver abscess). Cysts are the infective, non-invasive transmission stage.

Question 4

During ovulation:

A. Vaginal secretions become thick and sticky, and BBT decreases sharply
B. Vaginal secretions become clear and stretchy, and BBT experiences a slight increase
C. Vaginal secretions become scanty and dry, and BBT remains constant
D. Vaginal secretions have a creamy consistency, and BBT decreases gradually

✓ Correct Answer: B

Rationale:
At ovulation, cervical mucus becomes clear, stretchy (spinnbarkeit / egg-white consistency) under estrogen influence. Basal body temperature shows a slight rise (0.2–0.5°C) after ovulation due to progesterone.

Question 5

Mrs. P has been diagnosed with congestive heart failure. The nurse understands that the drug which has been used to reduce morbidity and mortality of this particular disease is:

A. Verapamil
B. Nifedipine
C. Digoxin
D. Captopril

✓ Correct Answer: D

Rationale:
ACE inhibitors (e.g., Captopril) have proven mortality and morbidity benefit in heart failure by reducing afterload, preventing remodeling, and improving survival. Digoxin improves symptoms and reduces hospitalizations but does not reduce mortality. Calcium-channel blockers are generally avoided in systolic HF.

Question 6

You have just entered the NBU and have been assigned 4 newborns to take care of. Which of the following newborns will you assess first?

A. A baby who is 36 weeks gestation reported as stable
B. A baby born with first minute APGAR score of 4, and 6 at 5 minutes
C. A baby who is reported to be 4.5 kg at birth
D. A baby who is reported to have been born with meconium grade III amniotic fluid

✓ Correct Answer: B

Rationale:
An APGAR of 4 at 1 minute and 6 at 5 minutes indicates significant depression at birth and ongoing compromise. This neonate requires immediate reassessment of airway, breathing, and circulation before the other relatively stable or at-risk but currently stable infants.

Question 7

Qualitative research primarily deals with which type of data?

A. Discrete data
B. Numerical data
C. Categorical data
D. Continuous data

✓ Correct Answer: C

Rationale:
Qualitative research explores meanings, experiences, and social phenomena using non-numerical data such as interviews, observations, and categorical descriptions. Discrete, continuous, and numerical data are quantitative.

Question 8

The process of learning and adopting other people's culture is through:

A. Socialization
B. Acculturation
C. Enculturation
D. Cultural immersion

✓ Correct Answer: B

Rationale:
Acculturation is the process of adopting cultural traits or social patterns of another group. Enculturation is learning one's own culture; socialization is broader social learning.

Question 9

The theory that nursing is a helping or assisting profession to persons who are wholly or partially dependent on those who are supposedly caring for them is attributed to:

A. Henderson
B. Orem
C. Roy
D. Nightingale

✓ Correct Answer: B

Rationale:
Dorothea Orem's Self-Care Deficit Theory describes nursing as assisting individuals who have a deficit in self-care (wholly or partially dependent). Henderson focused on 14 basic needs; Roy on adaptation; Nightingale on environment.

Question 10

A 17-year-old is bleeding between periods of less than 2 weeks. This condition is an abnormality in the menstrual cycle known as:

A. Metrorrhagia
B. Menorrhagia
C. Amenorrhea
D. Dysmenorrhea

✓ Correct Answer: A

Rationale:
Metrorrhagia is irregular uterine bleeding between normal menstrual periods. Menorrhagia is excessive/prolonged bleeding at regular intervals; amenorrhea is absence of menses; dysmenorrhea is painful menses.

Question 11

The following predispose to primary post-partum haemorrhage except:

A. Administration of prolonged or deep anesthesia to the mother
B. Multiple pregnancies
C. Prolonged labor
D. Puerperal sepsis

✓ Correct Answer: D

Rationale:
Primary PPH occurs within 24 hours of delivery. Risk factors include uterine atony from prolonged labor, multiple pregnancy, and anesthesia effects. Puerperal sepsis is a cause of secondary PPH (after 24 hours).

Question 12

A patient is brought to the emergency department after he was rescued while trying to commit suicide by hanging. Which is your next step in managing the patient?

A. Take history of the events leading to the attempt
B. Ask the client to talk about the experience
C. Examine the airway and the neck
D. Refer to a psychiatrist

✓ Correct Answer: C

Rationale:
Hanging can cause airway compromise, cervical spine injury, or vascular damage. Immediate assessment and stabilization of airway and neck (ABC + C-spine) take absolute priority over history or psychiatric referral.

Question 13

Which of the following is true about suicide?

A. Females try to commit suicide more often than men
B. Females have a higher success rate
C. A family history of suicide increases one's risk
D. A person with a suicide plan is at higher risk

✓ Correct Answer: A

Rationale:
Females attempt suicide more frequently than males, but males have a higher completion rate (more lethal methods). Family history and having a concrete plan both increase risk, but the most classically correct statement among the options is A regarding attempt rates.

Question 14

Second degree uterine prolapse is characterized by:

A. Complete protrusion of the uterus outside the introitus
B. Descent of the uterus within the vagina
C. Descent of the cervix below the ischial spines
D. Descent of the uterus outside the vulva during straining

✓ Correct Answer: B

Rationale:
In second-degree prolapse the cervix descends into the vagina (often to or beyond the introitus on straining) but the uterus is not completely outside. Complete protrusion outside the introitus is third-degree (procidentia).

Question 15

In management, Theory Y assumes that:

A. People are committed to organization's activities
B. People dislike work
C. People are poorly motivated
D. People achieve little satisfaction from work

✓ Correct Answer: A

Rationale:
McGregor's Theory Y posits that people are self-motivated, enjoy work, seek responsibility, and are committed to organizational goals when properly engaged. Theory X assumes the opposite (dislike of work, need for control).

Question 16

Which anticoagulant is considered the first line treatment for DVT during pregnancy?

A. Warfarin
B. Rivaroxaban
C. Low molecular weight heparin
D. Aspirin

✓ Correct Answer: C

Rationale:
LMWH (e.g., enoxaparin) does not cross the placenta and is the preferred anticoagulant for VTE in pregnancy. Warfarin is teratogenic; DOACs (rivaroxaban) lack sufficient safety data; aspirin alone is inadequate for treatment of DVT.

Question 17

Largest gland in the body:

A. Pituitary gland
B. Adrenal gland
C. Thyroid gland
D. Liver

✓ Correct Answer: D

Rationale:
The liver is the largest gland (and largest internal organ) in the human body. The pituitary, adrenal, and thyroid are much smaller endocrine glands. (Note: some keys list pituitary, but anatomically the liver is correct.)

Question 18

A patient's health assessment has resulted in a diagnosis of alopecia areata. What nursing diagnosis should the nurse likely associate with this health problem?

A. Chronic pain related to the disease
B. Impaired skin integrity related to the disease
C. Impaired tissue integrity observed on examination
D. Disturbed body image as reported by the patient

✓ Correct Answer: D

Rationale:
Alopecia areata causes patchy hair loss that is often cosmetically distressing. The primary nursing concern is usually disturbed body image / self-esteem rather than pain or tissue integrity (the scalp itself is intact).

Question 19

A patient in the medical outpatient clinic complained to the nurse of chest pain. What aspect of the patient's pain would be most suggestive of angina as the cause?

A. The pain is worse when the patient inhales deeply
B. The pain occurs immediately following physical exertion
C. The pain is worse when the patient coughs
D. The pain is most severe when the patient moves his upper body

✓ Correct Answer: B

Rationale:
Classic angina is precipitated by exertion or emotional stress and relieved by rest or nitroglycerin. Pain worsened by inspiration, cough, or movement is more consistent with musculoskeletal, pleural, or pericardial causes.

Question 20

Pancreatic juice:

A. Deactivates bile
B. Is secreted by pancreatic islet cells
C. Buffers chyme
D. Is released into the cystic duct

✓ Correct Answer: C

Rationale:
Pancreatic juice is rich in bicarbonate that neutralizes acidic chyme entering the duodenum. It is secreted by acinar and ductal cells (not islets), does not deactivate bile, and enters via the pancreatic duct into the duodenum (not cystic duct).

Question 21

What should the nurse suspect when hourly assessment of urine output on a patient post craniotomy exhibits urine output from a catheter of 1500 ml for 2 consecutive hours?

A. Cushing syndrome
B. Syndrome of inappropriate Anti-diuretic hormone
C. Adrenal crisis
D. Diabetes insipidus

✓ Correct Answer: D

Rationale:
Massive dilute urine output after neurosurgery is classic for central diabetes insipidus (deficient ADH). SIADH causes oliguria and hyponatremia; Cushing and adrenal crisis have different fluid patterns.

Question 22

In which situation is ego development evident?

A. A client calls out in pain to get his/her needs met
B. A client complains of poor self-esteem because of punishments from his/her past
C. A client exhibits the ability to assert himself/herself without anger/aggression
D. A client morally feels guilty about wanting to have sexual relations outside of marriage

✓ Correct Answer: C

Rationale:
A mature ego mediates between id impulses and superego demands, allowing assertive expression of needs without aggression or excessive guilt. Option C demonstrates healthy ego function.

Question 23

The nurse completes an orthopedic assessment of a 10-year-old child who has a new cast applied for fractured radius. Which of the following clinical manifestations is a priority for the nurse to report immediately to the physician?

A. A capillary refill of 3 seconds in the affected hand
B. The child states that the hand feels asleep
C. Skin around the cast is warm
D. None of the above

✓ Correct Answer: B

Rationale:
Paresthesia ('hand feels asleep') is an early sign of neurovascular compromise (compartment syndrome or cast tightness). Capillary refill of 3 seconds is borderline acceptable; warm skin is expected. Immediate physician notification is required for sensory changes.

Question 24

A pregnant school-age child undergoing chemotherapy experiences nausea and vomiting. Which of the following would be the best intervention to include in the child's plan of care?

A. Allow the child to choose what to eat for meals
B. Administer tube feedings
C. Offer small frequent meals
D. Offer fluids between meals

✓ Correct Answer: C

Rationale:
Small, frequent meals are better tolerated than large meals during chemotherapy-induced nausea. Allowing choice is helpful but secondary; tube feeding is invasive and not first-line; fluids between meals help prevent early satiety.

Question 25

In a client with substance-induced persisting dementia, the patient doesn't remember facts and fills gaps with imaginary information. This is called:

A. Flight of ideas
B. Confabulation
C. Concretism
D. Perseveration

✓ Correct Answer: B

Rationale:
Confabulation is the unconscious filling of memory gaps with fabricated but often plausible details, classic in Korsakoff's (alcohol-related) amnesia and other organic dementias.

Question 26

What is the most appropriate nursing response to a myocardial infarction patient who is fearful of dying?

A. Tell me about your feelings right now
B. When the doctor arrives everything will be fine
C. This is a bad situation but you will feel better soon
D. Please be assured we are doing everything we can to make you feel better

✓ Correct Answer: A

Rationale:
Open-ended exploration of feelings ('Tell me about your feelings…') validates the patient's fear and encourages therapeutic communication. False reassurance (B, C, D) blocks expression of anxiety.

Question 27

A medication given in alcohol withdrawal is?

A. Naloxone
B. Lorazepam
C. Haloperidol
D. Methadone

✓ Correct Answer: B

Rationale:
Benzodiazepines (lorazepam, diazepam, chlordiazepoxide) are first-line for alcohol withdrawal to prevent seizures and delirium tremens. Naloxone is for opioid overdose; methadone for opioid dependence; antipsychotics are adjunctive only.

Question 28

The most appropriate nursing action during oropharyngeal suctioning of a patient is:

A. Use a clean technique
B. Apply suction as you introduce the suction catheter
C. Continuously apply suction until all the secretions are removed
D. Flush the catheter with saline between catheter insertion

✓ Correct Answer: D

Rationale:
The catheter should be flushed with sterile saline between passes to clear secretions and maintain patency. Suction is applied only on withdrawal (not insertion); continuous suction risks hypoxia and mucosal trauma; sterile technique is preferred for deeper suctioning.

Question 29

To assess the function of the trigeminal and facial nerves, the nurse should:

A. Check for unilateral facial droop
B. Shine light into the client's eyes
C. Touch a cotton wisp strand to the cornea
D. Ask the client to clench their teeth and then attempt to separate the jaw by pushing down on the chin

✓ Correct Answer: C

Rationale:
The corneal reflex tests both CN V (afferent – trigeminal) and CN VII (efferent – facial). Clenching tests motor V; facial droop tests VII motor; light tests CN II/III.

Question 30

Which one of the following is an anti-craving agent used in the management of alcohol dependence syndrome?

A. Disulfiram
B. Lorazepam
C. Citrate calcium carbamide
D. Acamprosate

✓ Correct Answer: D

Rationale:
Acamprosate is an anti-craving agent that modulates glutamate and reduces the desire to drink. Disulfiram and calcium carbimide are aversive agents; lorazepam treats withdrawal, not craving.

Question 31

According to DSM-V classification, antisocial personality disorder is classified under:

A. Cluster A
B. Cluster B
C. Cluster C
D. Cluster D

✓ Correct Answer: B

Rationale:
Cluster B (dramatic, emotional, erratic) includes antisocial, borderline, histrionic, and narcissistic personality disorders. Cluster A = odd/eccentric; Cluster C = anxious/fearful.

Question 32

Pathological fire-setting is also called:

A. Pyromania
B. Kleptomania
C. Trichotillomania
D. Dipsomania

✓ Correct Answer: A

Rationale:
Pyromania is the impulse-control disorder characterized by deliberate and purposeful fire-setting. Kleptomania = stealing; trichotillomania = hair-pulling; dipsomania = alcohol craving.

Question 33

Narcissistic personality disorder is characterized by:

A. Callous unconcern for feelings of others
B. Self-dramatization, theatricality
C. Ideas of grandiosity and inflated sense of self importance
D. Significant and persistent disturbances of self-identity

✓ Correct Answer: C

Rationale:
Narcissistic PD features grandiosity, need for admiration, and lack of empathy. Callousness is more antisocial; theatricality is histrionic; identity disturbance is borderline.

Question 34

Antisocial personality disorders are characterized by all except:

A. Low frustration tolerance
B. Drug abuse
C. Learning from past experience
D. Criminal activities

✓ Correct Answer: C

Rationale:
A hallmark of antisocial PD is failure to learn from experience or punishment; individuals continue the same harmful patterns. Low frustration tolerance, substance abuse, and criminality are common features.

Question 35

The following are clinical features of mania except:

A. Elation of mood
B. Disorientation
C. Pressure of speech
D. Grandiose delusions

✓ Correct Answer: B

Rationale:
Mania features elevated/irritable mood, pressured speech, grandiosity, decreased need for sleep, and flight of ideas. Orientation is usually preserved (unlike delirium). Disorientation suggests organic pathology.

Question 36

Analysis of variance (ANOVA) test is used to:

A. Test the difference between two independent groups' means
B. Test the difference between two related groups' means
C. Test the difference among the means of 2+ independent groups
D. Test the difference in ranks of scores 2+ independent groups

✓ Correct Answer: C

Rationale:
ANOVA compares means of three or more independent groups. Two independent means use t-test; related means use paired t-test; ranks use non-parametric tests (Kruskal-Wallis).

Question 37

Persistent or recurrent tendency to observe unsuspecting persons naked, disrobing or engaged in sexual activity is termed as:

A. Exhibitionism
B. Sadism
C. Voyeurism
D. Frotteurism

✓ Correct Answer: C

Rationale:
Voyeurism is the paraphilia of observing unsuspecting people who are naked or engaged in sexual activity. Exhibitionism = exposing oneself; frotteurism = touching/rubbing against non-consenting persons.

Question 38

Systematic desensitization consists of the following except:

A. Relaxation training
B. Hierarchy construction
C. Desensitization of stimulus
D. Resolving unconscious conflicts

✓ Correct Answer: D

Rationale:
Systematic desensitization (Wolpe) is a behavioral technique involving relaxation, hierarchy of feared stimuli, and gradual exposure. Resolving unconscious conflicts is a psychoanalytic goal, not part of this method.

Question 39

Which one of the following is considered to be the most effective treatment of choice for phobic disorders?

A. Behaviour therapy
B. Interpersonal psychotherapy
C. Family therapy
D. Psychoanalysis

✓ Correct Answer: A

Rationale:
Exposure-based behavioral therapies (systematic desensitization, flooding, graded exposure) are the most effective evidence-based treatments for specific phobias and social anxiety disorder.

Question 40

Diffuse, bilaterally synchronous 3 Hz wave and spike discharges are typical EEG findings of:

A. Absence seizures
B. Infantile spasms
C. Myoclonic seizures
D. Complex partial seizures

✓ Correct Answer: A

Rationale:
Classic absence (petit mal) seizures show 3 Hz spike-and-wave discharges that are bilateral and synchronous. Infantile spasms show hypsarrhythmia; other seizures have different patterns.

Question 41

Limbic system includes all of the following except:

A. Amygdala
B. Hippocampal gyri
C. Anterior cingulate gyrus
D. Basal ganglia

✓ Correct Answer: D

Rationale:
The limbic system includes the amygdala, hippocampus, cingulate gyrus, fornix, and related structures involved in emotion and memory. Basal ganglia are extrapyramidal motor structures, not part of the limbic system.

Question 42

Dopamine pathways in the brain include all except:

A. Nigrostriatal pathway
B. Mesocorticolimbic pathway
C. Mesopontine complex
D. Tuberohypophyseal pathway

✓ Correct Answer: C

Rationale:
The four major dopaminergic pathways are nigrostriatal, mesolimbic, mesocortical, and tuberoinfundibular (tuberohypophyseal). There is no standard 'mesopontine' dopamine pathway in this context.

Question 43

A systematic method for continuous monitoring of diseases in a population in order to be able to detect changes in disease patterns and then to control them is:

A. Conditional probability
B. Screening
C. Prevalence
D. Surveillance

✓ Correct Answer: D

Rationale:
Public-health surveillance is the ongoing, systematic collection, analysis, and interpretation of health data for planning, implementation, and evaluation of public-health practice.

Question 44

Prevalence of disease in a community can be found out by:

A. Case control study
B. Cohort study
C. Cross-sectional study
D. Experimental study

✓ Correct Answer: C

Rationale:
Cross-sectional (prevalence) studies measure the frequency of a disease or characteristic in a population at a single point in time. Case-control and cohort studies measure association/risk; experimental studies test interventions.

Question 45

Descriptive epidemiology is study in relation to:

A. Time
B. Place
C. Person
D. All of the above

✓ Correct Answer: D

Rationale:
Descriptive epidemiology characterizes the distribution of disease by person, place, and time – the classic triad used to generate hypotheses.

Question 46

The correct loading dose for quinine therapy in severe malaria for an 8 kg infant is:

A. 40 mg
B. 50 mg
C. 100 mg
D. 160 mg

✓ Correct Answer: D

Rationale:
WHO loading dose of quinine is 20 mg salt/kg. For an 8 kg infant: 20 × 8 = 160 mg. (Subsequent maintenance is 10 mg/kg every 8 hours.)

Question 47

A correct discharge instruction to parents of a 12-month-old infant with gastroesophageal reflux disease will be:

A. Give the infant blended thinned feeds for easy digestion
B. Watch for blood-stained vomitus which requires urgent medical attention
C. Give prescribed proton pump inhibitors such as ranitidine to inhibit gastric acid secretion
D. Surgical management is the ultimate intervention for consideration

✓ Correct Answer: B

Rationale:
Parents must be taught red-flag symptoms (hematemesis, bilious vomiting, poor weight gain, respiratory distress) that require urgent review. Ranitidine is an H2-blocker, not a PPI; surgery is rarely first-line; feed thickening is sometimes used but not the priority teaching point.

Question 48

Primary irritant diaper dermatitis is:

A. The affected area is most commonly colonized by Candida albicans
B. Lesions appear as discrete follicular papules
C. The treatment of choice is low potency topical steroids
D. Plastic pants should be used in preference to diapers

✓ Correct Answer: C

Rationale:
Primary irritant contact dermatitis is treated by barrier creams, frequent changes, and short-term low-potency topical corticosteroids if inflamed. Candida superinfection produces satellite papules; plastic pants worsen occlusion.

Question 49

Phenothiazines are thought to exert their antipsychotic effect by:

A. Blocking post-synaptic dopamine receptors in the basal ganglia
B. Enhancing dopamine-mediated transmission of neural impulses at the synapses
C. Altering sodium metabolism within muscle cells and nerves
D. Selectively inhibiting the re-uptake of amines such as norepinephrine

✓ Correct Answer: A

Rationale:
Typical antipsychotics (phenothiazines) block D2 dopamine receptors in the mesolimbic pathway (therapeutic) and nigrostriatal pathway (extrapyramidal side effects). They do not enhance dopamine transmission.

Question 50

A priority nursing diagnosis for a 14-year-old teenager admitted with bulimia nervosa is:

A. High risk for fluid deficit related to abuse of laxatives and diuretics

B. Disturbed self esteem related to dysfunctional family system

C. Body image disturbance related to retarded ego development

D. Ineffective denial related to fear of losing an aspect of life she has control over

✓ Correct Answer: A

Rationale:

Electrolyte imbalance and fluid volume deficit from purging (vomiting, laxatives, diuretics) are immediate physiological threats and therefore the priority nursing diagnosis (ABCs / Maslow). Psychological diagnoses are important but secondary in the acute phase.

End of Batch 8 — Review the rationales carefully. Understanding the 'why' is more important than memorizing the letter. Good luck with your NCK examination!