

Power sums and special values of L-functions

Tung T. Nguyen

Western University, Algebra Seminar

March 5, 2021

Plans

- Some historical motivations.

Plans

- Some historical motivations.
- Hurwitz zeta functions.

Plans

- Some historical motivations.
- Hurwitz zeta functions.
- Special values of Hurwitz zeta functions.

- Some historical motivations.
- Hurwitz zeta functions.
- Special values of Hurwitz zeta functions.
- Some further topics for future investigations.

Pythagorean triples

A Pythagorean triple consists of three positive integers x, y, z such that $x^2 + y^2 = z^2$.

Pythagorean triples

A Pythagorean triple consists of three positive integers x, y, z such that $x^2 + y^2 = z^2$.

For example $(3, 4, 5)$ is a Pythagorean triple because

$$5^2 = 3^2 + 4^2.$$

Pythagorean triples

A Pythagorean triple consists of three positive integers x, y, z such that $x^2 + y^2 = z^2$.

For example $(3, 4, 5)$ is a Pythagorean triple because

$$5^2 = 3^2 + 4^2.$$

We can find all these triples by an elementary argument starting with

$$y^2 = z^2 - x^2 = (z - x)(z + x).$$

Pythagorean triples


A Pythagorean triple consists of three positive integers x, y, z such that $x^2 + y^2 = z^2$.

For example $(3, 4, 5)$ is a Pythagorean triple because

$$5^2 = 3^2 + 4^2.$$

We can find all these triples by an elementary argument starting with

$$y^2 = z^2 - x^2 = (z - x)(z + x).$$



A motivation: Fermat last theorem

- Fermat last theorem says that for odd prime p , the equation

$$x^p + y^p = z^p,$$

has no non-trivial solutions.

A motivation: Fermat last theorem

- Fermat last theorem says that for odd prime p , the equation

$$x^p + y^p = z^p,$$

has no non-trivial solutions.

- In the 19th century, Kummer “almost” solved this problem by using the following factorization in $\mathbb{Z}[\zeta_p]$

$$x^p = (z^p - y^p) = \prod_{i=0}^{p-1} (z - \zeta^i y).$$

A motivation: Fermat last theorem

- Fermat last theorem says that for odd prime p , the equation

$$x^p + y^p = z^p,$$

has no non-trivial solutions.

- In the 19th century, Kummer “almost” solved this problem by using the following factorization in $\mathbb{Z}[\zeta_p]$

$$x^p = (z^p - y^p) = \prod_{i=0}^{p-1} (z - \zeta^i y).$$

- The issue is that $\mathbb{Z}[\zeta_p]$ might not have the unique factorization property.

A motivation: Fermat last theorem

- Fermat last theorem says that for odd prime p , the equation

$$x^p + y^p = z^p,$$

has no non-trivial solutions.

- In the 19th century, Kummer “almost” solved this problem by using the following factorization in $\mathbb{Z}[\zeta_p]$

$$x^p = (z^p - y^p) = \prod_{i=0}^{p-1} (z - \zeta^i y).$$

- The issue is that $\mathbb{Z}[\zeta_p]$ might not have the unique factorization property.
- The class number h of $\mathbb{Z}[\zeta_p]$ measures the failure of the unique factorization property of $\mathbb{Z}[\zeta_p]$. When $p \nmid h$, the above elementary approach works.

A motivation: Fermat last theorem

- Fermat last theorem says that for odd prime p , the equation

$$x^p + y^p = z^p,$$

has no non-trivial solutions.

- In the 19th century, Kummer “almost” solved this problem by using the following factorization in $\mathbb{Z}[\zeta_p]$

$$x^p = (z^p - y^p) = \prod_{i=0}^{p-1} (z - \zeta^i y).$$

- The issue is that $\mathbb{Z}[\zeta_p]$ might not have the unique factorization property.
- The class number h of $\mathbb{Z}[\zeta_p]$ measures the failure of the unique factorization property of $\mathbb{Z}[\zeta_p]$. When $p \nmid h$, the above elementary approach works.

Class number formula-The algebraic side

Let K be a number field and \mathcal{O}_K be its ring of algebraic integers.

Let

Class number formula-The algebraic side

Let K be a number field and \mathcal{O}_K be its ring of algebraic integers.

Let

- r_1 (respectively r_2) be the number of real (respectively complex) places of K .

Class number formula-The algebraic side

Let K be a number field and \mathcal{O}_K be its ring of algebraic integers.

Let

- r_1 (respectively r_2) be the number of real (respectively complex) places of K .
- h_K : class number of K .

Class number formula-The algebraic side

Let K be a number field and \mathcal{O}_K be its ring of algebraic integers.

Let

- r_1 (respectively r_2) be the number of real (respectively complex) places of K .
- h_K : class number of K .
- w_K : number of roots of unity in K .

Class number formula-The algebraic side

Let K be a number field and \mathcal{O}_K be its ring of algebraic integers.

Let

- r_1 (respectively r_2) be the number of real (respectively complex) places of K .
- h_K : class number of K .
- w_K : number of roots of unity in K .
- R_K : the regulator of K .

Class number formula-The algebraic side

Let K be a number field and \mathcal{O}_K be its ring of algebraic integers.

Let

- r_1 (respectively r_2) be the number of real (respectively complex) places of K .
- h_K : class number of K .
- w_K : number of roots of unity in K .
- R_K : the regulator of K .
- D_K : the discriminant of K .

Class number formula-The algebraic side

Let K be a number field and \mathcal{O}_K be its ring of algebraic integers.

Let

- r_1 (respectively r_2) be the number of real (respectively complex) places of K .
- h_K : class number of K .
- w_K : number of roots of unity in K .
- R_K : the regulator of K .
- D_K : the discriminant of K .

These numbers contain arithmetic information about K .

Class number formula-The analytic side

The Dedekind zeta function for K is defined by

$$\zeta_K(s) = \sum_{\mathfrak{a} \subset \mathcal{O}_K} \frac{1}{N(\mathfrak{a})^s}.$$

It has a meromorphic continuation to \mathbb{C} with a unique simple pole at $s = 1$.

Class number formula-The analytic side

The Dedekind zeta function for K is defined by

$$\zeta_K(s) = \sum_{\mathfrak{a} \subset \mathcal{O}_K} \frac{1}{N(\mathfrak{a})^s}.$$

It has a meromorphic continuation to \mathbb{C} with a unique simple pole at $s = 1$.

Theorem (Class number formula)

$$\lim_{s \rightarrow 1} (s - 1) \zeta_K(s) = \frac{2^{r_1} (2\pi)^{r_2} R_K h_K}{w_K \sqrt{|D_K|}}.$$

Class number formula-The analytic side

The Dedekind zeta function for K is defined by

$$\zeta_K(s) = \sum_{\mathfrak{a} \subset \mathcal{O}_K} \frac{1}{N(\mathfrak{a})^s}.$$

It has a meromorphic continuation to \mathbb{C} with a unique simple pole at $s = 1$.

Theorem (Class number formula)

$$\lim_{s \rightarrow 1} (s - 1) \zeta_K(s) = \frac{2^{r_1} (2\pi)^{r_2} R_K h_K}{w_K \sqrt{|D_K|}}.$$

This formula provides a connection between the algebraic world and the analytic world.

Euler's discoveries

- In 1734, Leonhard Euler found the following remarkable formula

$$\zeta(2) = \zeta_{\mathbb{Q}}(2) = 1 + \frac{1}{2^2} + \frac{1}{3^2} + \dots = \frac{\pi^2}{6}.$$

Euler's discoveries

- In 1734, Leonhard Euler found the following remarkable formula

$$\zeta(2) = \zeta_{\mathbb{Q}}(2) = 1 + \frac{1}{2^2} + \frac{1}{3^2} + \dots = \frac{\pi^2}{6}.$$

- Indeed, Euler did much more. In particular, he showed that

$$\zeta(2k) = (-1)^{k+1} \frac{(2\pi)^{2k}}{2(2k)!} B_{2k},$$

where $\{B_n\}$ are the [Bernoulli numbers](#) defined by following Taylor's expansion

$$\frac{z}{e^z - 1} = \sum_{n=0}^{\infty} \frac{B_n}{n!} z^n.$$

Euler's discoveries

- In 1734, Leonhard Euler found the following remarkable formula

$$\zeta(2) = \zeta_{\mathbb{Q}}(2) = 1 + \frac{1}{2^2} + \frac{1}{3^2} + \dots = \frac{\pi^2}{6}.$$

- Indeed, Euler did much more. In particular, he showed that

$$\zeta(2k) = (-1)^{k+1} \frac{(2\pi)^{2k}}{2(2k)!} B_{2k},$$

where $\{B_n\}$ are the [Bernoulli numbers](#) defined by following Taylor's expansion

$$\frac{z}{e^z - 1} = \sum_{n=0}^{\infty} \frac{B_n}{n!} z^n.$$

- Using the [functional equation](#) for $\zeta(s)$ we have

$$\zeta(1-n) = \zeta(-n) = (-1)^{n-1} \frac{B_n}{n}.$$

Three phases of understanding of zeta values (according to K. Kato)

We see from the previous slide that $\zeta(1 - n)$ is a rational number.

Three phases of understanding of zeta values (according to K. Kato)

We see from the previous slide that $\zeta(1 - n)$ is a rational number. Generally speaking, the study of zeta values could be divided into three phases.

Three phases of understanding of zeta values (according to K. Kato)

We see from the previous slide that $\zeta(1 - n)$ is a rational number. Generally speaking, the study of zeta values could be divided into three phases.

1. Rationality of zeta values.

Three phases of understanding of zeta values (according to K. Kato)

We see from the previous slide that $\zeta(1 - n)$ is a rational number. Generally speaking, the study of zeta values could be divided into three phases.

1. Rationality of zeta values.
2. p -adic properties of zeta values.

Three phases of understanding of zeta values (according to K. Kato)

We see from the previous slide that $\zeta(1 - n)$ is a rational number. Generally speaking, the study of zeta values could be divided into three phases.

1. Rationality of zeta values.
2. p -adic properties of zeta values.
3. Arithmetic significances of zeta values.

Three phases of understanding of zeta values (according to K. Kato)

We see from the previous slide that $\zeta(1 - n)$ is a rational number. Generally speaking, the study of zeta values could be divided into three phases.

1. Rationality of zeta values.
2. p -adic properties of zeta values.
3. Arithmetic significances of zeta values.

Today talk is mostly about phase 1 and a little bit about phase 2 and phase 3.

The generalized class number formula (The Bloch-Kato's conjecture)

- Bloch-Kato made a precise conjecture about the general form of the class number formula. Their work generalizes the previous work of Beilinson, Deligne, Jannsen and many others.

The generalized class number formula (The Bloch-Kato's conjecture)

- Bloch-Kato made a precise conjecture about the general form of the class number formula. Their work generalizes the previous work of Beilinson, Deligne, Jannsen and many others.
- Roughly speaking, the generalized class number formula says that the many arithmetic information of an algebraic variety X can be read off from its L -function.

The generalized class number formula (The Bloch-Kato's conjecture)

- Bloch-Kato made a precise conjecture about the general form of the class number formula. Their work generalizes the previous work of Beilinson, Deligne, Jannsen and many others.
- Roughly speaking, the generalized class number formula says that the many arithmetic information of an algebraic variety X can be read off from its L -function.
- For example, if X is given by an cubic equation of the form

$$y^2 = x^3 + ax + b,$$

with $a, b \in \mathbb{Q}$ then the L -function $L(E, s)$ can conjecturally tell how large $X(\mathbb{Q})$ is. This is known as the Birch and Swinnerton-Dyer conjecture.

Hurwitz zeta function

- In 1882, Hurwitz defined the following infinite series

$$\zeta(s, a) = \sum_{n=0}^{\infty} \frac{1}{(n+a)^s},$$

where $0 < a \leq 1$.

Hurwitz zeta function

- In 1882, Hurwitz defined the following infinite series

$$\zeta(s, a) = \sum_{n=0}^{\infty} \frac{1}{(n+a)^s},$$

where $0 < a \leq 1$.

- When $a = 1$, we have $\zeta(s, 1) = \zeta(s)$.

Hurwitz zeta function

- In 1882, Hurwitz defined the following infinite series

$$\zeta(s, a) = \sum_{n=0}^{\infty} \frac{1}{(n+a)^s},$$

where $0 < a \leq 1$.

- When $a = 1$, we have $\zeta(s, 1) = \zeta(s)$.
- Like the Riemann zeta function, the Hurwitz zeta function has an analytic continuation to \mathbb{C} with a simple pole at $s = 1$.

Bernoulli polynomials

- Let $B_n(x)$ be the function given by the Taylor expansion

$$\frac{ze^{xz}}{e^z - 1} = \sum_{n=0}^{\infty} \frac{B_n(x)}{n!} z^n.$$

Bernoulli polynomials

- Let $B_n(x)$ be the function given by the Taylor expansion

$$\frac{ze^{xz}}{e^z - 1} = \sum_{n=0}^{\infty} \frac{B_n(x)}{n!} z^n.$$

- It turns out that $B_n(x)$ is a polynomial. In fact

$$B_n(x) = \sum_{k=0}^n \binom{n}{k} B_k(0) x^{n-k}.$$

Bernoulli polynomials

- Let $B_n(x)$ be the function given by the Taylor expansion

$$\frac{ze^{xz}}{e^z - 1} = \sum_{n=0}^{\infty} \frac{B_n(x)}{n!} z^n.$$

- It turns out that $B_n(x)$ is a polynomial. In fact

$$B_n(x) = \sum_{k=0}^n \binom{n}{k} B_k(0) x^{n-k}.$$

This can be seen by looking at the Taylor expansion

$$\frac{ze^{xz}}{e^z - 1} = e^{xz} \times \frac{z}{e^z - 1} = \left[\sum_{m=0}^{\infty} \frac{x^m}{m!} z^m \right] \times \left[\sum_{n=0}^{\infty} \frac{B_n(0)}{n!} z^n \right]$$

- $B_n(x)$ is called the n -th Bernoulli polynomial.

Special values of Hurwitz zeta functions, I

Using contour integrals, we can show that

Theorem (Special values of Hurwitz zeta functions)

For $n \geq 0$

$$\zeta(-n, a) = -\frac{B_{n+1}(a)}{n+1}.$$

Some examples of Bernoulli numbers

n	0	1	2	3	4	5	6	7	8	9	10	11	12
B_n	1	$-\frac{1}{2}$	$\frac{1}{6}$	0	$-\frac{1}{30}$	0	$\frac{1}{42}$	0	$-\frac{1}{30}$	0	$\frac{5}{66}$	0	$-\frac{691}{2730}$

Special values of Hurwitz zeta functions, II

The proof for special values of Hurwitz zeta functions uses complex analysis-which is not elementary. We explain another method to derive this formula without advanced tools.

Special values of Hurwitz zeta functions, II

The proof for special values of Hurwitz zeta functions uses complex analysis-which is not elementary. We explain another method to derive this formula without advanced tools. For this, we introduce the power-sum polynomials.

Special values of Hurwitz zeta functions, II

The proof for special values of Hurwitz zeta functions uses complex analysis-which is not elementary. We explain another method to derive this formula without advanced tools. For this, we introduce the power-sum polynomials.

Let us remind ourselves some familiar formulas in Calculus 1.

$$1 + 2 + \dots + (M - 1) =$$

Special values of Hurwitz zeta functions, II

The proof for special values of Hurwitz zeta functions uses complex analysis-which is not elementary. We explain another method to derive this formula without advanced tools. For this, we introduce the power-sum polynomials.

Let us remind ourselves some familiar formulas in Calculus 1.

$$1 + 2 + \dots + (M - 1) = \frac{M(M - 1)}{2}.$$

What about sum of the first M squares?

$$1^2 + 2^2 + \dots + (M - 1)^2 =$$

Special values of Hurwitz zeta functions, II

The proof for special values of Hurwitz zeta functions uses complex analysis-which is not elementary. We explain another method to derive this formula without advanced tools. For this, we introduce the power-sum polynomials.

Let us remind ourselves some familiar formulas in Calculus 1.

$$1 + 2 + \dots + (M - 1) = \frac{M(M - 1)}{2}.$$

What about sum of the first M squares?

$$1^2 + 2^2 + \dots + (M - 1)^2 = \frac{M(M - 1)(2M - 1)}{6}.$$

What about sum of the first n cubes?

$$1^3 + 2^3 + \dots + (M - 1)^3 =$$

Special values of Hurwitz zeta functions, II

The proof for special values of Hurwitz zeta functions uses complex analysis-which is not elementary. We explain another method to derive this formula without advanced tools. For this, we introduce the power-sum polynomials.

Let us remind ourselves some familiar formulas in Calculus 1.

$$1 + 2 + \dots + (M - 1) = \frac{M(M - 1)}{2}.$$

What about sum of the first M squares?

$$1^2 + 2^2 + \dots + (M - 1)^2 = \frac{M(M - 1)(2M - 1)}{6}.$$

What about sum of the first n cubes?

$$1^3 + 2^3 + \dots + (M - 1)^3 = \frac{M^2(M - 1)^2}{4}.$$

Power sum polynomials

From these numerical data, we can predict that for each positive integer k

$$S_n(M) = 1^k + 2^k + \dots + (M-1)^k,$$

is a polynomial in M of degree $k+1$.

Power sum polynomials

From these numerical data, we can predict that for each positive integer k

$$S_n(M) = 1^k + 2^k + \dots + (M-1)^k,$$

is a polynomial in M of degree $k+1$. This prediction turns out to be correct! In fact, we have the following theorem.

Theorem

$$S_n(x) = \frac{B_{n+1}(x) - B_{n+1}(1)}{n+1}.$$

Minac's theorem

In 1994, Jan Minac observed the following

$$\int_0^1 S_1(x) dx = \int_0^1 \frac{x(x-1)}{2} dx = -\frac{1}{12}.$$

Minac's theorem

In 1994, Jan Minac observed the following

$$\int_0^1 S_1(x) dx = \int_0^1 \frac{x(x-1)}{2} dx = -\frac{1}{12}.$$

$$\int_0^1 S_2(x) dx = \int_0^1 \frac{x(x-1)(2x-1)}{6} dx = 0.$$

Minac's theorem

In 1994, Jan Minac observed the following

$$\int_0^1 S_1(x) dx = \int_0^1 \frac{x(x-1)}{2} dx = -\frac{1}{12}.$$

$$\int_0^1 S_2(x) dx = \int_0^1 \frac{x(x-1)(2x-1)}{6} dx = 0.$$

$$\int_0^1 S_3(x) dx = \int_0^1 \frac{x^2(x-1)^2}{4} dx = \frac{1}{120}.$$

He observed that the numbers on the right hand sides are exactly zeta values $\zeta(-1), \zeta(-2), \zeta(-3)$.

From this observation, he proved that

Theorem

$$\zeta(-n) = \int_0^1 S_n(x) dx.$$

The first proof

There are two proofs for Minac's theorem.

The first proof

There are two proofs for Minac's theorem. The first one uses Bernoulli polynomials.

$$\begin{aligned}\int_0^1 S_n(x) dx &= \int_0^1 \frac{B_{n+1}(x) - B_{n+1}(1)}{n+1} dx \\ &= \int_0^1 \frac{B_{n+1}(x)}{n+1} dx - \frac{B_{n+1}(1)}{n+1}.\end{aligned}$$

The first proof

There are two proofs for Minac's theorem. The first one uses Bernoulli polynomials.

$$\begin{aligned}\int_0^1 S_n(x) dx &= \int_0^1 \frac{B_{n+1}(x) - B_{n+1}(1)}{n+1} dx \\ &= \int_0^1 \frac{B_{n+1}(x)}{n+1} dx - \frac{B_{n+1}(1)}{n+1}.\end{aligned}$$

It is known that $\int_a^{a+1} B_{n+1}(x) dx = a^{n+1}$ for all a . In particular, $\int_0^1 B_{n+1}(x) dx = 0$.

The first proof

There are two proofs for Minac's theorem. The first one uses Bernoulli polynomials.

$$\begin{aligned}\int_0^1 S_n(x) dx &= \int_0^1 \frac{B_{n+1}(x) - B_{n+1}(1)}{n+1} dx \\ &= \int_0^1 \frac{B_{n+1}(x)}{n+1} dx - \frac{B_{n+1}(1)}{n+1}.\end{aligned}$$

It is known that $\int_a^{a+1} B_{n+1}(x) dx = a^{n+1}$ for all a . In particular, $\int_0^1 B_{n+1}(x) dx = 0$. Additionally, by Euler's theorem

$$\zeta(-n) = -\frac{B_{n+1}(1)}{n+1}.$$

Combining these two facts, we have

$$\zeta(-n) = \int_0^1 S_n(x) dx.$$

The second proof

We can also prove Minac's theorem without using Bernoulli polynomials.

The second proof

We can also prove Minac's theorem without using Bernoulli polynomials. Instead, we use the following identity

$$1 = \sum_{q=0}^{\infty} \frac{(s-1) \dots (s+q-1)}{(q+1)!} (\zeta(s+q) - 1).$$

The second proof

We can also prove Minac's theorem without using Bernoulli polynomials. Instead, we use the following identity

$$1 = \sum_{q=0}^{\infty} \frac{(s-1) \dots (s+q-1)}{(q+1)!} (\zeta(s+q) - 1).$$

By plugging $s = -1, -2, \dots$, we get some combinatorial relations between $\zeta(-n)$ for $n \geq 0$.

The second proof

We can also prove Minac's theorem without using Bernoulli polynomials. Instead, we use the following identity

$$1 = \sum_{q=0}^{\infty} \frac{(s-1) \dots (s+q-1)}{(q+1)!} (\zeta(s+q) - 1).$$

By plugging $s = -1, -2, \dots$, we get some combinatorial relations between $\zeta(-n)$ for $n \geq 0$. Using the collapsing sum method, we have

$$\begin{aligned} (M-1)^{n+1} &= 1 + \sum_{m=2}^{M-1} (m^{n+1} - (m-1)^{n+1}) \\ &= \sum_{k=0}^n \binom{n+1}{k+1} (-1)^k S_{n-k}(M). \end{aligned}$$

The second proof

We can also prove Minac's theorem without using Bernoulli polynomials. Instead, we use the following identity

$$1 = \sum_{q=0}^{\infty} \frac{(s-1) \dots (s+q-1)}{(q+1)!} (\zeta(s+q) - 1).$$

By plugging $s = -1, -2, \dots$, we get some combinatorial relations between $\zeta(-n)$ for $n \geq 0$. Using the collapsing sum method, we have

$$\begin{aligned}(M-1)^{n+1} &= 1 + \sum_{m=2}^{M-1} (m^{n+1} - (m-1)^{n+1}) \\ &= \sum_{k=0}^n \binom{n+1}{k+1} (-1)^k S_{n-k}(M).\end{aligned}$$

Integrating both sides from 0 to 1, we easily see that $\int_0^1 S_n(x) dx$ satisfies a similar combinatorial relation.

Generalized power sums and special values of the Hurwitz zeta functions

Recently, we generalized Minac's theorem for the Hurwitz zeta functions. More precisely, let $S_{n,a}$ is the generalized power sum defined by

$$S_{n,a}(M) = a^n + (1+a)^n + (2+a)^n + \dots + (M+a-2)^n.$$

Generalized power sums and special values of the Hurwitz zeta functions

Recently, we generalized Minac's theorem for the Hurwitz zeta functions. More precisely, let $S_{n,a}$ is the generalized power sum defined by

$$S_{n,a}(M) = a^n + (1+a)^n + (2+a)^n + \dots + (M+a-2)^n.$$

Then we have the following

Theorem

$$\zeta(-n, a) = \int_{1-a}^{2-a} S_{n,a}(x) dx,$$

Generalized power sums and special values of the Hurwitz zeta functions

Recently, we generalized Minac's theorem for the Hurwitz zeta functions. More precisely, let $S_{n,a}$ is the generalized power sum defined by

$$S_{n,a}(M) = a^n + (1+a)^n + (2+a)^n + \dots + (M+a-2)^n.$$

Then we have the following

Theorem

$$\zeta(-n, a) = \int_{1-a}^{2-a} S_{n,a}(x) dx,$$

Note that when $a = 1$, this gives a direct generalization of Minac's theorem.

Sketchs of our proofs

We gave three different proofs for the above theorem.

- The first proof uses Bernoulli polynomials.

Sketchs of our proofs

We gave three different proofs for the above theorem.

- The first proof uses Bernoulli polynomials.
- The second proof uses a recursive relation between Hurwitz zeta values and classical zeta values.

Sketchs of our proofs

We gave three different proofs for the above theorem.

- The first proof uses Bernoulli polynomials.
- The second proof uses a recursive relation between Hurwitz zeta values and classical zeta values. More precisely, using the Taylor expansion of the term $\frac{1}{(n+a)^s}$, we have

$$\zeta(-n, a) = a^n - \frac{a^{n+1}}{n+1} + \sum_{k=0}^n \binom{n}{k} a^k \zeta(k-n).$$

Sketchs of our proofs

We gave three different proofs for the above theorem.

- The first proof uses Bernoulli polynomials.
- The second proof uses a recursive relation between Hurwitz zeta values and classical zeta values. More precisely, using the Taylor expansion of the term $\frac{1}{(n+a)^s}$, we have

$$\zeta(-n, a) = a^n - \frac{a^{n+1}}{n+1} + \sum_{k=0}^n \binom{n}{k} a^k \zeta(k-n).$$

Additionally, $S_{n,a}(x) = a^n S_0(x) + \sum_{k=0}^{n-1} \binom{n}{k} a^k S_{n-k}(x-1).$

Sketchs of our proofs

We gave three different proofs for the above theorem.

- The first proof uses Bernoulli polynomials.
- The second proof uses a recursive relation between Hurwitz zeta values and classical zeta values. More precisely, using the Taylor expansion of the term $\frac{1}{(n+a)^s}$, we have

$$\zeta(-n, a) = a^n - \frac{a^{n+1}}{n+1} + \sum_{k=0}^n \binom{n}{k} a^k \zeta(k-n).$$

Additionally, $S_{n,a}(x) = a^n S_0(x) + \sum_{k=0}^{n-1} \binom{n}{k} a^k S_{n-k}(x-1)$.

- We see that $\int_{1-a}^{2-a} S_{n,a}(x) dx$ satisfies a similar recursive relation.

Sketchs of our proofs

For the third proof, we use the recursive relation between Hurwitz zeta values.

Sketchs of our proofs

For the third proof, we use the recursive relation between Hurwitz zeta values. The key idea is the following observation

- We have

$$\begin{aligned}\frac{1}{(n+a-1)^{s-1}} - \frac{1}{(n+a)^{s-1}} &= \frac{1}{(n+a)^s} \left(\left(1 - \frac{1}{n+a}\right)^{-(s-1)} - 1 \right) \\ &= \sum_{k=0}^{\infty} \frac{\Gamma(s+k)}{\Gamma(s-1)(k+1)!} \frac{1}{(a+n)^{s+k}}.\end{aligned}$$

Sketchs of our proofs

For the third proof, we use the recursive relation between Hurwitz zeta values. The key idea is the following observation

- We have

$$\begin{aligned}\frac{1}{(n+a-1)^{s-1}} - \frac{1}{(n+a)^{s-1}} &= \frac{1}{(n+a)^s} \left(\left(1 - \frac{1}{n+a}\right)^{-(s-1)} - 1 \right) \\ &= \sum_{k=0}^{\infty} \frac{\Gamma(s+k)}{\Gamma(s-1)(k+1)!} \frac{1}{(a+n)^{s+k}}.\end{aligned}$$

- Using the collapsing sum, we have

$$\frac{1}{a^{s-1}} = \sum_{k=0}^{\infty} \frac{\Gamma(s+k)}{\Gamma(s-1)(k+1)!} \left(\zeta(s+k, a) - \frac{1}{a^{s+k}} \right).$$

Sketchs of our proofs

For the third proof, we use the recursive relation between Hurwitz zeta values. The key idea is the following observation

- We have

$$\begin{aligned}\frac{1}{(n+a-1)^{s-1}} - \frac{1}{(n+a)^{s-1}} &= \frac{1}{(n+a)^s} \left(\left(1 - \frac{1}{n+a}\right)^{-(s-1)} - 1 \right) \\ &= \sum_{k=0}^{\infty} \frac{\Gamma(s+k)}{\Gamma(s-1)(k+1)!} \frac{1}{(a+n)^{s+k}}.\end{aligned}$$

- Using the collapsing sum, we have

$$\frac{1}{a^{s-1}} = \sum_{k=0}^{\infty} \frac{\Gamma(s+k)}{\Gamma(s-1)(k+1)!} \left(\zeta(s+k, a) - \frac{1}{a^{s+k}} \right).$$

- Using the same collapsing technique, we can show that $\int_{1-a}^{2-a} S_{n,a}(x) dx$ satisfies a similar recursive relation.

p-adic properties of zeta values

By Fermat's little theorem, we know that if $s \equiv t \pmod{p-1}$ then

$$n^s \equiv n^t \pmod{p}.$$

p-adic properties of zeta values

By Fermat's little theorem, we know that if $s \equiv t \pmod{p-1}$ then

$$n^s \equiv n^t \pmod{p}.$$

Ignoring convergence issue we have

$$\zeta(-s) = \sum_{n=1}^{\infty} n^s \equiv \sum_{n=1}^{\infty} n^t = \zeta(-t) \pmod{p}.$$

p-adic properties of zeta values

By Fermat's little theorem, we know that if $s \equiv t \pmod{p-1}$ then

$$n^s \equiv n^t \pmod{p}.$$

Ignoring convergence issue we have

$$\zeta(-s) = \sum_{n=1}^{\infty} n^s \equiv \sum_{n=1}^{\infty} n^t = \zeta(-t) \pmod{p}.$$

Amazingly, we can make this intuitive argument become rigorous.

p-adic properties of zeta values

By Fermat's little theorem, we know that if $s \equiv t \pmod{p-1}$ then

$$n^s \equiv n^t \pmod{p}.$$

Ignoring convergence issue we have

$$\zeta(-s) = \sum_{n=1}^{\infty} n^s \equiv \sum_{n=1}^{\infty} n^t = \zeta(-t) \pmod{p}.$$

Amazingly, we can make this intuitive argument become rigorous. In fact, we have the following theorem.

Theorem (Kummer)

Let r_1, r_2 be two positive integers. Suppose r_1 is not a multiple of $p-1$. If $r_1 \equiv r_2 \pmod{(p-1)p^{n-1}}$ then

$$(1 - p^{r_1-1})\zeta(1 - r_1) \equiv (1 - p^{r_2-1})\zeta(1 - r_2) \pmod{p^n}.$$

Arithmetic significance of zeta values

Recall that Fermat last theorem holds if $p \nmid h$ where h is the class number of $\mathbb{Z}[\zeta_p]$.

Arithmetic significance of zeta values

Recall that Fermat last theorem holds if $p \nmid h$ where h is the class number of $\mathbb{Z}[\zeta_p]$. The following criterion is due to Kummer.

Theorem (Kummer)

If $p \nmid \prod_{i=1}^{\frac{p-3}{2}} \zeta(1-2i)$ then $p \nmid h$.

Arithmetic significance of zeta values

Recall that Fermat last theorem holds if $p \nmid h$ where h is the class number of $\mathbb{Z}[\zeta_p]$. The following criterion is due to Kummer.

Theorem (Kummer)

If $p \nmid \prod_{i=1}^{\frac{p-3}{2}} \zeta(1-2i)$ then $p \nmid h$.

There is a more refined version due to Herbrand-Ribet.

Arithmetic significance of zeta values

Recall that Fermat last theorem holds if $p \nmid h$ where h is the class number of $\mathbb{Z}[\zeta_p]$. The following criterion is due to Kummer.

Theorem (Kummer)

If $p \nmid \prod_{i=1}^{\frac{p-3}{2}} \zeta(1-2i)$ then $p \nmid h$.

There is a more refined version due to Herbrand-Ribet.

Ribet's proof is quite interesting: it uses Galois representations attached modular forms and congruence between modular forms. Further generalizations of his method have led to a proof of Fermat last theorem for all odd prime p (due to the work of Wiles, Taylor, and many others).

Future investigations

- It would be quite interesting to study p -adic properties of Hurwitz zeta values.

Future investigations

- It would be quite interesting to study p -adic properties of Hurwitz zeta values.
- Recent work by Lagarias shows that Hurwitz zeta functions are almost “motivic”. We can hope for a class number formula for Hurwitz zeta values.

Future investigations

- It would be quite interesting to study p -adic properties of Hurwitz zeta values.
- Recent work by Lagarias shows that Hurwitz zeta functions are almost “motivic”. We can hope for a class number formula for Hurwitz zeta values.
- In some ongoing joint projects with Lyle Muller, Jan Minac and many others, we are studying different aspects of zeta functions.

Future investigations

- It would be quite interesting to study p -adic properties of Hurwitz zeta values.
- Recent work by Lagarias shows that Hurwitz zeta functions are almost “motivic”. We can hope for a class number formula for Hurwitz zeta values.
- In some ongoing joint projects with Lyle Muller, Jan Minac and many others, we are studying different aspects of zeta functions. In particular, we are quite interested in the [Ihara zeta functions for graphs](#) and their applications to neuroscience.

Future investigations

- It would be quite interesting to study p -adic properties of Hurwitz zeta values.
- Recent work by Lagarias shows that Hurwitz zeta functions are almost “motivic”. We can hope for a class number formula for Hurwitz zeta values.
- In some ongoing joint projects with Lyle Muller, Jan Minac and many others, we are studying different aspects of zeta functions. In particular, we are quite interested in the [Ihara zeta functions for graphs](#) and their applications to neuroscience.
- The Galois groups play an important role in number theory. In joint work with Nguyen Duy Tan and Jan Minac, we are investigating the structures of [Galois groups over global fields](#).

Thank you!