

# Heights and Tamagawa numbers of motives

---

Tung T. Nguyen

Western University, Algebra Seminar

February 26, 2021

# Plans

- Some historical motivations.

# Plans

- Some historical motivations.
- Introduction to the Tamagawa number conjecture for motives.

# Plans

- Some historical motivations.
- Introduction to the Tamagawa number conjecture for motives.
- Heights of motives.

- Some historical motivations.
- Introduction to the Tamagawa number conjecture for motives.
- Heights of motives.
- Connection between heights and Tamagawa numbers of motives.

- Some historical motivations.
- Introduction to the Tamagawa number conjecture for motives.
- Heights of motives.
- Connection between heights and Tamagawa numbers of motives.
- Some ongoing joint work with Ján Mináč, Nguyễn Duy Tân, Lyle Muller and others.

## Euler's discoveries

- In 1734, Leonhard Euler found the following remarkable formula

$$1 + \frac{1}{2^2} + \frac{1}{3^2} + \dots = \frac{\pi^2}{6}.$$



## Euler's discoveries

- In 1734, Leonhard Euler found the following remarkable formula

$$1 + \frac{1}{2^2} + \frac{1}{3^2} + \dots = \frac{\pi^2}{6}.$$

- Indeed, Euler did much more. In particular, he showed that

$$1 + \frac{1}{2^{2k}} + \frac{1}{3^{2k}} + \dots = (-1)^{k+1} \frac{(2\pi)^{2k}}{2(2k)!} B_{2k},$$

where  $\{B_n\}$  are the Bernoulli numbers defined by following Taylor's expansion

$$\frac{z}{e^z - 1} = \sum_{n=0}^{\infty} \frac{B_n}{n!} z^n.$$

## Euler's discoveries

- In 1734, Leonhard Euler found the following remarkable formula

$$1 + \frac{1}{2^2} + \frac{1}{3^2} + \dots = \frac{\pi^2}{6}.$$

- Indeed, Euler did much more. In particular, he showed that

$$1 + \frac{1}{2^{2k}} + \frac{1}{3^{2k}} + \dots = (-1)^{k+1} \frac{(2\pi)^{2k}}{2(2k)!} B_{2k},$$

where  $\{B_n\}$  are the Bernoulli numbers defined by following Taylor's expansion

$$\frac{z}{e^z - 1} = \sum_{n=0}^{\infty} \frac{B_n}{n!} z^n.$$

- Euler's discoveries is the beginning of many wonderful and mysterious studies.

## Euler's discoveries from a modern perspective

In 1859, Bernhard Riemann studied the following (complex) function in his article “On the Number of Primes Less Than a Given Magnitude”

$$\zeta(s) = \sum_{n=1}^{\infty} n^{-s} = \frac{1}{1^s} + \frac{1}{2^s} + \frac{1}{3^s} + \cdots \quad .$$

## Euler's discoveries from a modern perspective

In 1859, Bernhard Riemann studied the following (complex) function in his article “On the Number of Primes Less Than a Given Magnitude”

$$\zeta(s) = \sum_{n=1}^{\infty} n^{-s} = \frac{1}{1^s} + \frac{1}{2^s} + \frac{1}{3^s} + \cdots \quad .$$

With this definition, Euler's formula could be written as

$$\zeta(2k) = (-1)^{k+1} \frac{(2\pi)^{2k}}{2(2k)!} B_{2k}.$$

## Euler's discoveries from a modern perspective

In 1859, Bernhard Riemann studied the following (complex) function in his article “On the Number of Primes Less Than a Given Magnitude”

$$\zeta(s) = \sum_{n=1}^{\infty} n^{-s} = \frac{1}{1^s} + \frac{1}{2^s} + \frac{1}{3^s} + \cdots \quad .$$

With this definition, Euler's formula could be written as

$$\zeta(2k) = (-1)^{k+1} \frac{(2\pi)^{2k}}{2(2k)!} B_{2k}.$$

Riemann shown that  $\zeta(s)$  can be extended to a **analytic function** on  $\mathbb{C}$  except for a simple pole at  $s = 1$ .

## Euler's discoveries from a modern perspective

In 1859, Bernhard Riemann studied the following (complex) function in his article “On the Number of Primes Less Than a Given Magnitude”

$$\zeta(s) = \sum_{n=1}^{\infty} n^{-s} = \frac{1}{1^s} + \frac{1}{2^s} + \frac{1}{3^s} + \cdots .$$

With this definition, Euler's formula could be written as

$$\zeta(2k) = (-1)^{k+1} \frac{(2\pi)^{2k}}{2(2k)!} B_{2k}.$$

Riemann shown that  $\zeta(s)$  can be extended to a **analytic function** on  $\mathbb{C}$  except for a simple pole at  $s = 1$ . Furthermore, it has a symmetric **functional equation**

$$\zeta(s) = 2^s \pi^{s-1} \sin\left(\frac{\pi s}{2}\right) \Gamma(1-s) \zeta(1-s).$$

## Euler's discoveries from a modern perspective

By Euler's formula, the values of the Riemann zeta functions at even integers are of the form

$$\zeta(2n) = (-1)^{k+1} \frac{(2\pi)^{2k}}{2(2k)!} B_{2k}.$$

## Euler's discoveries from a modern perspective

By Euler's formula, the values of the Riemann zeta functions at even integers are of the form

$$\zeta(2n) = (-1)^{k+1} \frac{(2\pi)^{2k}}{2(2k)!} B_{2k}.$$

Using the functional equation, we have

$$\zeta(1-n) = \zeta(-n) = (-1)^{n-1} \frac{B_n}{n}.$$



## Euler's discoveries from a modern perspective

By Euler's formula, the values of the Riemann zeta functions at even integers are of the form

$$\zeta(2n) = (-1)^{k+1} \frac{(2\pi)^{2k}}{2(2k)!} B_{2k}.$$

Using the functional equation, we have

$$\zeta(1-n) = \zeta(-n) = (-1)^{n-1} \frac{B_n}{n}.$$

This formula implies that  $\zeta(1-n)$  is a rational number for all  $n$ .

## Euler's discoveries from a modern perspective

By Euler's formula, the values of the Riemann zeta functions at even integers are of the form

$$\zeta(2n) = (-1)^{k+1} \frac{(2\pi)^{2k}}{2(2k)!} B_{2k}.$$

Using the functional equation, we have

$$\zeta(1-n) = \zeta(-n) = (-1)^{n-1} \frac{B_n}{n}.$$

This formula implies that  $\zeta(1-n)$  is a rational number for all  $n$ . The values  $\zeta(1-n)$  are often called **zeta values**.

## Euler's discoveries from a modern perspective

By Euler's formula, the values of the Riemann zeta functions at even integers are of the form

$$\zeta(2n) = (-1)^{k+1} \frac{(2\pi)^{2k}}{2(2k)!} B_{2k}.$$

Using the functional equation, we have

$$\zeta(1-n) = \zeta(-n) = (-1)^{n-1} \frac{B_n}{n}.$$

This formula implies that  $\zeta(1-n)$  is a rational number for all  $n$ . The values  $\zeta(1-n)$  are often called **zeta values**.

### Question

*What is the arithmetic significance of the zeta values  $\zeta(1-n)$ ?*

## Class number formula

Let  $K$  be a number field and  $\mathcal{O}_K$  be its ring of algebraic integers.

Let

## Class number formula

Let  $K$  be a number field and  $\mathcal{O}_K$  be its ring of algebraic integers.

Let

- $r_1$  (respectively  $r_2$ ) be the number of real (respectively complex) places of  $K$ .

## Class number formula

Let  $K$  be a number field and  $\mathcal{O}_K$  be its ring of algebraic integers.

Let

- $r_1$  (respectively  $r_2$ ) be the number of real (respectively complex) places of  $K$ .
- $h_K$ : class number of  $K$ .

## Class number formula

Let  $K$  be a number field and  $\mathcal{O}_K$  be its ring of algebraic integers.

Let

- $r_1$  (respectively  $r_2$ ) be the number of real (respectively complex) places of  $K$ .
- $h_K$ : class number of  $K$ .
- $w_K$ : number of roots of unity in  $K$ .

## Class number formula

Let  $K$  be a number field and  $\mathcal{O}_K$  be its ring of algebraic integers.

Let

- $r_1$  (respectively  $r_2$ ) be the number of real (respectively complex) places of  $K$ .
- $h_K$ : class number of  $K$ .
- $w_K$ : number of roots of unity in  $K$ .
- $R_K$ : the regulator of  $K$ .



## Class number formula

Let  $K$  be a number field and  $\mathcal{O}_K$  be its ring of algebraic integers.

Let

- $r_1$  (respectively  $r_2$ ) be the number of real (respectively complex) places of  $K$ .
- $h_K$ : class number of  $K$ .
- $w_K$ : number of roots of unity in  $K$ .
- $R_K$ : the regulator of  $K$ .
- $D_K$ : the discriminant of  $K$ .

## Class number formula

Let  $K$  be a number field and  $\mathcal{O}_K$  be its ring of algebraic integers.

Let

- $r_1$  (respectively  $r_2$ ) be the number of real (respectively complex) places of  $K$ .
- $h_K$ : class number of  $K$ .
- $w_K$ : number of roots of unity in  $K$ .
- $R_K$ : the regulator of  $K$ .
- $D_K$ : the discriminant of  $K$ .

These numbers contain arithmetic information about  $K$ .

## Class number formula

A quite natural generalization of the Riemann zeta function is the Dedekind zeta function for  $K$  defined by

$$\zeta_K(s) = \sum_{\mathfrak{a} \subset \mathcal{O}_K} \frac{1}{N(\mathfrak{a})^s}.$$

It has a meromorphic continuation to  $\mathbb{C}$  with a unique simple pole at  $s = 1$ .

## Class number formula

A quite natural generalization of the Riemann zeta function is the Dedekind zeta function for  $K$  defined by

$$\zeta_K(s) = \sum_{\mathfrak{a} \subset \mathcal{O}_K} \frac{1}{N(\mathfrak{a})^s}.$$

It has a meromorphic continuation to  $\mathbb{C}$  with a unique simple pole at  $s = 1$ .

### Theorem (Class number formula)

$$\lim_{s \rightarrow 1} (s - 1) \zeta_K(s) = \frac{2^{r_1} (2\pi)^{r_2} R_K h_K}{w_K \sqrt{|D_K|}}.$$

## Class number formula

A quite natural generalization of the Riemann zeta function is the Dedekind zeta function for  $K$  defined by

$$\zeta_K(s) = \sum_{\mathfrak{a} \subset \mathcal{O}_K} \frac{1}{N(\mathfrak{a})^s}.$$

It has a meromorphic continuation to  $\mathbb{C}$  with a unique simple pole at  $s = 1$ .

### Theorem (Class number formula)

$$\lim_{s \rightarrow 1} (s - 1) \zeta_K(s) = \frac{2^{r_1} (2\pi)^{r_2} R_K h_K}{w_K \sqrt{|D_K|}}.$$

This formula provides a connection between the algebraic world and the analytic world.

## Birch and Swinnerton-Dyer conjecture

- From a modern point of view, the Riemann zeta function and its generalization, the Dedekind zeta functions, concern the arithmetics/geometry of zero dimensional objects, namely  $\text{Spec}(K)$ .

## Birch and Swinnerton-Dyer conjecture

- From a modern point of view, the Riemann zeta function and its generalization, the Dedekind zeta functions, concern the arithmetics/geometry of zero dimensional objects, namely  $\text{Spec}(K)$ .
- It is quite natural to generalize this study to higher dimensional objects.

## Birch and Swinnerton-Dyer conjecture

- From a modern point of view, the Riemann zeta function and its generalization, the Dedekind zeta functions, concern the arithmetics/geometry of zero dimensional objects, namely  $\text{Spec}(K)$ .
- It is quite natural to generalize this study to higher dimensional objects.
- An elliptic curve is a plane algebraic curve given by an equation of the form

$$y^2 = x^3 + ax + b.$$



## Birch and Swinnerton-Dyer conjecture

- From a modern point of view, the Riemann zeta function and its generalization, the Dedekind zeta functions, concern the arithmetics/geometry of zero dimensional objects, namely  $\text{Spec}(K)$ .
- It is quite natural to generalize this study to higher dimensional objects.
- An elliptic curve is a plane algebraic curve given by an equation of the form

$$y^2 = x^3 + ax + b.$$

The  $L$ -function of  $E$  is defined by the Euler product

$$L(s, E) = \prod_p L_p(s, E)^{-1}.$$

# The Birch and Swinnerton-Dyer conjecture

Let  $E(\mathbb{Q})$  be the set of all rational solutions of  $E$ , i.e.

$$E(\mathbb{Q}) = \{(x, y) \in \mathbb{Q}^2 \mid y^2 = x^3 + ax + b\}.$$

# The Birch and Swinnerton-Dyer conjecture

Let  $E(\mathbb{Q})$  be the set of all rational solutions of  $E$ , i.e.

$$E(\mathbb{Q}) = \{(x, y) \in \mathbb{Q}^2 \mid y^2 = x^3 + ax + b\}.$$

The Mordell-Weil theorem says that the set  $E(\mathbb{Q})$  is a finite generated abelian group of finite rank  $r_{al}$ .

# The Birch and Swinnerton-Dyer conjecture

Let  $E(\mathbb{Q})$  be the set of all rational solutions of  $E$ , i.e.

$$E(\mathbb{Q}) = \{(x, y) \in \mathbb{Q}^2 \mid y^2 = x^3 + ax + b\}.$$

The Mordell-Weil theorem says that the set  $E(\mathbb{Q})$  is a finite generated abelian group of finite rank  $r_{a/b}$ . In the early 1960s Birch and Swinnerton-Dyer did lots of numerical computations with elliptic curves. These computations lead them to a remarkable conjecture.

# The Birch and Swinnerton-Dyer conjecture

Let  $E(\mathbb{Q})$  be the set of all rational solutions of  $E$ , i.e.

$$E(\mathbb{Q}) = \{(x, y) \in \mathbb{Q}^2 \mid y^2 = x^3 + ax + b\}.$$

The Mordell-Weil theorem says that the set  $E(\mathbb{Q})$  is a finite generated abelian group of finite rank  $r_{al}$ . In the early 1960s Birch and Swinnerton-Dyer did lots of numerical computations with elliptic curves. These computations lead them to a remarkable conjecture.

## Conjecture (BSD conjecture, weak form)

$$r_{al} = r_{an},$$

where  $r_{an}$  is the order of vanishing of the  $L$ -function  $L(E, s)$  at  $s = 1$ .

# Generalizations

- In the 70-80s, mathematicians realized that these two examples should be special cases of a more general phenomenon.

# Generalizations

- In the 70-80s, mathematicians realized that these two examples should be special cases of a more general phenomenon.
- Beilinson and Deligne made a breakthrough by putting this question in the framework of mixed motives and motivic cohomology. However, their work only predicts  $L$ -values up to an undetermined rational factor. Beilinson's approach is inspired by the work of Bloch on  $K_2$  of CM elliptic curves.

# Generalizations

- In the 70-80s, mathematicians realized that these two examples should be special cases of a more general phenomenon.
- Beilinson and Deligne made a breakthrough by putting this question in the framework of mixed motives and motivic cohomology. However, their work only predicts  $L$ -values up to an undetermined rational factor. Beilinson's approach is inspired by the work of Bloch on  $K_2$  of CM elliptic curves.
- Bloch and Kato formulated a more precise conjecture using tools from  $p$ -adic Hodge theory discovered by Fontaine.



## What is a (mixed) motive?

- There is still no universally agreed definition of mixed motives even up to this day. Progress has been made by many mathematicians, notably Grothendieck, Jannsen, Bloch, Beilinson, Deligne and many others.

## What is a (mixed) motive?

- There is still no universally agreed definition of mixed motives even up to this day. Progress has been made by many mathematicians, notably Grothendieck, Jannsen, Bloch, Beilinson, Deligne and many others.
- A folklore definition:  $H^n(X)(r)$  where  $X$  is a smooth variety over  $\mathbb{Q}$ . It has different realizations depending on different cohomology theories (de Rham, Betti, etale, crystalline.) These realizations are connected by comparison isomorphisms. For example, if  $X$  is smooth and projective, there is an isomorphism between Betti (singular) cohomology and etale cohomology

$$H_{et}^n(X_{\bar{\mathbb{Q}}}, \mathbb{Z}_\ell) \cong H_B^n(X(\mathbb{C}), \mathbb{Z}_\ell).$$

## A working definition

- Jannsen's definition: a mixed motive with  $\mathbb{Q}$ -coefficients is a collection of realizations with weight and Hodge filtrations. These realizations are related by comparison isomorphisms that are compatible with both of these filtrations.

## A working definition

- Jannsen's definition: a mixed motive with  $\mathbb{Q}$ -coefficients is a collection of realizations with weight and Hodge filtrations. These realizations are related by comparison isomorphisms that are compatible with both of these filtrations.
- A mixed motive with  $\mathbb{Z}$ -coefficients is a pair  $(M, \Theta)$  where
  1.  $M$  is a mixed motive with  $\mathbb{Q}$  coefficients.
  2.  $\Theta$  is a free  $\mathbb{Z}$ -module of finite rank equipped with a linear action of  $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$  on  $\hat{\Theta} = \hat{\mathbb{Z}} \otimes \Theta$ .
  3. There is an isomorphism  $\mathbb{Q} \otimes \Theta \cong M_B$  where  $M_B$  is the Betti realization of  $M$ .

## Period rings and the local Bloch-Kato Selmer groups

Let  $V$  be a  $\mathbb{Q}_\ell$ -vector space equipped with an action of  $\text{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$  and  $T \subset V$  a Galois stable lattice. When  $\ell = p$ , we define the following subgroups of  $H^1(\mathbb{Q}_p, V)$  :

# Period rings and the local Bloch-Kato Selmer groups

Let  $V$  be a  $\mathbb{Q}_\ell$ -vector space equipped with an action of  $\text{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$  and  $T \subset V$  a Galois stable lattice. When  $\ell = p$ , we define the following subgroups of  $H^1(\mathbb{Q}_p, V)$  :

- $H_e^1(\mathbb{Q}_p, V) = \ker(H^1(\mathbb{Q}_p, V) \rightarrow H^1(\mathbb{Q}_p, V \otimes B_{\text{crys}}^{\varphi=1}))$ .
- $H_f^1(\mathbb{Q}_p, V) = \ker(H^1(\mathbb{Q}_p, V) \rightarrow H^1(\mathbb{Q}_p, V \otimes B_{\text{crys}}))$ .
- $H_g^1(\mathbb{Q}_p, V) = \ker(H^1(\mathbb{Q}_p, V) \rightarrow H^1(\mathbb{Q}_p, V \otimes B_{\text{dR}}))$ .

These are called the exponential, finite, and geometric parts of  $H^1(\mathbb{Q}_p, V)$ .

## Local Bloch-Kato Selmer groups

Let  $* \in \{e, f, g\}$ . We define  $H_*^1(\mathbb{Q}_p, T)$  to be the pre-image of  $H_*^1(\mathbb{Q}_p, V)$  under the canonical map

$$H^1(\mathbb{Q}_p, T) \rightarrow H^1(\mathbb{Q}_p, V).$$

## Local Bloch-Kato Selmer groups

Let  $* \in \{e, f, g\}$ . We define  $H_*^1(\mathbb{Q}_p, T)$  to be the pre-image of  $H_*^1(\mathbb{Q}_p, V)$  under the canonical map

$$H^1(\mathbb{Q}_p, T) \rightarrow H^1(\mathbb{Q}_p, V).$$

If  $M$  is a  $\widehat{\mathbb{Z}}$ -module with a continuous  $\text{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$  action, we define

$$H_*^1(\mathbb{Q}_p, M) = \prod_{\ell} H_*^1(\mathbb{Q}_p, M_{\ell}).$$



# Tamagawa measure

We define the following Selmer conditions on local Galois cohomology.

$$A(\mathbb{Q}_p) = \begin{cases} H_f^1(\mathbb{Q}_p, \hat{\Theta}) & \text{if } p < \infty \\ (D_\infty \otimes_{\mathbb{R}} \mathbb{C}) / (\text{Fil}^0 D_\infty \otimes_{\mathbb{R}} \mathbb{C}) + \Theta)^+ & \text{if } p = \infty. \end{cases}$$

# Tamagawa measure

We define the following Selmer conditions on local Galois cohomology.

$$A(\mathbb{Q}_p) = \begin{cases} H_f^1(\mathbb{Q}_p, \hat{\Theta}) & \text{if } p < \infty \\ (D_\infty \otimes_{\mathbb{R}} \mathbb{C}) / (\text{Fil}^0 D_\infty \otimes_{\mathbb{R}} \mathbb{C}) + \Theta)^+ & \text{if } p = \infty. \end{cases}$$

The exponential map

$$\exp : D_p / D_p^0 \rightarrow A(\mathbb{Q}_p),$$

is an isomorphism if  $\omega(M) \leq -1$  where  $D_p$  is the  $p$ -adic de Rham realization of  $M$ .

# Tamagawa measure

By fixing a basis for  $\det(D)$ , we can define the Tamagawa measure on  $\mu(A(\mathbb{Q}_p))$ .

# Tamagawa measure

By fixing a basis for  $\det(D)$ , we can define the Tamagawa measure on  $\mu(A(\mathbb{Q}_p))$ . Furthermore, under some technical conditions, we have

$$\mu(A(\mathbb{Q}_p)) = L_p(V, 0)^{-1},$$

where  $L_p(V, p^{-s})$  is the local Euler factor at  $p$ .

# Tamagawa measure

By fixing a basis for  $\det(D)$ , we can define the Tamagawa measure on  $\mu(A(\mathbb{Q}_p))$ . Furthermore, under some technical conditions, we have

$$\mu(A(\mathbb{Q}_p)) = L_p(V, 0)^{-1},$$

where  $L_p(V, p^{-s})$  is the local Euler factor at  $p$ .

The induced measure  $\mu$  on the adelic space  $\prod'_p A(\mathbb{Q}_p)$  is called the Tamagawa measure of  $M$ .

The Tamagawa number of  $M$  is defined as

$$\mathrm{Tam}(M) = \mu \left( \prod'_{p \leq \infty} A(\mathbb{Q}_p) / A(\mathbb{Q}) \right),$$

where  $A(\mathbb{Q})$  can be thought as  $\mathrm{Ext}^1(\mathbb{Z}, (M, \Theta))$  in the category of motives with  $\mathbb{Z}$ -coefficients satisfying certain Selmer conditions.

The Tamagawa number of  $M$  is defined as

$$\mathrm{Tam}(M) = \mu \left( \prod'_{p \leq \infty} A(\mathbb{Q}_p) / A(\mathbb{Q}) \right),$$

where  $A(\mathbb{Q})$  can be thought as  $\mathrm{Ext}^1(\mathbb{Z}, (M, \Theta))$  in the category of motives with  $\mathbb{Z}$ -coefficients satisfying certain Selmer conditions.

If  $M = H^n(X)(r)$ , then we should have

$A(\mathbb{Q}) \subset A(\mathbb{Q}) \otimes \mathbb{Q} = \mathrm{gr}^r K_{2n-r-1}(\mathfrak{X}) \otimes \mathbb{Q}$  where  $\mathfrak{X}/\mathbb{Z}$  is a regular flat model of  $X$ .

# Tamagawa number conjecture, I

## Conjecture (Tamagawa number conjecture)

$$\mathrm{Tam}(M) = \frac{\#H^0(\mathbb{Q}, M^* \otimes \mathbb{Q}/\mathbb{Z}(1))}{\#\mathrm{III}(M)},$$

where  $\mathrm{III}(M)$  is the Tate-Shafarevich group associated with  $M$ .  
It is conjectured to be a finite group.



## Some known results

- $M = \mathbb{Z}(r)$ . The class number formula is a special case of this when  $r = 1$ . (Bloch-Kato)
- $M = H^1(E)(2)$  where  $E$  is an elliptic curve with CM (Bloch-Kato)
- $M = H^1(E)(1)$  in some special cases by work of Coates-Wiles, Kolyvagin, Rubin, Kato, and many others. This case is equivalent to the BSD conjecture.
- $M$  is the adjoint motive of a modular form (Diamond, Flach, Guo, Loeffler-Lei-Zerbes and others).
- Some progress for motives associated with automorphic forms on  $GSp_4$ ,  $GL_2 \times GL_2$  (Loeffler-Zerbes and others.)
- Most of these works construct special elements in  $A(\mathbb{Q})$ .
- Iwasawa theory plays an important role.

## Tamagawa number conjecture, II

- In their work, Bloch-Kato mentioned that the Tamagawa number conjecture could be formulated using  $B(\mathbb{Q})$  and  $B(\mathbb{Q}_p)$  where  $B(\mathbb{Q}_p)$  is the inverse image in  $H_g^1(\mathbb{Q}_p, \hat{\Theta})$  of

$$\text{Im} \left( \Psi \rightarrow H_g^1(\mathbb{Q}_p, \Theta \otimes \mathbf{A}_{\mathbb{Q}}^f) / H_f^1(\mathbb{Q}_p, \Theta \otimes \mathbf{A}_{\mathbb{Q}}^f) \right),$$

and  $B(\mathbb{R}) = A(\mathbb{R})$ .

## Tamagawa number conjecture, II

- In their work, Bloch-Kato mentioned that the Tamagawa number conjecture could be formulated using  $B(\mathbb{Q})$  and  $B(\mathbb{Q}_p)$  where  $B(\mathbb{Q}_p)$  is the inverse image in  $H_g^1(\mathbb{Q}_p, \hat{\Theta})$  of

$$\text{Im} \left( \psi \rightarrow H_g^1(\mathbb{Q}_p, \Theta \otimes \mathbf{A}_{\mathbb{Q}}^f) / H_f^1(\mathbb{Q}_p, \Theta \otimes \mathbf{A}_{\mathbb{Q}}^f) \right),$$

and  $B(\mathbb{R}) = A(\mathbb{R})$ .

- We define  $B(\mathbb{Q})$  similarly. As before,  $B(\mathbb{Q})$  should be considered as the extension group  $\text{Ext}^1(\mathbb{Z}, (M, \Theta))$  in the category of motives with  $\mathbb{Z}$ -coefficients (with no Selmer conditions).

## Tamagawa number conjecture, II

- In their work, Bloch-Kato mentioned that the Tamagawa number conjecture could be formulated using  $B(\mathbb{Q})$  and  $B(\mathbb{Q}_p)$  where  $B(\mathbb{Q}_p)$  is the inverse image in  $H_g^1(\mathbb{Q}_p, \hat{\Theta})$  of

$$\text{Im} \left( \psi \rightarrow H_g^1(\mathbb{Q}_p, \Theta \otimes \mathbf{A}_{\mathbb{Q}}^f) / H_f^1(\mathbb{Q}_p, \Theta \otimes \mathbf{A}_{\mathbb{Q}}^f) \right),$$

and  $B(\mathbb{R}) = A(\mathbb{R})$ .

- We define  $B(\mathbb{Q})$  similarly. As before,  $B(\mathbb{Q})$  should be considered as the extension group  $\text{Ext}^1(\mathbb{Z}, (M, \Theta))$  in the category of motives with  $\mathbb{Z}$ -coefficients (with no Selmer conditions).
- When  $M = H^n(X)(r)$ , we should take

$$\psi = \text{gr}^r K_{2r-m-1}(X) \otimes \mathbb{Q}.$$

## Height functions

- A height function is a function that quantifies the complexity of a mathematical object.

## Height functions

- A height function is a function that quantifies the complexity of a mathematical object.
- Let us take an example of a height function. If  $r = \frac{p}{q}$  is a rational number written in the reduced form then the multiplicative height of  $r$  is defined to be

$$H\left(\frac{p}{q}\right) = \max\{|p|, |q|\}.$$

# Height functions

- A height function is a function that quantifies the complexity of a mathematical object.
- Let us take an example of a height function. If  $r = \frac{p}{q}$  is a rational number written in the reduced form then the multiplicative height of  $r$  is defined to be

$$H\left(\frac{p}{q}\right) = \max\{|p|, |q|\}.$$

- Height functions play important roles in many parts of mathematics. For example, G. Faltings defined height functions for abelian varieties and used them to give a proof of the Mordell conjecture.

# Height functions

- A height function is a function that quantifies the complexity of a mathematical object.
- Let us take an example of a height function. If  $r = \frac{p}{q}$  is a rational number written in the reduced form then the multiplicative height of  $r$  is defined to be

$$H\left(\frac{p}{q}\right) = \max\{|p|, |q|\}.$$

- Height functions play important roles in many parts of mathematics. For example, G. Faltings defined height functions for abelian varieties and used them to give a proof of the Mordell conjecture.
- K. Kato recently defined height functions for motives. He dreams that these definitions will solve important problems in number theory.



## Local height functions

Recall that we have the following inclusions

$$A(\mathbb{Q}_p) := H_f^1(\mathbb{Q}_p, \hat{\Theta}) \subset B(\mathbb{Q}_p) \subset H_g^1(\mathbb{Q}_p, \hat{\Theta}).$$

Furthermore, under some technical conditions then

$$H_g^1(\mathbb{Q}_p, M_\ell) / H_f^1(\mathbb{Q}_p, M_\ell) = \begin{cases} ((\mathrm{gr}_{-2}^{\mathcal{W}} M_\ell(-1))_{\mathrm{prim}})^{\varphi=1} & \text{if } \ell \neq p \\ ((\mathrm{gr}_{-2}^{\mathcal{W}} D_{\mathrm{st}}(M_p))_{\mathrm{prim}}(-1))^{\varphi=1} & \text{if } \ell = p. \end{cases}$$

We use this identification and the polarization of  $M$  to define the height function

$$H_{\diamond, v, d} : B(\mathbb{Q}_p) \rightarrow \mathbb{R}_{\geq 0}.$$

By definition,  $H_{\diamond, v, d}$  is trivial on  $A(\mathbb{Q}_p)$ . Furthermore, it is expected that  $H_{\diamond, v, d}$  is induced by a positive-definite quadratic form on  $B(\mathbb{Q}_p)/A(\mathbb{Q}_p)$ .

## Local height functions

Under some conjectures as in Bloch-Kato's article, we have  $B(\mathbb{Q}_p)/A(\mathbb{Q}_p) \cong \mathbb{Z}^{r_p}$  as topological groups. By choosing a basis, we have

$$B(\mathbb{Q}_p) = A(\mathbb{Q}_p) \times \mathbb{Z}^{s_p}.$$

The local Tamagawa measure  $\mu$  on  $B(\mathbb{Q}_p)$  is defined to be the product measure of the Tamagawa measure on  $A(\mathbb{Q}_p)$  and the counting measure on  $\mathbb{Z}^{s_p}$ .

The following statement is straightforward.

### Proposition

*For each positive real number  $B$*

$$\mu(a \in B(\mathbb{Q}_p) | H_{\diamond, p, d}(a) \leq B) < \infty.$$

# Adelic and global height functions

We define

$$H_{\diamond, d} : \prod'_{p \leq \infty} B(\mathbb{Q}_p) \rightarrow \mathbb{R}_{\geq 0}$$

as the product of local height functions.

Similarly, we define the height function on  $B(\mathbb{Q})$  using the canonical embedding

$$B(\mathbb{Q}) \rightarrow \prod_{p \leq \infty} B(\mathbb{Q}_p).$$

By definition  $H_{\diamond, d}$  is trivial on  $A(\mathbb{Q})$ .

We also define the Tamagawa measure on  $\prod'_{p \leq \infty} B(\mathbb{Q}_p)$  as the product of the local Tamagawa measure on  $B(\mathbb{Q}_p)$ .

# Adelic and global height functions

A simple observation.

## Proposition

*For each positive real number  $B$ , the set*

$$\{a \in \prod'_{p \leq \infty} B(\mathbb{Q}_p) \mid H_{\diamond, d}(a) \leq B\}.$$

*is compact. In particular,*

$$\mu(a \in \prod'_{p \leq \infty} B(\mathbb{Q}_p) \mid H_{\diamond, d}(a) \leq B) < \infty,$$

*and*

$$\#\{a \in B(\mathbb{Q}) \mid H_{\diamond, d}(a) \leq B\} < \infty.$$

We have the the following theorem.

**Theorem (Nguyen)**

*Let  $M$  be a pure motive with  $\mathbb{Z}$ -coefficients of weight  $-d$  such that  $d \geq 3$ . We assume further that  $M$  has semistable reduction at all places. Then*

$$\lim_{B \rightarrow \infty} \frac{\#\{x \in B(\mathbb{Q}) \mid H_{\diamond, d}(x) \leq B\}}{\mu\left(x \in \prod'_{p \leq \infty} B(\mathbb{Q}_p) \mid H_{\diamond, d}(x) \leq B\right)} = \frac{1}{\text{Tam}(M)}.$$

**Main idea of the proof:** For a symmetric, convex region  $A$  in  $\mathbb{R}^n$ , the number of integer points in  $A$  is asymptotic to the area of  $A$ .

## Recent and future work

- I hope my presentation convinces the audience that zeta functions are beautiful and mysterious.

## Recent and future work

- I hope my presentation convinces the audience that zeta functions are beautiful and mysterious.
- In our recent joint work with Jan Minac and Nguyen Duy Tan, we studied some generalization of the Riemann zeta functions. We discovered some quite interesting formulas for zeta values.

## Recent and future work

- I hope my presentation convinces the audience that zeta functions are beautiful and mysterious.
- In our recent joint work with Jan Minac and Nguyen Duy Tan, we studied some generalization of the Riemann zeta functions. We discovered some quite interesting formulas for zeta values.
- In an ongoing project with Lyle Muller et al, we are investigating the Ihara zeta functions for graphs, random graphs and their applications to neuroscience.



Thank you!