

Prevention Quality Indicator 08 (PQI 08) Heart Failure Admission Rate

October 2016 Area-Level Indicator Type of Score: Rate

Prepared by:

Agency for Healthcare Research and Quality U.S. Department of Health and Human Services www.qualityindicators.ahrq.gov

DESCRIPTION

Admissions with a principal diagnosis of heart failure per 100,000 population, ages 18 years and older. Excludes cardiac procedure admissions, obstetric admissions, and transfers from other institutions.

[NOTE: The software provides the rate per population. However, common practice reports the measure as per 100,000 population. The user must multiply the rate obtained from the software by 100,000 to report admissions per 100,000 population.]

October 2016 1 of 4

AHRQ QI[™] ICD-9-CM Specification Version 6.0 PQI 08 Heart Failure Admission Rate www.qualityindicators.ahrq.gov

NUMERATOR

Discharges, for patients ages 18 years and older, with a principal ICD-9-CM diagnosis code for heart failure.

[NOTE: By definition, discharges with a principal diagnosis of heart failure are precluded from an assignment of MDC 14 by grouper software. Thus, obstetric discharges should not be considered in the PQI rate, though the AHRQ QITM software does not explicitly exclude obstetric cases.]

Heart failure diagnosis codes (MRTCHFD):

| 39891 | RHEUMATIC HEART FAILURE | 42821 | AC SYSTOLIC HRT FAILURE |
|-------|--------------------------|-------|--------------------------|
| 40201 | MAL HYPERT HRT DIS W CHF | 42822 | CHR SYSTOLIC HRT FAILURE |
| 40211 | BENIGN HYP HRT DIS W CHF | 42823 | AC ON CHR SYST HRT FAIL |
| 40291 | HYPERTEN HEART DIS W CHF | 42830 | DIASTOLC HRT FAILURE NOS |
| 40401 | MAL HYPER HRT/REN W CHF | 42831 | AC DIASTOLIC HRT FAILURE |
| 40403 | MAL HYP HRT/REN W CHF&RF | 42832 | CHR DIASTOLIC HRT FAIL |
| 40411 | BEN HYPER HRT/REN W CHF | 42833 | AC ON CHR DIAST HRT FAIL |
| 40413 | BEN HYP HRT/REN W CHF&RF | 42840 | SYST/DIAST HRT FAIL NOS |
| 40491 | HYPER HRT/REN NOS W CHF | 42841 | AC SYST/DIASTOL HRT FAIL |
| 40493 | HYP HT/REN NOS W CHF&RF | 42842 | CHR SYST/DIASTL HRT FAIL |
| 4280 | CONGESTIVE HEART FAILURE | 42843 | AC/CHR SYST/DIA HRT FAIL |
| 4281 | LEFT HEART FAILURE | 4289 | HEART FAILURE NOS |
| 42820 | SYSTOLIC HRT FAILURE NOS | | |
| | | | |

October 2016 2 of 4

AHRQ QI™ ICD-9-CM Specification Version 6.0 PQI 08 Heart Failure Admission Rate www.qualityindicators.ahrq.gov

NUMERATOR EXCLUSIONS

Exclude cases:

- with any-listed ICD-9-CM procedure codes for cardiac procedure
- transfer from a hospital (different facility)
- transfer from a Skilled Nursing Facility (SNF) or Intermediate Care Facility (ICF)
- transfer from another health care facility
- with missing gender (SEX=missing), age (AGE=missing), quarter (DQTR=missing), year (YEAR=missing), principal diagnosis (DX1=missing), or county (PSTCO=missing)

<u>Appendix A – Admission Codes for Transfers</u> <u>Appendix B – Cardiac Procedure Codes</u>

October 2016 3 of 4

AHRQ QI[™] ICD-9-CM Specification Version 6.0 PQI 08 Heart Failure Admission Rate www.qualityindicators.ahrq.gov

DENOMINATOR

Population ages 18 years and older in metropolitan area† or county. Discharges in the numerator are assigned to the denominator based on the metropolitan area or county of the patient residence, not the metropolitan area or county of the hospital where the discharge occurred.

† The term "metropolitan area" (MA) was adopted by the U.S. Census in 1990 and referred collectively to metropolitan statistical areas (MSAs), consolidated metropolitan statistical areas (CMSAs) and primary metropolitan statistical areas (PMSAs). In addition, "area" could refer to either 1) FIPS county, 2) modified FIPS county, 3) 1999 OMB Metropolitan Statistical Area or 4) 2003 OMB Metropolitan Statistical Area. Micropolitan Statistical Areas are not used in the QI software.