

A choice-free Lean formalization of a profile-based dominated convergence theorem in presented Bishop–Cheng measure theory

Toshihisa Urahigashi

Draft of 2026-08-31
Artifact version 0.6.0

Abstract

Bishop and Cheng derive the dominated convergence theorem of their integral-first constructive measure theory through profiles, that is, through monotone functionals on families of continuous ramp functions. All but countably many levels of such a profile are *smooth* in the sense the memoir gives that word, the constructive stand-in for a point of continuity of the distribution function. The step “choose a level outside a countable exceptional set” is exactly where a choice-free version of the theory is known to be obstructed: Spitters observed that the profile theorem implies the uncountability of the reals, which fails in an intuitionistic sheaf model. This paper reports a Lean 4 formalization of the profile route to Bishop–Cheng dominated convergence: convergence in measure, together with domination on full sets that may depend on the index, implies convergence of the integrals. The theorem is proved over *presented* reals—regular sequences of rationals with Bishop equality—and against an explicitly quantified integration-space interface, rather than for an abstract constructive real-number object. The method is to enumerate the exceptional levels as an explicit sequence and to step around them by Bishop’s nested-interval construction, whose case split is decided from rational approximation data and returned as data. For the named public declarations, the kernel reports an axiom footprint confined to `[propext, Quot.sound]`; a static scan of the source closure and a set of checksums accompany it. The artifact also packages a Rocq `Print Assumptions` audit of Séméria’s CoRN declarations, reproduced by the author from a pinned CoRN snapshot. No priority is claimed over that formalization. Relative to the development’s encodings of the ambient notions, the result covers every integration space in Bishop and Cheng’s sense: each field of the working interface is derived, hypothesis-free, from a clause-by-clause transcription of their Definition 1.1 at the memoir’s own gauge. In the other direction, a finitely supported model, unconditional and normalized at two points, exhibits an instance whose measure is not concentrated at a single point.

Keywords: constructive measure theory; dominated convergence; Lean 4; proof assistants; presented reals; choice principles.

How to read this paper. Part I (Sections 1–6) is mathematical. It fixes the setting, states the definitions, states the theorem, and gives the proof, and it can be read without any knowledge of Lean or of proof assistants: no formal identifier occurs in a definition or in the statement of a theorem. Throughout the paper, “the source” refers to Bishop and Cheng’s memoir *Constructive Measure Theory* [2]; Bishop’s *Foundations of Constructive Analysis* [1] is always cited by name. Part II (Sections 7–13) concerns the formalization: the shape of the public interface, the relation of the result to the known no-go theorems, the comparison with earlier formalizations, the audit, the size of the development and how much of it the claims reach, and the limitations and reproduction procedure. Appendix A tabulates the correspondence between the numbering of this paper and that

of the source, and between the mathematical objects of Part I and the Lean declarations of Part II. Appendix B states, as ordinary mathematics and without Lean vocabulary, the constructions that the formalization had to supply beyond what the source provides, together with a classification of which parts are routine, which are known, and which were the actual difficulties. Part I suffices for the mathematical content; Appendix B is addressed to a reader who wants to know what the formalization concretely consisted of.

Part I

The mathematics

1 Introduction

Dominated convergence is a basic passage from convergence of functions to convergence of their integrals. In the integral-first constructive measure theory of Bishop and Cheng, the proof is not simply a translation of the classical almost-everywhere argument. It proceeds through profiles of integrable functions, integrability of level sets away from an explicit exceptional sequence, and a uniform estimate on the complement of a suitable integrable set.

The reason is worth stating at once, because the whole paper turns on it. Classically, one proves that a monotone function $\mathbb{R} \rightarrow \mathbb{R}$ has at most countably many discontinuities, and one then picks a level t at which the distribution function $t \mapsto \mu(\{|h| > t\})$ is continuous. Both halves of that argument are problematic constructively. Bishop and Cheng replace the first half by the theory of profiles, which returns not an abstract countable set of bad levels but an *explicit sequence* of them. The second half—picking a level different from every member of an explicitly given sequence—then becomes a genuine construction rather than a triviality, and it is precisely the step at which choice-free constructive measure theory is known to encounter an obstruction: in an unrestricted constructive setting, the relevant profile theorem entails an uncountability principle that fails in an intuitionistic sheaf model [22].

This paper studies what can nevertheless be formalized when the real numbers are supplied with presentations. A presented real is not merely an element of an abstract ordered field. It carries rational approximation data and a modulus, and it returns as data the strict-order cotransitivity required by Bishop’s nested-interval construction. The formalization uses that representation to build a point apart from every member of a given sequence, applies the construction to the exceptional levels of a profile, and then follows the Bishop–Cheng route to dominated convergence.

Here and below, *choice-free* has a deliberately audit-oriented meaning. It means that the named public Lean declarations, including their transitive proof-term dependencies, do not use `Classical.choice`, `Classical.choose`, `sorryAx`, or a comparable selector axiom. It does not mean that the theorems have no hypotheses, that every imported module is constructive, or that the profile theorem has been proved for every possible constructive real-number object. The reason for fixing the meaning this narrowly is stated in the literature: Berger and Seisenberger observe that “relativized choice—though trivially provable in constructive type theory—is in general not realizable” [17, §29.5]. Derivability inside a type theory therefore does not by itself guarantee realizability or the intended computational content, and a claim of this kind has to be attached to a named dependency closure and its axiom output (Section 8, Section 10).

The contributions of the artifact are as follows.

- (1) It gives an explicit Lean 4 theorem surface for a profile-based Bishop–Cheng dominated-convergence route over regular-sequence presented reals and an explicitly quantified

integration-space structure.

- (2) It gives a Lean implementation of the countable-avoidance step via Bishop’s nested-interval construction: convergence data for the endpoints, an apart limiting point, and the resulting smooth-level data are constructed inside the audited dependency closure.
- (3) It supplies a model of the interface in which the measure is not concentrated at a single point: finitely many points with weights, discharging every clause of Definition 1.1, with an unconditional normalized instance at two points. For a single point the continuity clause is the identity; for several, the point at which the series stays below the function has to be found, and it is found from the positivity witness of the strict inequality without any appeal to choice.
- (4) It separates a proof-relevant Type-valued route, automatic wrappers that synthesize smooth-level data, and a separately proved Prop-facing theorem. The Type-valued route includes both global pointwise-majorant wrappers and a full-set variant in which the carrier and fullness proof for every index are explicit inputs. This makes the computational content visible without requiring every caller to supply the low-level data explicitly.
- (5) It provides reproducible Lean and Rocq audits, a pinned comparison with Séméria’s earlier CoRN development, and explicit qualifications concerning predicativity, the integration-space interface, concrete models, and priority. The paper separately reports the size of the development (Section 11).

Each principal claim is stated together with an explicit qualification. The following table pairs them; every row is developed in the section cited.

Claim	Direct evidence	Qualification maintained
A profile-route Bishop–Cheng dominated convergence theorem over presented reals (Theorem 4.1)	the Prop-facing public declaration and the separately proved Type-valued route (Section 7)	no priority over the CoRN formalization (Section 9)
The working interface is derived hypothesis-free—all fourteen fields—from a clause-by-clause transcription of Definition 1.1 (Remark 2.2)	<code>toIntSpaceC</code> and <code>cutSmall_tendsto_of_src</code> ; <code>logs/reading_layer_axioms.txt</code>	the memoir’s own partial functions and its integral on L ; the apartness structure on X is not represented
Full-set, index-dependent domination is the source’s own §1 convention with the quantifiers displayed (Definition 3.4)	the memoir’s §1 notational convention	not a weakening introduced by this paper
The named public declarations report <code>[propext, Quot.sound]</code> (Section 10)	kernel output, static source scan, checksums, shipped logs	no claim about ZF, about predicativity, or about mathlib as a whole
The countable-avoidance step is discharged constructively (Section 8)	the explicit exception sequence and the nested-interval construction (Appendix B)	the no-go results are not overturned; the claim is presented-reals-relative
The axioms admit a model not concentrated at a single point (Section 7)	<code>Mathdemo.FiniteWeightedIntegrationsSpace</code> with <code>twoPointDef11</code> unconditional	the weights realized are of the form k/m ; no model over a continuum is constructed

2 The setting

2.1 Presented reals

The scalar field is represented by presented real numbers: regular rational approximation sequences equipped with Bishop equality as the working equality relation [1, 3]. A presented real is thus a sequence (x_n) of rationals together with the regularity condition $|x_m - x_n| \leq m^{-1} + n^{-1}$, and two presented reals x, y are *equal* ($x \approx y$) when $|x_n - y_n| \leq 2/n$ for every n : equality of the numbers represented, weaker than literal agreement of the two sequences. A strict inequality $x < y$ is likewise always carried with a witness—an index and a gauge from which the difference stays positive—so that “how far apart, and from when” is part of the datum. Equality and convergence remain distinct: a sequence of presented reals converges to l when its terms approach l at every gauge with a stated rate, not because its terms are Bishop-equal to l . (This section follows the $1/n$ convention of Chapter 2 of Bishop’s *Foundations of Constructive Analysis*; the implementation and Appendix B use the equivalent regularity condition with the dyadic gauge 2^{-k} , and the two are translated by an explicit reindexing—Section 11.2. Equality is taken in the implementation in the tail form $\forall k \exists N \forall n \geq N: |x_n - y_n| \leq 2^{-k}$ —the form stated in Definition B.1, and the relation `relEventually` in the code. For regular sequences this is equivalent to the displayed condition—one direction is immediate, and the other follows from regularity, since $|x_n - y_n| \leq 2 \cdot 2^{-n} + 2 \cdot 2^{-m} + 2^{-k}$ for every $m \geq N$ —but the development adopts the tail form as its definition rather than deriving the equivalence internally.) The development does not identify Bishop-equal approximation sequences: the carrier is the type of regular sequences, not a quotient of it, and every statement is a relation between sequences rather than between equivalence classes. This is a transcription of Bishop’s own position in the *Foundations* rather than an implementation convenience: the notes to that chapter state that a real number *is* the regular sequence, and not an equivalence class of regular sequences, arguing that the equivalence-class reading would be inaccurate or meaningless. Analytic statements are formulated so that they respect the Bishop equivalence relation.

Two consequences matter later. First, every real carries its own modulus by construction. Second, given a strict inequality $a < b$ and a third presented real c , the required cotransitivity branch can be *decided from rational approximation data and returned as data*: from sufficiently good rational approximations one can tell which of $a < c$, $c < b$ holds, and one may return that information rather than only the proposition “ $a < c$ or $c < b$ ”. It is this data form of cotransitivity that makes Bishop’s nested-interval construction (Lemma 5.3 below) a construction rather than an appeal to a choice principle; the mechanism by which the branch is decided is written out in Appendix B.2.

That second consequence is Richman’s construction, quoted here because it is precisely the step at which a classical argument divides into cases: “if $a < b$ are rational numbers, then we can show that either $x < b$ or $x > a$. Indeed let x' be a rational number within $(b - a)/3$ of x . If $x' \leq (b + a)/2$, then $x < b$; while if $x' \geq (b + a)/2$, then $x > a$ ” [7]. The branch is settled by one rational comparison at a computable accuracy, and that outcome is the datum returned by the order layer (Appendix B.2). Richman also states the moral this development follows throughout: a case split “only says that the *argument* is discontinuous as a function of the data, not that this discontinuity is an essential feature of the theorem” [7].

Supplying such a datum is the standard device of the subject rather than a local expedient—“we can constructivize a classical notion by supplying additional information”, and moduli witness not only Cauchyness but uniform continuity, uniform convergence, differentiability and monotonicity as well [15, §27.2.1]—and it is not free. Bickford states the price exactly: a theorem proved under a uniform-continuity hypothesis “requires a modulus of continuity as an input”, whereas a hypothesis

with no constructive content needs none [16, §28.1]. The Type-valued route of Section 7 makes that payment explicit; the Prop-facing surface hides it again from callers who do not need it.

2.2 Integration spaces

The measure-theoretic layer follows a Bishop–Cheng style route [2]. We work with *partial* functions: an element of the ambient function class is a pair consisting of a map into the presented reals and a domain, and the sum of two partial functions is defined on the intersection of their domains.

Both features are the source’s own choices. A measure-first constructive development is possible—it is carried out in the memoir itself—but the earlier book “adopted the much more convenient Daniell-integral approach”, postulating L and E side by side and building measurability and the measure from them [9, §13.1.2]. Admitting functions defined only almost everywhere is equally deliberate, and Chan names its cost: it “does add the small task of checking, whenever we create an integrable function from prior ones, that the condition of being defined a.e. is duly inherited” [9, §13.1.2]. That task is a substantial part of the partial-function bookkeeping below.

All results below are proved relative to an explicitly quantified structure, an *integration space* in the following working sense.

Definition 2.1 (Integration space, working form; cf. the source’s Definition 1.1). An integration space is a triple (X, L, I) consisting of a type X , a class L of partial functions on X with values in the presented reals, and a functional $I : L \rightarrow \mathbb{R}$, such that:

- (A1) L and I respect Bishop equality of partial functions;
- (A2) L is closed under addition, multiplication by a scalar, and absolute value— $f + g$, αf , and $|f|$ lie in L whenever $f, g \in L$ and $\alpha \in \mathbb{R}$ —and under truncation at a constant carrying a positivity witness: $\min\{f, \alpha\}$ lies in L whenever $f \in L$ and $0 < \alpha$ is held as data. Truncation at 0 is not a field but a derived lemma: $\min\{f, 0\} = \frac{1}{2}(f - |f|)$, the negated negative part of f , follows from linear combination, absolute value, and respect for equality (Appendix B.6);
- (A3) I is linear: $I(f + g) = I(f) + I(g)$ and $I(\alpha f) = \alpha I(f)$;
- (A4) I is positive: $I(f) \geq 0$ whenever $f \in L$ is pointwise nonnegative on its domain;
- (A5) the two truncation limits hold for every $f \in L$:

$$I(\min\{f, n\}) \longrightarrow I(f), \quad I(\min\{|f|, n^{-1}\}) \longrightarrow 0 \quad (n \geq 1);$$

(the implementation phrases the small truncation with the dyadic gauge 2^{-n} ; the difference of index domains is absorbed by the gauge translation of Section 11.2);

- (A6) (*constructive continuity*) if $f \in L$, if (f_n) is a sequence of pointwise nonnegative members of L , if $\sum_n I(f_n)$ converges, and if $\sum_n I(f_n) < I(f)$, then there is a point x lying in the domain of f and of every f_n such that $\sum_n f_n(x)$ converges and $\sum_n f_n(x) < f(x)$.

Axiom (A6) is Bishop and Cheng’s clause (2), the constructive analogue of the continuity condition for a Daniell integral, stated in the positive form which returns a point at which a series of nonnegative functions stays below a given function. The two limits in (A5) are the same pair of limits that Spitters isolates in his definition of an integration f -algebra [22].

Remark 2.2 (Fidelity: a clause-level strict weakening of Definition 1.1). Definition 2.1 is *not* a verbatim rendering of Bishop and Cheng’s Definition 1.1 [2]. It is *strictly weaker*, at the level of the displayed clauses, than Definition 1.1 as encoded in this development—the encoding, and the two scope qualifications it carries, are made precise below. Here and below, “clauses (1)–(4)” refer to the four clauses of the source’s Definition 1.1—(1) closure under linear combination, absolute value, and $\min\{f, 1\}$, together with linearity of I ; (2) series continuity, yielding a point; (3) normalization $I(p) = 1$; (4) the two truncation limits—while (A1)–(A6) refer to Definition 2.1.

In two respects Definition 2.1 is weaker. The source requires in addition that X be inhabited and that there exist $p \in L$ with $I(p) = 1$; neither requirement appears above. Definition 2.1 is therefore satisfied by a degenerate zero-integral model, shipped with the artifact (Section 7.3). A second, *normalized* example is point evaluation at a fixed point (`Mathdemo.DiracIntegrationSpace`), for which the constant function 1 lies in L with $I(1) = 1$, so that clause (3) holds. Both source-level limits of clause (4) are proved, so the model is unconditional. For point evaluation they reduce to statements about the truncations of a single real: $\min\{c, n\} \rightarrow c$ holds with a constant modulus once n passes an Archimedean bound for c , and $\min\{|c|, 1/(n+1)\} \rightarrow 0$ holds because the truncation is squeezed between 0 and $1/(n+1)$; the module also proves the auxiliary dyadic limit $\min\{|c|, 2^{-n}\} \rightarrow 0$. The Archimedean bound is the witness $2^{\text{mulArchBound}(c)}$ of `exists_nat_geC` written out explicitly, since that lemma is a `Prop`-valued existential and cannot supply a modulus, which is data. Since the adapter is hypothesis-free as of v0.4.1 (below), this clause-by-clause encoded model yields a model of Definition 2.1 with no further hypothesis.

The closure clause of (A2) is stated for constants carrying a positivity witness, and the restriction is not cosmetic. Clause (1) asks only that $\min\{f, 1\} \in L$, and the constants recoverable from it are positive: the source’s own remark derives $\min\{f, n\} = n \cdot \min\{n^{-1}f, 1\}$. Closure under truncation at an *arbitrary* constant is not available, and the obstruction is visible in Bishop and Cheng’s own canonical model $(X, C(X), \mu)$ of their Definition 7.1— X locally compact, $C(X)$ the continuous functions of compact support, μ a positive measure: one has $\min\{f, \alpha\} = \alpha$ off the support of f , so for $\alpha < 0$ and non-compact X the truncation leaves $C(X)$ (Appendix B.6). An interface demanding truncation at arbitrary constants would therefore not be implied by Definition 1.1, and the two would not be ordered by logical strength.

Restricting the clause to $\alpha \geq 0$ would not suffice either. The recovery route the source itself supplies is $\min\{f, n\} = n \cdot \min\{n^{-1}f, 1\}$, which requires α^{-1} and hence a witness that α is apart from 0; the case $\alpha = 0$ is recovered separately, since $\min\{f, 0\} = (f - |f|)/2$ follows from clause (1) and closure under $|\cdot|$ by a lattice identity that needs no case analysis (Appendix B.6). Constructively one cannot decide between $\alpha = 0$ and $\alpha > 0$, and we have found no uniform construction from $\alpha \geq 0$ —that is, from $\neg(\alpha < 0)$ —alone. The source-derivable form, sufficient for every use in the present development, is therefore truncation at constants *carrying a positivity witness*, or at 0: exactly the two kinds of constant the development truncates at. The closure clause of Definition 2.1 is accordingly the field `cutPos_mem`, truncation at a constant accompanied by positivity data (`PosEventuallyData`); truncation at 0 is the derived lemma `cutZero_mem` rather than a field; and a witness type `CutConstWitnessC`, with a positive branch and a zero branch, wires the evidence through the development. Every use site goes through with the canonical witness; a survey of all use sites found none truncating at a general constant. We emphasize that we have not proved that $\alpha \geq 0$ is insufficient; the claim is only that the source’s route needs a witness and that no other constructive route is evident to us.

The adapter is hypothesis-free. The module `Mathdemo.SourceIntegrationSpaceDef11` contains a clause-by-clause transcription `IntegrationSpaceDef11` of the displayed Definition 1.1 together with an adapter `toIntSpaceC` into Definition 2.1: *all fourteen fields of the working interface are machine-derived, without extra hypotheses, from the clause-by-clause transcription*. Thus Def-

inition 1.1 supplies every field required by Definition 2.1. (No gauge convention is carried across: the transcription states the small-cut limit of clause (4) at the source’s own gauge n^{-1} , built as a presented real from a positivity witness and with no rational inverse assumed, and the passage to the dyadic gauge 2^{-n} used by Definition 2.1 is itself machine-checked as `IntegrationSpaceDef11.cutSmall_tendsto_of_src`, the reverse gauge comparison `srcGauge_le_halfPow` is used by the point-evaluation model.) Two scope qualifications bound this claim. First, the ambient function class is rendered as the memoir renders it: an element of `BFunc` is a domain together with a map defined *only on that domain*, and the integral is a map defined *only on L* , so that the transcription asks for no datum the source does not supply. What is still not represented is the memoir’s apartness structure on X and the strong extensionality it builds into $F(X)$: those are omissions, not additional demands. Second, at the level of the four displayed clauses the only remaining differences are the omissions of inhabitedness and normalization, both pure weakenings. Consequently the current Definition 2.1 is strictly weaker than the transcribed Definition 1.1, and *every integration space in the sense of Definition 1.1 satisfies Definition 2.1 and the theorems of this paper apply to it*. The clause-level weakening is proper: the zero-integral model on `PUnit` satisfies Definition 2.1 but, having $I \equiv 0$, cannot satisfy the normalization clause $I(p) = 1$ of Definition 1.1, since 1 is apart from 0 in the presented reals. The intermediate steps follow the source: clause (1), whose linear-combination part is the assertion $\alpha f + \beta g \in L$ together with $I(\alpha f + \beta g) = \alpha I(f) + \beta I(g)$, yields closure under addition and under scalar multiplication and the additivity and homogeneity of I (each through the equality of $F(X)$); clause (2) yields Corollary 1.3 at $k = 1$ (`pos_witness`) and from it `I_nonneg`; and clause (1)’s $\min\{\cdot, 1\}$ yields `cutPos_mem` by the same algebra as the source’s positive-integer remark, generalized to constants carrying positivity data. Every one of these declarations reports `[propext, Quot.sound]`.

Two further observations complete the inventory. First, Definition 2.1 carries `I_nonneg` as a field, but need not: it is derivable from clause (2), by the derivation the source performs in Lemma 1.2, Corollary 1.3, and the sentence following Corollary 1.3, and that derivation is machine-checked (Appendix B.6). Second, and this is *not* a difference: clause (2) is rendered with a point-returning structure rather than a `Prop`-valued $\exists$, and the limits of clause (4) with an explicit modulus rather than an ε - δ statement. That is faithfulness rather than strengthening, since Bishop’s $\exists$ asserts under the BHK reading that the object can be constructed, and a Bishop limit always carries a modulus; the `Prop`-valued reading would be the weaker one. As a Lean interface, however, the data-carrying rendering is a data refinement of the corresponding `Prop`-valued statement rather than an equivalent restatement (Section 7.1).

The full field list is machine-readable in the artifact, so a reader can check exactly which hypotheses the theorems consume.

3 Integrable sets, full sets, profiles

This section collects the notions used in the statement of the theorem. All of them are Bishop and Cheng’s [2]; we recall them because the theorem cannot be read without them.

Definition 3.1 (Full set; the source’s Definition 1.9). A subset $A \subseteq X$ is a *full set* if it contains a countable intersection of domains of integrable functions.

Definition 3.2 (Complemented set; the source’s Definitions 2.1, 2.2(d)(e), 2.3). A *complemented set* is an ordered pair $A \equiv (A^1, A^2)$ of subsets of X such that $x \in A^1$ and $y \in A^2$ imply $x \neq y$. We write $x \in A$ for $x \in A^1$ and $x \in -A$ for $x \in A^2$, and put $-A \equiv (A^2, A^1)$ and $A - B \equiv A \wedge (-B)$. A complemented set is *integrable* when its characteristic function is, and its *measure* $\mu(A)$ is the integral of that characteristic function.

Complemented sets carry their own witnesses of non-membership; this is what replaces the classical complement, for which one has no constructive access.

Definition 3.3 (Convergence in measure; the source’s Definition 4.11). A sequence (f_n) of measurable functions *converges in measure* to a function f defined on a full set $D(f)$ if for every integrable set A and every $\varepsilon > 0$ there is N such that for every $n \geq N$ there is an integrable set B with

$$B^1 \subseteq A^1 \cap D(f) \cap D(f_n), \quad \mu(A - B) < \varepsilon, \quad |f(x) - f_n(x)| < \varepsilon \text{ for all } x \in B.$$

Note that the good set B , its integrability, the small-complement estimate, and the pointwise closeness all appear in the statement; nothing is left to an external library convention.

Definition 3.4 (Domination on full sets; the source’s §1 convention with the quantifiers displayed). In §1 of the memoir, Bishop and Cheng adopt the notational convention that, for arbitrary $f, g \in F(X)$, the inequality $f \leq g$ means that $f \leq g$ holds *on some full set*. The domination hypothesis $|f_n| \leq g$ of their Theorem 4.15 is therefore already a full-set inequality on the source’s own reading, and since the inequalities for different n are separate statements, the full set may depend on n . The present definition displays that quantification. A sequence (f_n) of integrable functions is *dominated on full sets* by an integrable g if for every n there is a full set F_n such that $|f_n(x)| \leq g(x)$ for every $x \in F_n$ at which the relevant values are defined. The full set is allowed to depend on n .

Finally, the notion at the heart of the proof.

Definition 3.5 (Profile; the source’s Definition 3.1). Let $[a, b] \subset \mathbb{R}$ be a compact proper interval and let $C[a, b]$ be the continuous functions on it. Let $F \subseteq C[a, b]$ satisfy

- (a) $0 \leq f \leq 1$ for every $f \in F$;
- (b) the constant functions 0 and 1 belong to F ;
- (c) if $a \leq u < v \leq b$ then some $f \in F$ satisfies $f(t) = 0$ for $t \leq u$ and $f(t) = 1$ for $t \geq v$.

Let $\lambda : F \rightarrow \mathbb{R}$ be non-decreasing, i.e. $\lambda(f) \leq \lambda(g)$ whenever $f \leq g$ pointwise on $[a, b]$. Then (F, λ) is a *profile* on $[a, b]$.

The example to keep in mind is $\lambda(f) = \int_a^b f(x) dx$; this sentence is the source’s own. A profile is thus a monotone functional on a supply of ramp functions, and condition (c) says that the supply is rich enough to separate any two levels. (The formalization states the definition in a coded form in which condition (c) returns the separating function; Appendix B.4. The coded structure records (a)–(c) and the monotonicity of λ , and carries *no* continuity clause: where the source takes $F \subseteq C[a, b]$, the formalized profile requires of its members only the displayed conditions. This weakens the hypothesis, so the formalized Theorems 5.1 and 5.2 are correspondingly more general than the source’s. The structure has no continuity field, so nothing downstream can depend on continuity.)

Definition 3.6 (Smooth level; introduced in the source inside the text of its Theorem 3.5). A point $t \in [a, b]$ is *smooth* for the profile (F, λ) if there is a number $\bar{\lambda}(t)$ such that for every $\varepsilon > 0$ there is $\delta > 0$ with the property that

$$|\lambda(f) - \bar{\lambda}(t)| < \varepsilon$$

for every $f \in F$ with $f(x) = 1$ for all $x \geq t + \delta$ and $f(x) = 0$ for all $x \leq t - \delta$.

For the model example $\lambda(f) = \int f$, smoothness at t says that ramps which turn on inside a small window around t all have nearly the same integral: the profile does not jump at t . Smoothness is therefore the constructive substitute for continuity of a monotone function at t , and $\bar{\lambda}(t)$ is the common value.

4 The theorem

Theorem 4.1 (Dominated convergence; the source’s Theorem 4.15). *Let (X, L, I) be an integration space in the sense of Definition 2.1. Let f and f_n ($n \in \mathbb{N}$) be integrable functions on X , and assume:*

- (i) $f_n \rightarrow f$ in measure, in the sense of Definition 3.3; and
- (ii) there is an integrable g dominating (f_n) on full sets, in the sense of Definition 3.4.

Then

$$\forall \varepsilon > 0 \exists N \forall n \geq N, \quad |I(f_n) - I(f)| < \varepsilon.$$

On numbering: this paper numbers items by section, so the statement above is Theorem 4.1, and it is the same assertion as Theorem 4.15 of the memoir (the differences in how the hypotheses are phrased are discussed in Remark 4.3 and Section 6); that both numbers happen to begin with “4” carries no meaning. Appendix A.1 tabulates the correspondence between the numbering of this paper and that of the source. In the sequel, “Theorem 4.1” refers to this paper’s numbering and “Theorem 4.15” to the source’s.

Remark 4.2 (Scope visible in the statement). The limit f is itself assumed integrable; the theorem does not derive the integrability of an otherwise merely measurable limit. The result is also relative to the structure (X, L, I) and to presented reals. These are mathematical hypotheses of the theorem, not axioms hidden from Lean’s dependency report.

Remark 4.3 (The mode of the hypotheses). Two features of the hypotheses are worth flagging for comparison with other formulations. First, the mode of convergence assumed is convergence in measure, not pointwise convergence. Second, the domination hypothesis is a full-set hypothesis, and the full set is allowed to vary with the index n ; a single global pointwise bound is not required. The second feature is *not* a weakening introduced by this paper: it renders explicit what the source’s §1 convention already means under Theorem 4.15 (Definition 3.4). Likewise, the difference between the error majorant $H = |g| + |f|$ used in the proof below and the source’s $2g$ is not a difference of hypotheses but a difference of proof route: the memoir’s displayed estimate uses $2g$, which presupposes an inherited full-set bound $|f| \leq g$; that inheritance is classically standard but is neither a stated hypothesis of Theorem 4.15 nor established in the displayed proof, and the verified development does not derive it from convergence in measure and the bounds on the f_n , so it uses the immediately available majorant instead (Section 6). Section 9.1 compares the hypothesis with the corresponding CoRN hypothesis.

5 Profiles, level sets, and countable avoidance

The proof of Theorem 4.1 rests on three results. The first two are Bishop and Cheng’s; the third is the constructive content of the step that the classical proof takes for granted.

Theorem 5.1 (Smoothness away from an explicit sequence; the source’s Theorem 3.5). *Let (F, λ) be a profile on $[a, b]$. Then all but countably many points of $[a, b]$ are smooth in the sense of Definition 3.6.*

The proof is what matters here. Bishop and Cheng choose an increasing sequence $\{n(k)\}$ of positive integers with $(n(k) + 1)k^{-1} > \lambda(1) - \lambda(0)$, collect the finitely many points $\{t_1^k, \dots, t_{n(k)}^k\}$ excluded at stage k , and set $T = \{t_j^k\}$. Every t distinct from every member of T is then shown to be smooth, with $\bar{\lambda}(t)$ constructed explicitly. Thus the conclusion is not that a certain set “is

countable” with an existentially hidden enumeration: the exceptional levels arrive as an *explicit sequence*, constructed by the source’s own proof. The formalization’s contribution at this point is packaging rather than strengthening: it flattens the proof-internal data into a first-class $\mathbb{N}$ -indexed sequence that later constructions consume directly (Appendix B.5).

Theorem 5.2 (Integrability of level sets; the source’s Theorem 3.6). *Let (X, L, I) be an integration space and h an integrable function. Then for all but countably many $t > 0$ the complemented sets*

$$A \equiv (\{x : h(x) \geq t\}, \{x : h(x) < t\}), \quad B \equiv (\{x : h(x) > t\}, \{x : h(x) \leq t\})$$

are integrable and have the same measure.

The bridge between the two theorems is the profile built from h . For $0 < u < v$ let $f_{u,v}$ be the ramp which is 0 below u , rises linearly on $[u, v]$, and is 1 above v , and set $\lambda_0(f_{u,v}) = I(f_{u,v} \circ h)$; the composite is integrable because

$$f_{u,v} \circ h = \min\{(v - u)^{-1}(h - \min\{h, u\}), 1\}.$$

Restricting to $[a, b]$ gives a profile in the sense of Definition 3.5, and $\bar{\lambda}(t)$ is the common measure of A and B . So the distribution function of h appears as a profile, its jump levels appear as the explicit sequence T , and integrability of a level set is available exactly at levels away from T . (Constructively, the two pairs A and B are built independently: since $\leq$ is the negation of $<$, neither pair is the complement of the other, and the classical one-line passage from A to B is not available; Appendix B.8.)

This is where a constructive proof must do work that the classical proof does not.

Lemma 5.3 (Countable avoidance by nested intervals; Bishop). *Let $a < b$ be presented reals and let (q_n) be a sequence of presented reals. Then one can construct t with $a < t < b$ and t apart from q_n for every n .*

The construction is the familiar one: shrink a nested sequence of closed intervals inside (a, b) , at stage n choosing a subinterval that avoids q_n , and take the limit. The step “choose a subinterval avoiding q_n ” is a case distinction on the position of q_n relative to two candidate subintervals, and it is exactly here that presented reals are used: the cotransitivity split is decided from rational approximations and returned as data, so the sequence of intervals is a construction and not a sequence of arbitrary choices. The limit exists because the left endpoints form a Cauchy sequence with an explicit modulus. A complete proof, including the reason why the classical midpoint bisection is not available and the margin that replaces it, is given in Appendix B.3.

Applying Lemma 5.3 to the exceptional sequence T of Theorem 5.1 constructs an admissible level, and hence, by Theorem 5.2, integrable level sets at that level. The following consequence is the form in which the profile theory enters the convergence argument.

Proposition 5.4 (Uniform complement estimate). *Let H be a nonnegative integrable function and let (e_n) be a sequence of nonnegative integrable functions such that for each n there is a full set on which $e_n \leq H$ (in the application to Theorem 4.1, $e_n = |f_n - f|$ and $H = |g| + |f|$). Then for every $\varepsilon > 0$ one can construct an integrable set A , an index bound N , and a measure threshold $\delta > 0$ such that for every $n \geq N$ and every integrable complemented set C ,*

$$\mu(A - C) < \delta \implies I(e_n - \chi_C e_n) < \varepsilon.$$

Proposition 5.4 corresponds to the hypotheses of the source’s Lemma 4.14 together with the construction inside its Theorem 4.13, under the equivalent reparameterization $B := -C$: the source’s smallness condition $\mu(A \wedge B) < \delta$ is the condition $\mu(A - C) < \delta$ above, since $A - C = A \wedge (-C)$. It is what makes the “outside a good set” half of the classical proof work without an almost-everywhere argument: the smallness of the integral off A is obtained from the measure of a level set of H at an admissible level, and admissibility is supplied by Lemma 5.3. The triple (A, N, δ) is returned as data because the sequel consumes it (Appendix B.10).

6 Proof of Theorem 4.1

The proof follows the decomposition below. The arrows indicate constructed data or a theorem application, rather than an informal implication between black-box library results.

profile partition data and an explicit exception sequence
 $\Downarrow$ presented-real countable avoidance
 an apart threshold and integrable level sets
 $\Downarrow$ profile estimates
 a uniform small-complement integral bound
 convergence in measure + full-set domination
 $\Downarrow$ good-set/complement decomposition
 $I(|f_n - f|) \rightarrow 0$
 $\Downarrow |I(f_n) - I(f)| \leq I(|f_n - f|)$
 $I(f_n) \rightarrow I(f).$

More concretely, the proof performs the following steps.

- (1) For each n , form the absolute-error function $e_n = |f_n - f|$. From the full-set domination of f_n by g , derive a full-set bound of e_n by the nonnegative integrable function

$$H = |g| + |f|.$$

The appearance of $|f|$ is why the verified error majorant is not advertised simply as $2g$: the source’s own chain passes through $2g$, which presupposes $|f| \leq g$, and the verified route does not establish that inherited bound, so it uses the immediately available majorant instead.

- (2) From H , construct the dyadic smooth-level data needed by the profile argument. Internally, Theorem 5.1 returns a concrete exception sequence; Lemma 5.3 constructs a presented real apart from that sequence, and hence an admissible level at which the relevant upper and lower level sets are integrable by Theorem 5.2.
- (3) The profile estimates at these levels yield Proposition 5.4: for a prescribed positive tolerance, there are an integrable set A , an index bound, and a positive measure threshold such that every sufficiently late error function has small integral outside each sufficiently large integrable subset of A .
- (4) Convergence in measure (Definition 3.3) is invoked at the minimum of the profile threshold and a good-set bound. It supplies, for each sufficiently large n , an integrable good set $C \subseteq A$ on which f_n is close to f and whose complement in A has small measure. One then estimates $I(e_n)$ by its contribution on C plus its contribution outside C ; this is the two-part estimate corresponding to Bishop and Cheng’s Lemma 4.14.

(5) Finally, the constructive triangle inequality

$$|I(f_n) - I(f)| \leq I(|f_n - f|)$$

gives the conclusion of Theorem 4.1. □

Part II

The formalization and its audit

7 The Lean development

The development is carried out in Lean 4 [31], toolchain v4.30.0. The carrier of the presented reals is a concrete type,

$$\text{BishopCReal.CReal} = \text{RegularSeq},$$

not a typeclass interface for an abstract constructive real-number structure; the rational layer is itself a quotient constructed inside the development. A quotient of the presented reals by Bishop equality, `BishopCReal.CRealQuot`, is also constructed, but it is used only as a transport device inside proofs: a ring identity between presented reals is discharged by `ring` in the quotient and the conclusion is pulled back to the representatives by `Quotient.exact`, so no statement on the theorem surface is formulated about equivalence classes. Internally the quotient does occur in statements—the closure contains declarations about `CRealQuot` itself, in the layer that establishes its ring structure—but none of the public aliases, the reading layer, or the measure-theoretic development quantifies over equivalence classes. `Quot.sound` in the audit output comes from the kernel’s quotient machinery: these two constructions, and function extensionality, which Lean derives from quotients and which the development uses through set extensionality. The data form of cotransitivity discussed in Section 2 is the declaration

$$\text{lt_cotrans_data} : \text{lt } a \ b \rightarrow \forall c, \text{ PSum } (\text{lt } a \ c) (\text{lt } c \ b),$$

which returns the branch rather than only its Prop-valued disjunction. The displayed declaration is the field `BishopC.COF.lt_cotrans_data` of the ordered-field interface, instantiated at the rational layer; it is the only data-valued branch the avoidance construction consumes. Its presented-real counterpart, `regularSeqLtData_cotrans`, has the same shape one level up.

Integration spaces in the sense of Definition 2.1 are the structure `BishopSec1P.IntSpaceC X`, whose fields are

$$\text{L, I, L_resp, I_resp, add_mem, smul_mem, abs_mem, cutPos_mem, I_add, I_smul, cutNat_tendsto_raw, cutSmall_tendsto_raw, I_nonneg, continuity,}$$

corresponding to the axioms (A1)–(A6) as tabulated in Appendix A. The truncation field is `cutPos\_mem`, which takes the positivity data `PosEventuallyData`; truncation at 0 is the derived lemma `cutZero\_mem`, and the witness type `CutConstWitnessC` wires the evidence to the use sites (Remark 2.2). Integrable representatives, integrable complemented sets, relative integrals, profiles, and convergence in measure are represented by explicit structures or by Prop-facing predicates whose existential witnesses are opened only inside Prop goals. The final dominated-convergence statement is reached through an explicit majorant/profile route rather than by invoking a measurable-selection principle.

Theorem 4.1 is the implementation theorem

BishopSec3P.bishop_cheng_thm_4_15_propC,

exported under the paper-facing alias

ChoiceFreeMeasureDCT.bishop_cheng_dominated_convergence_propC.

Its hypothesis (i) is `ConvergeInMeasureC`, which unfolds exactly to Definition 3.3; its hypothesis (ii) is `DominatedOnFullC`, which unfolds to Definition 3.4; and its conclusion is `RepSeriesTendstoEpsPropC`, the ordinary epsilon formulation displayed in Theorem 4.1.

7.1 Three-layer public interface

The public API has three layers:

Layer I: Type-valued convergence and profile data, data-carrying DCT route	Layer III: Prop-facing full-set DCT
↓	
Layer II: automatic dyadic smooth-level construction	

The two columns are deliberately not joined by an arrow: Layer III is not obtained by applying the Layer I theorem, and Layer I is not obtained from Layer III either. The distinction between the layers is the distinction between a proof that merely asserts convergence and a proof that returns the modulus. In Lean, a statement in `Prop` is proof-irrelevant, and an arbitrary witness hidden in a `Prop`-valued existential cannot in general be eliminated into `Type` (data can nevertheless be recomputed when an independent decidable search produces a witness—Appendix B.12); a statement in `Type` is proof-relevant, and its inhabitants are data that later constructions may consume. A mathematician may read Layer III as the theorem and Layers I–II as the same theorem with the witnesses kept. More precisely, the Type-valued surface is a data refinement rather than an equivalent restatement: it demands and returns witnesses. A direct wrapper from the Prop-facing hypotheses to that surface would have to extract indexed witnesses hidden in `Prop` and would therefore require an additional choice principle. The artifact instead proves the Prop-facing and Type-valued routes separately; this is an interface and implementation claim, not a theorem of logical non-derivability between the two complete statements. “Returns data” means that a proof-relevant term exists; efficient program extraction is not claimed, and the fact that some public aliases are marked `noncomputable` is not evidence of choice either way (Section 10).

Layer I: Type-valued data-carrying route. The declarations

```
ChoiceFreeMeasureDCT.l1_error_convergence_from_majorant_measure_convergenceC
ChoiceFreeMeasureDCT.integral_convergence_from_majorant_measure_convergenceC
ChoiceFreeMeasureDCT.dominated_convergence_from_error_majorant_profileC
ChoiceFreeMeasureDCT.dominated_convergence_from_pointwise_majorant_profileC
ChoiceFreeMeasureDCT.dominated_convergence_from_pointwise_majorant_good_set_profileC
ChoiceFreeMeasureDCT.l1_error_convergence_from_majorant_on_full_dataC
ChoiceFreeMeasureDCT.integral_convergence_from_majorant_on_full_dataC
ChoiceFreeMeasureDCT.dominated_convergence_from_pointwise_majorant_on_full_dataC
```

consume explicit proof-relevant data: convergence moduli, good-set data, integrability witnesses, point-evaluation witnesses, profile/majorant data, smooth-level data, and Type-valued convergence output. In the full-set route, the carrier may depend on the sequence index. A Type-valued convergence interface is not claimed to be novel, since CoRN’s `CvMeasure` is also Type-valued.

Layer II: automatic Type-valued wrappers. The public declarations ending in `_autoC` internally construct the dyadic smooth-level data by `BishopSec3P.lemma43DyadicSmoothDataC_construct`. They therefore avoid exposing an external `Lemma43DyadicSmoothDataC` argument. This includes `dominated_convergence_from_pointwise_majorant_on_full_data_autoC` and the variant combining explicit domination full sets with explicit convergence good sets; the two indexed set families remain distinct.

Layer III: Prop-facing theorem. The declaration `ChoiceFreeMeasureDCT.bishop_cheng_dominated_convergence_propC` accepts Prop-valued `ConvergeInMeasureC` and an existential integrable majorant satisfying `DominatedOnFullC`. It concludes Prop-valued epsilon convergence of the integral sequence. Existential witnesses from convergence and full-set domination are opened only inside Prop proof goals, not assembled into the selector required by the Type-valued full-set route. Consequently the final theorem does not manufacture a global Type-valued choice function. A separate Type-valued interface accepts precisely such data as `DominatedOnFullDataC`: for each index it contains a carrier, a proof that the carrier is full, and the domination proof on that carrier. It then returns `RepSeriesTendsto`, retaining the convergence modulus.

The two routes are therefore proved separately, and the separation is visible in the source. From the uniform-complement estimate onward the assembly exists twice, once producing a `Prop` and once producing a structure: some of the paired declarations differ only in the type of the hypothesis, the type of the conclusion, how the witness is opened, and how the result is assembled; others diverge further—the Prop-valued L^1 endpoint is an explicit proof, while its Type-valued counterpart composes data-valued lemmas. The two assemblies are structurally parallel rather than textually identical. What is not duplicated is the substance: the Lemma 4.3 apparatus—dyadic smooth-level data, level-set sequences, countable avoidance, the stratum where the known obstruction lives—exists once and is called by both. The repeated tail is an implementation consequence of the chosen interfaces: a direct wrapper would have to extract indexed witnesses hidden in `Prop`. The duplication makes that interface boundary visible, but is not by itself a proof that every choice-free implementation must duplicate the tail, or that the two complete statements are logically independent.

7.2 Formal counterparts of Section 5

Theorem 5.1 is exported as

`ChoiceFreeMeasureDCT.profile_smooth_away_from_sequenceC`

(implementation `BishopSec3P.thm_3_5_smooth_aeC`); the partition data underlying it, which is the source’s Lemma 3.4 rather than its Theorem 3.5, is exported separately as

`ChoiceFreeMeasureDCT.profile_partition_dataC`

(implementation `BishopSec3P.lemma_3_4DataC`). Theorem 5.2 is exported in the source’s global form as

`ChoiceFreeMeasureDCT.profile_level_sets_integrable_apart_globalC,`

implemented by `BishopSec3P.thm_3_6_all_posC`: it produces one exceptional sequence $T : \mathbb{N} \rightarrow \mathbb{R}$ good for every positive threshold at once, obtained as the source obtains it, by covering the positive half-line with the intervals $((n+2)^{-1}, n+2)$ and flattening the per-interval sequences. The interval-local core it is assembled from is exported as well, as `profile_level_sets_integrable_apartC,`

since that is the form the dominated-convergence route consumes; it packages the conclusion that apart level thresholds give integrable upper and lower level sets with the expected profile measure. The word *apart* in those names is the load-bearing one: the thresholds are required to be apart from the explicit exception sequence returned by Theorem 5.1, and such thresholds are obtained by Lemma 5.3. Proposition 5.4 is exposed as

`ChoiceFreeMeasuredCT.uniform_complement_from_profile_levelsC,`

implemented by `BishopSec3P.lemma43UniformComplementData_of_majorantC`; the variant whose domination hypothesis is itself a full-set hypothesis, as in the statement of Proposition 5.4, is `lemma43UniformComplementData_of_majorantOnFullDataC`. Lemma 5.3 is formalized on the presented reals as the structure `Thm36B1ApartPointDataC`—a point t with $a < t < b$ together with, for every n , a witness that t is apart from the n -th term of the prescribed sequence—and its canonical inhabitant `thm36B1_apartPointDataC`, obtained from the data-valued nested-interval construction: the interval sequence, monotonicity of the endpoints, width shrinkage, Cauchy data for the left endpoints, and the limiting apart point. `thm36B_smoothPointDataC_construct` applies it to the profile exception sequence, and `lemma43DyadicSmoothDataC_construct` iterates that at the dyadic gauges to supply the smooth-level data consumed by Proposition 5.4; the latter is the declaration the public `_autoC` route calls. The generic-route counterparts `thm36B1_exists_apart_in_interval`, `thm36B1_apart_data` and `lemma_4_3_A_n_apart_data` are earlier declarations over an abstract ordered field and are not on the presented-real path taken by the public theorems. Step (4) of the proof in Section 6 is `lemma_4_14_local_two_epsilonC`, and step (5) is `integral_diff_lt_of_abs_error_integral_ltC`.

7.3 Artifact additions in this development line

This development line separates the public L1 error-convergence endpoint from the final integral-convergence endpoint. It adds the explicit full-set Type-valued route described above and a good-set convergence wrapper: the convergence data carries the good set, its integrability, membership in the ambient set, Type-valued point-evaluation witnesses, small complement measure, and the pointwise error estimate restricted to the good set.

The artifact further includes CReal-native support for the countable-avoidance construction used in the smooth-level interface. The formalized support constructs the data-valued nested interval step, the resulting interval sequence, left/right monotonicity, interval-width shrink, Cauchy data for the left endpoints, the limiting apart point, and the dyadic smooth-level data constructor.

`Mathdemo.ChoiceFreeDCTExamples` gives abstract application examples showing that both the explicit-smooth and automatic public aliases can be invoked as theorem inputs in proof terms. The file `Mathdemo.ChoiceFreeDCTConcreteExamples` adds a nonempty concrete zero-integral example on `PUnit`. It constructs the required good-set convergence and domination data and invokes the automatic good-set wrapper. This concrete example is intentionally degenerate—it is admissible precisely because Definition 2.1 imposes no normalization axiom (Remark 2.2). The file `Mathdemo.DiracIntegrationSpace` adds the normalized point-evaluation model, in which every clause of Definition 1.1 is discharged with no hypotheses, and which therefore also yields a model of Definition 2.1 through the adapter. It is a nondegenerate concrete model, although a simple one: the underlying functional is evaluation at a single point. `Mathdemo.FiniteWeightedIntegrationSpace` lifts that to finitely many points: for a list of points, repetition realizing integer weights, the functional $I(f) = \sum_{i \leq N} f(p_i)$ discharges every clause of Definition 1.1, and `twoPointDef11` instantiates it at two points with the normalizing constant $1/2$, so that the measure takes more values than a single point mass allows. The mathematical content beyond the Dirac model is clause (2): for a single

point the continuity clause is the identity, while for several the point at which the series stays below the function has to be *found*. It is found by cotransitivity of the strict order along the finite sum, from the positivity witness carried by the strict inequality, with no appeal to choice. Finally, the module `Mathdemo.MathematicalInterface` restates the public surface in the vocabulary of Part I, as a reading layer for a mathematician.

8 Relation to the no-go results

The constructive content of this section is written out, from the side of the constructions, in Appendix B.5. The core of the answer is a *confluence* of two conditions: (a) the input to the avoidance step is at hand as data—the explicit family that the source’s own proof constructs—rather than a “countable set” hidden behind a proposition; and (b) the avoidance construction for that sequence (nested intervals) can be carried out over the order data of presented reals. The explicit-sequence form by itself does not evade the obstruction: over an abstract real-number object, a general principle taking an explicit sequence and returning a real apart from all of its terms would itself be a sequence-avoidance principle of exactly the kind the no-go results concern. What makes the avoidance possible is the restriction of the object, (b); condition (a) is the formal precondition that makes it applicable.

The route formalized here passes through the profile theorem, and the profile theorem is precisely where the choice-free programme is known to be obstructed. A choice-free claim for this route must therefore specify the real-number object and mathematical interface relative to which it is made.

Spitters reports Richman’s formulation that measure theory and the spectral theorem are the major challenges for a development without countable choice [22, §1], attributing it to a 2005 note of Richman’s that we have not consulted; Richman’s published case studies of that programme are the Hilbert syzygy theorem, trace-class operators, and the fundamental theorem of algebra [20], the last of which he carried out in full without choice [18]; the weak choice principles that remain in play are analysed in [19]. Spitters took up Richman’s challenge for integration theory [22]. In the choice-free version he observes that the profile theorem—the positive form of the classical fact that an increasing function on the reals has at most countably many discontinuities—*implies the uncountability of the reals*: for the integration space

$$I(f) = \sum_n 2^{-n} f(q_n)$$

determined by a sequence (q_n) of points of $[0, 1]$, and for $f = \text{id}$, a smooth point of the profile of f must be distinct from—in his sense, which is positive distance, hence apart from—every q_n . He then exhibits a sheaf model over $\mathbb{R}$, built from a sequence of linear functions forming a dense 45° lattice—that is, lines of slope ± 1 —in which the statement “for every sequence of reals there is a real apart from all of them” fails, and concludes that a proof of the profile theorem without choice is not to be expected; his development consequently obtains the approximation results it needs from a pointfree Stone representation theorem instead of from Bishop and Cheng’s approximation theorem. Petrakis and Zeuner record the same obstruction. Their formulation of what remains open is broader than a single theorem: noting—in their words—that Spitters gave a proof of a choice-free version of the profile theorem by the pointfree route, they ask whether the basic *applications* of profile theory can be recovered inside predicative Bishop–Cheng measure theory through choice-free variants of its basic notions and results [25, §11]. Their phrase is stronger than what the cited note claims for itself: Spitters writes that there is no hope of proving the profile theorem without choice, and that of its approximation consequence it is not clear to him whether it

can be proved without choice; what he supplies by the pointfree route is that consequence, not the theorem [22]. One level down, Lubarsky shows that intuitionistic set theory without choice does not prove that the constructive Cauchy reals are Cauchy complete, and records Richman’s observation that what is actually provable is the completeness of *sequence–modulus pairs* rather than of the reals themselves [21]. In his later survey the alternative is named in passing: excluded middle enters that construction as a case split “because the alternative is to use Countable Choice, and most would automatically avoid that as just too dishonest”, and formulations of “almost Cauchy” that are “mutually equivalent under Countable Choice” are shown there to come apart without it [13, §22.2.2, §22.7]. Distinctions of that kind are invisible while choice is available; the profile route is one more place where withdrawing it makes one visible. Richman puts the resulting order of motivations in its strongest form: “I have come around to thinking that sequences are undesirable, therefore I don’t want to use countable choice, rather than that I don’t want to use countable choice, therefore I must abandon sequences” [11, §19.2]. Behind all of it stands Diaconescu’s theorem: in a topos, where AC reads “Every epimorphism has a section”, “an intuitionistic model of set theory with the axiom of choice has to be a classical one” [6]. Choice in that form is therefore not a convenience a constructive development may help itself to, and a foundation may be designed to *lack* it rather than merely to avoid using it.

The present formalization neither refutes these results nor settles the open problem. It is compatible with them, and the mechanism is worth stating precisely, because it is the content of the word “presented” in the title.

- (i) **The reals are presented.** A real is a regular sequence of rationals, so it carries its own modulus. Given $a < b$ and c , the cotransitivity split needed by the construction is returned as data from rational approximations. Bishop’s nested-interval argument (Lemma 5.3) is therefore available as a *construction*, not as a choice principle. The artifact formalizes it and audits it with the same axiom output as the rest of the development. The mechanism by which the branch is returned as data—the one-point certificate and strong gauge extraction—is written out in Appendix B.12. Maietti and Sambin diagnose the same step from the other side: turning a formal point of the formal topology of the reals into a Cauchy sequence needs dependent choice in the setoid model over Martin-Löf type theory, and in a foundation without choice “this proof cannot be carried out”—because “what actually happens in MLTT is that the splitting of points is already given with an operation choosing an interval where the point is, and hence from this choice a definition by cases can be given similarly to that done classically” [12, §20.4.3]. Here that operation is not inherited from the ambient type theory but supplied, as part of the presentation.
- (ii) **The uncountability step is discharged, not avoided.** Theorem 5.1 as formalized does not assert that the set of non-smooth levels “is countable” with an existentially hidden enumeration; it returns an *explicit exception sequence* and concludes smoothness at every level apart from it. The countable-avoidance construction of (i) is then applied to that exception sequence to obtain an admissible level threshold. In short, the development pays exactly the price that Spitters identifies, and pays it constructively over presented reals.
- (iii) **Completeness is used only in representation-carrying form.** Limits are constructed from an explicitly supplied modulus and are never extracted from a bare Cauchy property. This is precisely the distinction that Lubarsky attributes to Richman, and it is why his independence result does not bear on the development.
- (iv) **Scope of the claim.** Both Spitters and Lubarsky note that, in the relevant models, the Cauchy reals and the Dedekind reals need not coincide [22, §2]; in the models Lubarsky

cites, even the smallest Cauchy-complete set of reals can be a proper subset of the Dedekind reals [21, §7]. We do not undertake here a model-theoretic analysis of which real-number object the counterexample family inhabits. Our claim is the correspondingly modest one: *over presented reals, and for the explicitly quantified integration-space interface of Definition 2.1, the Bishop–Cheng profile route can be carried out in Lean’s type theory with no appeal to Lean’s `Classical.choice`*. This is not a claim that the profile theorem is provable without choice over an arbitrary constructive real-number object; it is not a predicative result; and it does not address the open problem of [25], which concerns the recovery of the applications of profile theory inside a predicative setting.

Two limits on the word itself should be recorded here. A foundation may be built to *lack* choice: the Minimalist Foundation is “characterized by the lack of whatsoever choice principle, including the so-called axiom of unique choice” [12, §20.1], and in it the reals form no set. Lean is not such a foundation—unique choice is available once `Classical.choice` is admitted—so the audit certifies not that no choice principle is valid but that the named declarations do not *invoke* one. That weaker property is the checkable one, and it is checkable because Lean’s `Exists` is `Prop`-valued rather than a Σ -type: the propositions-as-sets identification that makes choice provable in MLTT is unavailable, so extraction requires the axiom and its absence is mechanically detectable (Section 10). Second, realizability is not elimination. Berger and Seisenberger prove countable and dependent choice realizable and then withhold the conclusion, “since in their proofs the very same principles are used”, adding that one cannot expect elimination for “choices of elements in abstract domains for which no computational methods are available” [17, §29.3]. That last clause names the hypothesis replaced here: over presented reals the domain is not abstract, and the branch is computed from rational approximations.

9 Related work

Bishop–Cheng measure theory and its reconstructions

Bishop and Cheng’s memoir *Constructive Measure Theory* is the mathematical source for the measure-theoretic route formalized here [2]; the surrounding constructive analysis is that of [1, 3], and the integral-first tradition it belongs to goes back to Daniell [4]. Chan’s recent monograph develops constructive probability theory on the same basis and contains an extended treatment of integration and measure [5]. Coquand and Palmgren give a pointfree, measure-algebraic account [23]. Ciraulo develops a more recent pointfree route toward measure theory, including a measure on σ -sublocales; that chapter uses countable choice freely, and notes that carrying the treatment over to choice-free frameworks would require a different definition of the frame of reals [10].

Petrakis and Zeuner have taken, in their own words, the first steps towards a predicative reconstruction of Bishop–Cheng measure theory: they replace the quantification over proper classes in the L^1 construction by set-indexed families of partial functions, and prove that the resulting family of canonically integrable functions is itself a pre-integration space and an appropriate completion of the original one—so the completion is recast predicatively rather than discarded [24, 25]; their development also avoids countable choice. The present work does not claim a predicative reconstruction. The ambient theory has an impredicative universe of propositions: a $\forall$ or an $\exists$ whose body is a proposition is itself a proposition, whatever is quantified over, so that no universe is climbed. Sets are accordingly $X \rightarrow \mathbf{Prop}$ and separation is available impredicatively, and no

attempt has been made to check that each definition of the development stays inside a predicative fragment. Fullness, the order relations on presented reals, and the `Prop`-facing chain of Section 7 are propositions; several of these notions also have `Type`-valued forms, which live in the predicative universes—the data form of the order, and integrable sets, which are structures. The presence of those forms does not make the development predicative, since the impredicative universe is available throughout.

That disclaimer is compatible with a more precise description of the relationship, which is complementary rather than merely disjoint. First, the `Type`-valued face of the principal theorem chain carries as data, in the predicative `Type` universes, exactly what has to be produced: convergence moduli, the carrier chosen at each index, and integrability representations. The remaining fields of those records are propositions—that the carrier is full, that the domination inequality holds on it, and that the terms are close at each gauge—so the `Type`-valued route is not free of `Prop`; what it does not contain is a step that extracts a witness from a `Prop`-valued existential by a choice principle. Fullness is one of those propositions, and the sharpest case: `IsFullC` is a `Prop`-valued existential over sequences of integrable representations, and the data-facing structures carry a proof of it rather than exposing its generating sequence. That is consistent with the preceding sentence rather than an exception to it: the existential is destructured only while proving `Prop`-valued facts about full sets—closure under intersection, and the passage to a single integrable representative—and is never eliminated into a `Type`-valued result or selector. The chain’s only passages from propositions to data are explicit supply of data and decidable search. The negative half of this is certified mechanically by the choice-free audit: in Lean, unconditional extraction of data from a `Prop`-valued existential requires `Classical.choice` (Section 10), so an axiom output confined to `[propext, Quot.sound]` proves the absence of choice-based extraction throughout the audited closure. The positive classification—that every passage from propositions to data proceeds by explicit supply of data or by decidable search—comes from source inspection, and its principal instance is written out in Appendix B.12. Second, integrable functions are handled as representation data from the outset and never pass through the impredicative L^1 -completion; this runs in the same direction as Petrakis and Zeuner’s central device of recasting the completion through set-indexed families—and it was they who carried that device out, as theorems, inside a predicative foundation.

The two contributions therefore divide the two known obstacles of the memoir between them: predicativity is addressed by Petrakis and Zeuner at the level of the theory—in their own words, the first steps towards a predicative reconstruction—while freedom from choice is established here, at the level of machine-checked proofs, for a Lean development over presented reals (for the machine-checked precedent in Coq, see the CoRN comparison below). Determining, inside Lean, which fragments of the present development admit a predicative reading is left as future work, alongside their open problem (Section 8).

Integration in proof assistants

Sacerdoti Coen and Tassi formalized a constructive proof of Lebesgue’s dominated convergence theorem in Matita [29]. Boldo, Clément and Leclerc formalized the Bochner integral in Coq [30].

One of the most substantial formalizations of Bishop-style analysis in a proof assistant is Bickford’s: “We have formalized in Nuprl all of Chapter 2 of Bishop and Bridges, ‘Constructive analysis’” [16, §28.1]. That report is a particularly relevant neighbouring formalization on the mathematical side, and it also marks the boundary that the present work sits next to. Its concluding section states which part of the programme was carried out and which was not: the numerical-analysis methods “can be adapted to yield provably correct, efficient algorithms for operations on Bishop’s real numbers. We have illustrated how this can be done for the functions defined in Chap-

ter 2 of Bishop and Bridges, with the notable exception of general definite integrals” [16, §28.14]. Bickford’s target is thus Bishop–Bridges analysis rather than the Bishop–Cheng memoir, and the general integral is explicitly outside what was achieved there; the route formalized in the present paper begins on the other side of that line. One further observation of his bears on a choice made here: a first attempt that generalized Bishop’s regular sequences to admit arbitrary moduli was withdrawn—“we found that this choice of representation was a mistake” [16, §28.2]—in favour of the source’s own fixed convention, which is the convention kept here (Section 2.1). He also separates constructive existence from executability, reporting a modulus obtainable in principle from Brouwer’s principles that the extracted code could not compute in practice [16, §28.13]; the audit of Section 10 concerns axiom dependence, not running time, and no performance claim is made here. On the classical side, Lean’s `mathlib` contains a dominated convergence theorem for the Bochner integral over a measure space [34, 35]; that development is classical and is not comparable to the present one at the level of hypotheses. The closest prior formalization of the Bishop–Cheng route itself is S  m  ria’s, discussed below.

Constructive reals in proof assistants

Miyamoto’s survey of exact real arithmetic is the closest published account of the representation used here—its Definition 27.1 is the same object, with the regular sequences as the case in which the modulus is the identity—and its treatment of uniformly continuous functions makes the design choice of the present artifact explicit: rather than invoking the fan theorem to obtain the missing bounds, “we require additional information to compute the lower and upper bounds . . . so that we don’t rely on the fan theorem” [15, §27.2.1, Def. 27.13]. Replacing an appeal to an extra principle by an extra field of data is the move the presented reals make for cotransitivity. In the theory of computable functionals behind Minlog the analogue of the `Prop/Type` split appears as an explicit declaration: predicates “can be declared to be either computationally relevant (c.r.) or else noncomputational (n.c.)” [14, §26.3], and the extracted term is determined by that declaration.

Geuvers and Niqui axiomatized the constructive reals of the FTA project—the development that later grew into CoRN—and proved the axiomatization categorical [27]. S  m  ria’s JFLA paper discusses Cauchy-data constructive reals, setoid equality, and the distinction between the constructive and classical real-number layers in Coq’s constructive-real infrastructure [28]; it is cited here for that infrastructure, not as a source for the dominated-convergence theorem.

Booij’s locators [26] approach the same proposition/data axis from the opposite side of the representation divide. There the reals are the extensional Dedekind reals of univalent type theory, carrying no presentation, and computational content is carried per real as additional structure: a *locator*, which replaces the disjunction in the locatedness clause $\forall q < r. (q < x) \vee (x < r)$ by an untruncated sum. His Theorem 3.9.5—for a real number, locators and signed-digit representations are interdefinable—is the dictionary between the two sides at the level of structure; read through it, the presented reals used here live on the locator-carrying side. (His Theorem 3.9.7 states the corresponding equivalence for *mere* existence: that a locator exists, that a signed-digit representation exists, that a rational Cauchy sequence with limit x exists, and that x is a Cauchy real, are equivalent. He reserves “to have” for untruncated structure and “to exist” for the truncation, and item 1 is written $\| \text{locator}(x) \|$.) The exact correspondence—the same decidable-search engine, the same normal form as the one-point certificate, dual answers to the same extensionality obstruction, and the same strategy of replacing choice principles by carried structure—is set out in Appendix B.13. Among proof-assistant neighbours his theory thus occupies a third coordinate, the extensional side with carried structure, distinct both from CoRN’s setoid representations and from the raw presentations used here.

The closest prior formalization: CoRN

The closest prior proof-assistant formalization identified for Bishop–Cheng dominated convergence is Séméria’s Coq/Rocq development in CoRN [36]. The audited community-repository snapshot is CoRN commit `ada7c0b497ff15dd67cf7932c6f20e143a2aee2f`. At that commit, the pinned source file `CMTMeasurableFunctions.v` contains `CvMeasure`, the source’s Lemma 4.14-style declaration `DominatedMeasureCvZero`, and `DominatedConvergence`. The reproduced Rocq 9.0.1 audit packaged with this artifact reports all three declarations as **Closed under the global context**. For each declaration, `Print Assumptions` listed no global axioms in its dependency closure. This describes the declarations’ global axiom footprint; it does not make the theorems free of mathematical hypotheses. Their statements remain parameterized by the explicitly quantified `IntegrationSpace` and `ConstructiveReals` structures, whose fields specify the mathematical setting in which the results are proved. The same reading applies symmetrically to the Lean audit reported here, whose statements are parameterized by the integration space of Definition 2.1.

CoRN also makes the countable-avoidance step explicit and data-valued. In the pinned `CMTprofile.v`, `CRuncountable` and its diagonal variants return a real in a prescribed interval apart from every member of a supplied sequence. The three-index variant feeds the jump-point enumeration used to prove integrability of level sets away from an explicit exception sequence. Its proof covers the supplied sequence by open intervals of controlled total measure in a concrete real-line integration space and extracts a point from the positive-measure remainder. The present Lean development instead implements Bishop’s nested-interval construction (Lemma 5.3) over fixed regular-sequence presentations. Thus both developments discharge the same countable-avoidance obligation by explicit constructions, but by different routes. The two routes also differ in the depth of their prerequisites: the covering argument runs inside a concrete real-line integration space, at a stage where the measure apparatus is already constructed, whereas the nested-interval construction uses only the order structure and rational approximations of the presentations, before any measure theory is in place. The same obligation is discharged at different depths of the respective developments; the contrast is a difference of position, not a ranking.

9.1 Comparison of domination assumptions

The theorem interfaces also differ materially. Both final DCT routes assume convergence in measure, not pointwise convergence. CoRN’s `CvMeasure` hypothesis is Type-valued, and `partialFuncLe` states domination on the intersection of the partial-function domains. The Lean artifact exposes both a Type-valued full-set route, with an explicit indexed selector of full carriers, and a Prop-facing full-set theorem. CoRN quantifies over an abstract `ConstructiveReals` implementation, whereas Lean fixes regular-sequence presented reals; CoRN’s `IntegrationSpace` also contains a distinguished integrable unit of integral one, whereas Definition 2.1 has no normalization field and admits the degenerate model described in Remark 2.2. These differences are not ordered by logical strength, and neither complete theorem surface is claimed to be a formal generalization of the other.

Concretely, CoRN’s public `DominatedConvergence` assumes global pointwise domination:

$$\text{forall } n, \text{ partialFuncLe } (X\text{abs } (fn \ n)) \ g.$$

The Lean artifact exposes the full-set hypothesis in two forms. The Prop-facing theorem assumes that for each n , domination holds on a full set, which may depend on the index (Definition 3.4). The Type-valued theorem takes those carriers, fullness proofs, and bounds as the explicit selector `DominatedOnFullDataC` and preserves a convergence modulus in its conclusion. The formal bridge

`DominatedOnFullDataC.ofGlobal` packages every global pointwise domination family as full-set data; no converse is asserted. Thus, relative to the domination clause of CoRN’s public theorem surface alone, full-set domination is a weaker hypothesis, while the Type-valued presentation asks the caller to supply more reusable proof-relevant data than the Prop presentation. (By Bishop and Cheng’s own notational convention the domination hypothesis of their Theorem 4.15 is also a full-set condition—Remark 4.3; the comparison in this paragraph is with the CoRN declaration, not with the source.)

We do not claim a formal strict generalization of the CoRN theorem, because the ambient real-number and integration-space interfaces differ. CoRN’s `CR_cv` and the Lean Type-valued full-set conclusion are both data-valued, but the two complete theorem statements are not ordered solely by logical strength. Across the Lean pointwise-majorant wrappers, the verified error majorant is $g + |f|$ or $|g| + |f|$, depending on the route, not $2g$, unless a separate proof of $|f| \leq g$ is supplied.

A summary comparison:

Aspect	This development	CoRN
Reals	presented regular sequences	abstract <code>ConstructiveReals</code>
Integration space	Definition 2.1, no normalization	abstract <code>IntegrationSpace</code> , unit of integral 1
Convergence API	Type layers + Prop surface	Type-valued <code>CvMeasure</code>
Domination	per-index full set	global pointwise
Countable avoidance	nested intervals over presentations	interval cover of controlled measure
Audit output	<code>[propext, Quot.sound]</code>	<code>Closed under the global context</code>
Comparison caveat	not strictly ordered	not strictly ordered

The apparent asymmetry of the audit outputs—`[propext, Quot.sound]` on the Lean side against `Closed under the global context` on the Rocq side—should not be read as a comparison of axiom counts. As Carneiro’s axiomatization of Lean records [32], Lean builds quotient types into the kernel, with definitional computation rules and the axiom `Quot.sound`, and derives function extensionality from quotients together with propositional extensionality; Coq does not support quotient types, and its ecosystem compensates with setoids. At least one of the two Lean axioms thus pays, as an axiom, for what the Coq side obtains from setoid machinery and quantified interfaces. The shape of the audit output reflects a difference in foundational design, not a difference in mathematical strength.

Accordingly, the present work does not claim to be the first proof-assistant formalization, the first constructive formalization, or the first choice-free formalization of Bishop–Cheng dominated convergence. To our knowledge, within Lean 4, this is the first choice-free formalization of the Bishop–Cheng dominated convergence theorem (mathlib’s dominated convergence theorem is classical). This statement is not based on a systematic literature search; the main contribution is not priority. The two published surveys consulted above are consistent with it without establishing it: Miyamoto’s list of Bishop-style formalizations records Nuprl, CoRN and Minlog and does not include Bishop–Cheng measure theory [15, §27.3.4], and Bickford, reporting one of the most substantial of those developments, states that the general definite integral was the one part of his programme not carried through [16, §28.14]. The contribution is a Lean 4 realization over presented reals, Type-valued global and full-set theorem routes, automatic smooth-data wrappers, a Prop-facing full-set theorem, a Lean implementation of countable avoidance via Bishop’s nested-interval construction, and a reproducible Lean audit whose named public declarations have declaration-level axiom output confined to `[propext, Quot.sound]`.

10 Choice-free audit

The development declares `mathlib` [34] as a dependency, and `mathlib` modules do occur in the transitive import closure of the public roots. Importing is not the same as depending on a result: Lean’s `#print axioms` follows the dependencies actually used by a declaration rather than every axiom made available by its imports. The real-number and measure-theoretic layers used above are constructed inside the artifact; in particular, the audited declarations do not consume `mathlib`’s `Real` or `MeasureTheory` hierarchy. This source-level separation, together with the declaration-level audit, explains why the reported axiom output is compatible with the presence of `mathlib` in the import closure.

The claim is restricted to the public theorem aliases, the named implementation declarations, and the public transitive import closure of `choicelfree-bc-dct`; it is not a claim about any file outside that closure. The build script prints axioms for the public declarations and runs a static audit after stripping Lean comments. `Quot.sound` is expected from the quotient constructions described in Section 7—the presented rational layer and the transport quotient of the presented reals; quotient extraction and classical selector axioms are not expected in the audited declarations.

What the claim is relative to. Collecting the qualifications made above, the audited statement is: for every type X and every integration space S on X in the sense of Definition 2.1, the named public declarations hold, and their proof terms use no axioms beyond `propext` and `Quot.sound`. The quantified structure S plays the same role that `IntegrationSpace` plays in CoRN, and the reals are the concrete presented reals of Section 2 rather than an abstract real-number interface. The countable-avoidance step required by the profile route is discharged inside this audit, not assumed (Section 8). What is *not* claimed is that Definition 2.1 is itself a rendering of Bishop and Cheng’s Definition 1.1 clause for clause (a clause-by-clause transcription is supplied separately, together with a hypothesis-free adapter deriving every field of Definition 2.1 from it—Remark 2.2), nor a model over a continuum, nor one realizing arbitrary positive weights, nor a predicative development, nor any statement about real-number objects other than the presented reals.

The CoRN assumptions audit and the Lean axiom audit are different mechanisms in different proof assistants. The CoRN audit reports `Closed under the global context` for the inspected declarations. The Lean audit reports `[propext, Quot.sound]` for the named public declarations. Both statements are relative to the explicit theorem interfaces being audited.

Three complementary checks

The audit is deliberately not based on a single text search.

- (1) **Kernel dependency output.** Dedicated Lean files invoke `#print axioms` on the public aliases and the implementation declarations used by them. The reported axiom names are confined to the set `{propext, Quot.sound}`, and the principal public DCT declarations report both. In particular, the fact that an alias is marked `noncomputable` is not treated as evidence for or against choice; the criterion is the dependency output of its elaborated term.
- (2) **Transitive source-closure scan.** `tools/static_no_choice_audit.py` computes the import closure of six public roots, strips Lean comments, and rejects forbidden tokens in the resulting source: the selectors (`Classical.choice`, `Classical.choose`, any `Classical.`, `native_decide`, `Quot.out`, `Quotient.out`, `open Classical` and the `classical` tactic), the placeholders (`sorry`, `admit`), the escapes from kernel checking (`unsafe`, a declaration-initial axiom, `partial def`, `implemented.by`, `extern`, `debug.skipKernelTC`), and a conclusion of

True. Seven were added in v0.4.2—the declaration-initial `axiom`, `Quotient.out`, `partial def`, `implemented_by`, `extern`, `debug.skipKernelTC`, and the `True` conclusion; `unsafe` was scanned from the outset. Each reports zero on the closure. The recorded run traverses 513 Lean files and passes (`closure_files: 513, tracked_lean_files: 513, STATIC AUDIT PASSED`). The audited closure and the Lean paths recorded in `SHA256SUMS` are compared as sets in both directions, and the run fails if they differ, or if an import under a project-local prefix is absent from the tree, or if an import is neither project-local nor on the allowed-external list: a missing file is a packaging error and must not be read as an external dependency, which would silently shrink the closure being audited. Because the audit roots include the aggregation module `Mathdemo`, this closure covers the four reading-layer modules: the mathematical reading interface `Mathdemo.MathematicalInterface`, the Definition 1.1 transcription `Mathdemo.SourceIntegrationSpaceDef11`, and the two concrete modules `Mathdemo.DiracIntegrationSpace` and `Mathdemo.FiniteWeightedIntegrationSpace`. The scan complements the declaration-level check by detecting suspicious source in the public closure even when it is not reached by one selected theorem.

- (3) **Integrity and cross-system provenance.** Root checksums cover the tracked artifact, while a separate checksum file protects the reproduced CoRN audit package. The CoRN commands pin both the source commit and Rocq version. Thus the Lean claim and the prior-work comparison can be reproduced independently rather than inferred from prose in this paper.

The checks answer different questions. A source scan cannot replace kernel dependency analysis, because the presence of an imported classical declaration does not show that a proof term uses it. Conversely, a short list of printed axioms does not by itself establish that the intended source tree, versions, and audit inputs were the ones inspected. The artifact therefore reports both kinds of evidence and protects their inputs with checksums.

Remark 10.1 (What the axiom footprint means mathematically). The audit output—the axioms `propext` and `Quot.sound`—can be read against Carneiro’s axiomatization of Lean’s type theory [32]. Each named declaration whose axiom footprint is reported in this paper is a theorem of a dependent type theory with inductive types, a hierarchy of universes, and a definitionally proof-irrelevant impredicative universe of propositions, extended by exactly two axioms: propositional extensionality (propositions that imply each other are equal) and the soundness axiom of quotient types. Function extensionality is derivable from these two, so in a mathematician’s vocabulary the axioms used amount to: extensionality of propositions, existence of quotients, and, as a consequence, extensionality of functions. In Lean the law of excluded middle is a *theorem* derived from `Classical.choice`, not an independent axiom; not using choice therefore entails that excluded middle is not used as an axiom either.

Set-theoretically, in the model underlying Carneiro’s soundness theorem—universes interpreted as V_{κ_n} for inaccessible cardinals κ_n , propositions as the two-valued proof-irrelevant universe $\{\emptyset, \{\bullet\}\}$ —the only interpretation clause that references the fixed choice function on the interpreting set is the clause for `Classical.choice`. The interpretation of the theorems audited here therefore does not depend on the selection of a choice function, and their consistency is relative to ZFC together with finitely many inaccessible cardinals. Carneiro’s published result is stated for each finite n : the fragment with n universes is interpreted relative to ZFC plus n inaccessibles. A fixed finite development such as this artifact uses only finitely many universe levels, so a concrete finite bound on its universe usage is to be expected; we have not computed one here.

In particular, unconditional extraction of data from a `Prop`-valued existential is exactly what `Classical.choice` licenses; an axiom output without it therefore certifies the absence of any such

extraction from the declarations whose axioms are printed, together with everything their proof terms depend on. That is a smaller set than the closure: the shipped log carries axiom output for 1,573 distinct names, while the closure declares 3,948. What covers the remaining declarations is the token-level scan of Section 12, which is a different and weaker instrument. What is *not* claimed: this reading does not establish that the theorems of the development hold in ZF. Carneiro’s soundness proof is itself carried out in classical set theory with choice (ZFC with inaccessibles); what is established is the object-level absence of choice and the independence of the interpretation from the choice function. The ambient type theory also has an impredicative **Prop** and definitional proof irrelevance with uniqueness of identity proofs built in, so it differs both from Bishop’s informal practice and from homotopy type theory with univalence, with which UIP is incompatible. The constructive content of the development—moduli, explicit exception sequences, witnesses as data—is carried by the design of the development over presented reals, not by the axiom system.

This remark is complementary to Section 8, and the final observation is a heuristic picture rather than a theorem. Section 8 explains why the general and abstract forms of the statements are not provable in the choice-free setting (in Spitters’ sheaf model, sequence avoidance fails); the present remark says inside which axiom system the theorems proved here do hold. Carneiro’s construction provides a classical two-valued interpretation of the audited Lean fragment, in which the reals are uncountable and the general forms are true, while the non-classical models on the no-go side explain why unrestricted formulations cannot be expected in weaker constructive settings. The contrast between the two model families makes the shape of the theorem—presented reals, coded profiles, explicit exception sequences—look mathematically motivated rather than stylistic. We emphasize that no published result establishes that Spitters’ model interprets Lean’s full type theory (impredicative **Prop**, definitional proof irrelevance, quotients), which is why this paragraph is offered as a picture and not as a proof.

11 The size of the development

The artifact shipped with this paper is the code the theorems of this paper depend on: 513 files, 99,988 lines and 3,948 declarations, against the 49 numbered items (definitions, lemmas, propositions, corollaries, theorems) that the source states from Definition 1.1 through Theorem 4.15.

Two remarks on the counting. A declaration is a source line matching

```
^(private )?(noncomputable )?(theorem|def|lemma|structure|abbrev|instance)
```

so declarations carrying other modifiers, or not starting at the beginning of a line, are missed; and the pattern is purely lexical, so it also matches the occasional line inside a block comment. The figures are the raw counts.

11.1 What fixes the contents

The contents were fixed mechanically rather than by judgement. The starting point is reachability: seed with every declaration of the modules that carry a claim of this paper—the public aliases of Section 7, the reading interface, the transcription of Definition 1.1 with its adapter, the three concrete models, and the axiom-check modules—and take the transitive closure of the constants occurring in the type and in the value of each declaration.

That closure is not by itself a tree that compiles, and the gap is not small. Of the 20,817 declarations of the development it selects 2,542, and a further 2,859 have to be added before the type checker accepts the result: material reached through elaboration rather than through a proof

term—instances, `simp` lemmas, notation—and declarations that the audit modules name in `#check` and `#print axioms` commands, which are not proof terms at all. The contents are therefore the least set containing the closure that the compiler accepts, reached by removing the complement, compiling, and restoring what the failure names, until the build succeeds. Every declaration kept is connected, through some chain of references, to a theorem this paper states.

Being connected is weaker than being needed, so the result is not claimed to be minimal: a proof term can mention a lemma the proof would go through without, and reachability keeps it. What is claimed is checked. Two invariants are machine-checked and shipped: the elaborated type and the kernel axiom footprint of each of the 21 public aliases (`tools/public_surface_check.lean`, recorded in `logs/public_surface.txt`), and the presence of every identifier this paper cites by name (`tools/check_frozen_names.sh`, against the 114 names of `tools/frozen_names.txt`). Both hold of the shipped tree, and both are the invariants any later reorganization of the artifact must preserve.

11.2 What the bulk of the development consists of

The formal-to-informal difference is not uniform overhead; at least three kinds are mixed. First, notational overhead: rewritings that a person performs in one line, placed as explicit lemmas. This part is genuinely redundant. Second, material the source presupposes: the foundational layer, which is not a difference from the source but a difference of reference target. Third, constructive content that the source suppresses: moduli of convergence, domain management for partial functions, apartness data. Where Bishop and Cheng write “the series converges”, the formalization must carry the rate. Only the third kind is mathematically interesting, and separating the three cleanly remains open. The second kind, however, can be decomposed against the reference itself.

Compared clause by clause with Chapter 2 of *Foundations of Constructive Analysis*, the foundational layer consists of a transcription of that chapter—regular sequences and Bishop equality with the carrier left unquotiented, the standard bound, the index-shifted arithmetic operations and their laws, the one-point positivity condition of its Definition 3 and the tail form of its Lemma 2, the two orders, and limits with moduli—together with five kinds of newly paid cost: (i) the construction of the rational layer itself, as a decidable ordered field, which is the reference of the reference; (ii) the substitution of the dyadic gauge 2^{-k} for the $1/n$ gauge, translated in both directions by the reindexing $x'_k := x_{2^k}$ (inverse $k(n) := \lceil \log_2 n \rceil$), with every margin constant redesigned under the new gauge; (iii) the discharge, at every operation and predicate, of respect for Bishop equality—the *Foundations* disposes of this in a single remark, that the map from a real to its n -th rational approximation “is not a function”; (iv) the carrying and recovery of existential witnesses—the *Foundations* declares once that in all its existence proofs the witness is understood to have been extracted, and that for positivity “the n was implicitly there all along”; Appendix B.12 is the machine-checked statement that this looseness is harmless without choice; and (v) an explicit modulus for every convergence statement. In one sentence: *most of the foundational stratum is the cost of discharging, at every point of use, conventions that Bishop declared once in prose—witnesses count as extracted, representation-level operations are not functions, a real is not an equivalence class.*

The comparison with Séméria closes the accounting. His CoRN development quantifies over an abstract `ConstructiveReals` interface whose implementations and theory live in the Coq standard library—the subject of his JFLA paper [28]—so the counterpart of that layer sits *outside* his artifact, while `Print Assumptions` still descends into it and reports no axioms. Bishop and Cheng refer to *Foundations*; Séméria refers to the standard library; the present development carries its own. That layer is the cost of internalizing the reference. (These three are all on the representation side of

the divide; the fourth coordinate, reaching the same proposition/data tension from the extensional side, is Booi’s locator theory—Appendix B.13.)

12 Limitations and validation boundary

The result should be read with the following limitations in view.

- (1) **Integration-space fidelity.** Definition 2.1 differs from the source’s Definition 1.1 by the absence of the inhabitedness and normalization requirements—a pure weakening, in the direction that widens the range of application: (Remark 2.2) every integration space in the sense of the displayed Definition 1.1 satisfies Definition 2.1, and the theorems apply to it. The adapter from the clause-by-clause encoded transcription is hypothesis-free, and `Mathdemo.SourceIntegrationSpaceDef11` derives all fourteen fields of Definition 2.1 from Definition 1.1 (Remark 2.2). We note also that the field `I_nonneg` of Definition 2.1 is derivable from the continuity clause, a derivation machine-checked in the artifact, so that carrying it as a field is unnecessary.
- (2) **Concrete semantics.** Three concrete examples are shipped: the degenerate zero-integral model on `PUnit`, which verifies that the public wrapper can be applied end to end; the normalized point-evaluation model `Mathdemo.DiracIntegrationSpace` (Remark 2.2); and the finitely supported model `Mathdemo.FiniteWeightedIntegrationSpace`, whose two-point normalized instance is unconditional. The weights it realizes are those of the form k/m obtained by repeating a point k times among m slots, not arbitrary positive reals, and no model over a continuum is constructed. What the three certify is that the interface is satisfiable, that normalization is realizable, that the adapter applies unconditionally, and that the axioms admit a model in which the measure is not concentrated at a single point; they are not offered as a survey of expressive range.
- (3) **Hypothesis on the limit.** The public theorem assumes that the limit f is already an integrable representative. It proves convergence of the integrals under that hypothesis; it does not derive integrability of a merely measurable limit.
- (4) **Logical and real-number scope.** The development uses Lean’s impredicative `Prop` and regular-sequence presented reals. It is neither a predicative reconstruction nor a theorem over arbitrary constructive real-number objects, and it does not overturn the model-theoretic no-go results discussed in Section 8.
- (5) **Comparison and priority.** Séméria’s CoRN development already contains a Bishop–Cheng dominated-convergence theorem, and the two theorem surfaces live over different real-number and integration-space interfaces. No claim of priority or strict formal generalization is made.
- (6) **Implementation maturity.** The artifact is fixed by reachability from the public claims (Section 11.1) and grouped by the layer each module builds. That criterion keeps what the theorems depend on; it does not produce a library with a designed interface, and the implementation modules beneath the public surface are not a public API. The public aliases and principal proof route have been reviewed and audited, while detailed manual review of every low-level internal declaration remains ongoing. A declaration-level scan of the closure for declarations whose conclusion or body is `True` now returns none. Five were present through v0.4.1 and were removed in v0.4.2: `ContinuousOn`, `lemma33_lt_of_not_le`, `lemma34_out_exists_cell`, `fatou_type_stub_not_source_4_14`, and—the one whose name most invites

misreading—`thm_4_15_dominated_convergence`, a `def` carrying the hypotheses of dominated convergence and concluding `True`. All five were vestiges of earlier generic routes, none was referenced anywhere in the closure, and a declaration whose conclusion is `True` carries no information, so none could contribute to any proof; the theorem of this paper is `bishop_cheng_thm_4_15_propC`, which is unrelated to that vestige and is stated and proved in full. They are recorded here because at the time they were present the token-level scan of the previous item detected vacuous *proofs* (`sorry`, `admit`) and not vacuous *statements*. That scan now also rejects a conclusion of `True`, which is the form all five took. It is a lexical guard, not a complete check: it matches the pattern `: True :=` on a single line, so a vacuous statement whose type is written across lines, or whose conclusion merely reduces to `True`, would escape it. A complete check would have to enumerate the elaborated declarations of the environment and test each conclusion for reducibility to `True`; that is not implemented here.

A complete `./build_audit.sh` run from the `v0.6.0` tree finished on 2026-09-02 with `BUILD_AUDIT_EXIT=0`. The shipped log carries the banner `artifact_version=0.6.0` and records the successful build stages (2856, 2857, 2858, 3 and 2870 jobs), the declaration-level axiom checks reporting `[proptext, Quot.sound]`, the reading-layer axiom check over 94 declarations, and the static audit traversing its closure within the same run. All four shipped logs record that run; none of them names a path from before the implementation modules were regrouped into `Mathdemo/Internal/Rat, Real, Measure and Sec4`.

The scope of the build audit should be delimited. The three reading-layer modules—`Mathdemo.MathematicalInterface`, `Mathdemo.SourceIntegrationSpaceDef11`, and `Mathdemo.DiracIntegrationSpace`—are not in the dependency cone of `Mathdemo.CheckSec3PortAxioms`: they are a layer for *reading* the development, not a layer the development *depends on*, so they are not built in the initial theorem-root stages; they are built by the later aggregate stage. They are audited by two other routes. First, the static source closure of Section 10 covers them, since the audit roots include the aggregation module. Second, a separate aggregated build, `lake build Mathdemo`, completes, and a dedicated reading-layer audit covers all 94 declarations—including the hypothesis-free adapter `toIntSpaceC` and the gauge bridge—of which 93 report `[proptext, Quot.sound]` while `BishopCheng.ComplementedSet` reports no axioms at all. As of `v0.4.1` this aggregate build is part of `build_audit.sh`; the reading-layer axiom audit is run from a check file outside the tracked closure, so that the audited closure is unchanged by the check itself, and its output is shipped (Section 13).

The artifact is tagged `v0.6.0`. The requirement that the tagged release receive a complete `./build_audit.sh` run is met by the run from the release tree just described. The software deposit [37], concept DOI 10.5281/zenodo.21850965, which resolves to the latest version, carries the Lean code and the audit apparatus only; the paper source is excluded from it by `.gitattributes` and is not part of the deposit.

Use of AI-assisted tools

Generative AI and AI-assisted coding tools were used extensively in the development, refactoring, documentation, and review of the Lean source code, and in drafting and revising the prose of this paper. The author determined the mathematical scope and formalization goals, reviewed the public theorem statements and proof architecture, reviewed and approved the manuscript text, and executed the reported build, static, checksum, and axiom audits. The author assumes full

responsibility for the contents of the paper and artifact. Detailed manual review of the low-level implementation is ongoing. AI systems are not listed as authors.

13 Artifact interface and reproducibility

The paper cites the public aliases in `ChoiceFreeMeasureDCTPublic.lean`. That file is the whole of the intended interface: 21 declarations in the namespace `ChoiceFreeMeasureDCT`, each carrying a doc-string that states what it is. The supplementary file `SupplementChoiceFreeMeasureDCT.lean` checks and prints axioms for both the aliases and their implementation declarations. The artifact is [37].

A reader can inspect the result at three levels.

- (1) **Public surface.** Begin with the following two files:

```
ChoiceFreeMeasureDCTPublic.lean
Mathdemo/ChoiceFreeDCTExamples.lean.
```

They display the callable interfaces without requiring navigation through the internal iteration history.

- (2) **Short checks.** Run the static audit, the root checksum command, and then the independent checksum command inside `audits/corn/`:

```
python3 tools/static_no_choice_audit.py
sha256sum -c SHA256SUMS
(cd audits/corn && sha256sum -c SHA256SUMS)
```

These checks do not rebuild Lean but verify the public source closure and the integrity of the shipped evidence.

- (3) **Complete Lean verification.** Run `./build_audit.sh`. The script builds the principal theorem root, public support modules, the concrete example, and the public alias files; invokes the dedicated axiom checks; and finishes with the strengthened static audit. A successful run ends with `BUILD_AUDIT_EXIT=0`. The script also performs the aggregate `lake build Mathdemo`, the second of the two audit routes described in Section 12, whose record is shipped as `logs/mathdemo_build.txt`; it then generates a temporary reading-layer check file outside the tracked closure, runs `lake env lean` on it to print the axiom footprint of all 94 reading-layer declarations—the facade’s 36, the transcription’s 27 including the hypothesis-free adapter and the gauge bridge, the point-evaluation model’s 10 and the finitely supported model’s 21—and verifies that 93 report `[propext, Quot.sound]` and one reports no axioms, shipping the output as `logs/reading_layer_axioms.txt`.

Local reruns are written to:

```
logs/build_audit.rerun.txt
logs/static_audit.rerun.txt.
```

The shipped reference logs are kept stable so that checksum verification remains meaningful after rerunning the audit. The independently packaged CoRN reproduction is documented in `audits/corn/CORN_ASSUMPTIONS_AUDIT.md`; it can be checked without trusting the Lean build or the comparison prose in this paper.

14 Conclusion

The artifact demonstrates a complete profile-based route from convergence in measure and full-set domination to convergence of integrals over presented reals, relative to the explicit integration-space interface of Definition 2.1. Its central constructive step is not the suppression of the exceptional levels returned by profile theory, but their explicit enumeration followed by a presented-real countable-avoidance construction. That distinction explains both how the formal proof proceeds and why it does not contradict the known obstruction for unrestricted constructive real-number objects.

The result is exposed through proof-relevant global and full-set interfaces, automatically synthesized wrappers, and a Prop-facing interface, and the final public declarations are accompanied by kernel dependency output, a transitive source audit, checksums, and a pinned reproduction of the closest CoRN prior work. The limitations are equally explicit: the limit is assumed integrable; the shipped concrete models reach finitely supported measures with weights of the form k/m , and no model over a continuum is constructed; the integration-space interface is a clause-level strict weakening of the encoded Bishop–Cheng Definition 1.1 (as of v0.4.1 every one of its fields is derived, hypothesis-free, from a clause-by-clause transcription that states clause (4) at the memoir’s own gauge and fixes the development’s ambient encodings; the remaining clause-level differences are the omitted inhabitedness and normalization requirements—Remark 2.2); and the development is not predicative. These boundaries leave clear next tasks while keeping the present theorem and its audit precisely scoped.

A From the mathematics to the Lean declarations

The tables in this appendix let a reader move between Part I, the source, and the artifact. They are the only place in which the correspondence is asserted; the statements in Part I are self-contained mathematics, and the declarations in Part II are self-contained Lean.

A.1 Numbering: this paper and the source

This paper numbers items by section, so its numbers do not coincide with those of Bishop and Cheng’s memoir; since both schemes contain items numbered “3. x ” and “4. x ”, the correspondence is collected here. The headings in Part I also carry the source numbers.

Notes in the last column assert agreement of mathematical content with the memoir; they do not assert word-for-word agreement with its English text.

This paper	Content	Source	Note
Definition 2.1	integration space (working form)	Definition 1.1	clause-level strict weakening of Definition 1.1: all fourteen fields derived, hypothesis-free, from the transcription at the memoir’s own gauge, with partial functions and the integral on L as the memoir has them (Remark 2.2; Appendix B.6)
Definition 3.1	full set	Definition 1.9	same statement
Definition 3.2	complemented set	Definitions 2.1, 2.2(d)(e), 2.3	same core statement; merges $-A$, $A - B$, integrability, measure
Definition 3.3	convergence in measure	Definition 4.11	same statement
Definition 3.4	domination on full sets	§1 notational convention	the convention with the quantifiers displayed; index-dependent full sets already licensed by it—not a weakening
Definition 3.5	profile	Definition 3.1	stated by the formalization in coded form (Appendix B.4); no continuity clause—a weaker hypothesis, hence more general
Definition 3.6	smooth level	—	introduced in the source inside the text of its Theorem 3.5
Theorem 4.1	dominated convergence	Theorem 4.15	same assertion; hypotheses phrased differently (Remark 4.3, Section 6)
Theorem 5.1	smoothness away from an explicit sequence	Theorem 3.5	the formalized statement returns the explicit family constructed by the source’s proof (Appendix B.5); the tail conditions are quantified over $[a, b]$, as the source’s $F \subseteq C[a, b]$ has them; Part I keeps the source’s wording—not a strengthening (Appendix B.5)
Theorem 5.2	integrability of level sets	Theorem 3.6	A pair and B pair constructed independently (Appendix B.8); exported in the source’s global form, one exceptional sequence for all $t > 0$, obtained as the source obtains it, by covering $\mathbb{R}^+$ with the intervals $((n+2)^{-1}, n+2)$ and flattening the per-interval sequences
Lemma 5.3	countable avoidance	—	not a separate stage in the source; contained in the transition from its Theorem 3.5 to its Theorem 3.6 (Appendix B.3)
Proposition 5.4	uniform complement estimate	hypotheses of Lemma 4.14 + construction of Theorem 4.13	returns (A, N, δ) as data; reparameterization $B := -C$ (Appendix B.10)
Proof step (4), Section 6	two-part estimate	Lemma 4.14	same two-part estimate
Appendix B	Lemma 1.2, Corollary 1.3, Definition 7.1, and others	same numbers	the appendix retraces the source’s arguments and cites the source’s numbers directly

A.2 From mathematical objects to Lean declarations

Axiom of Definition 2.1	IntSpaceC field
(A1) respect for Bishop equality	<code>L_resp, I_resp</code>
(A2) closure of L	<code>add_mem, smul_mem, abs_mem, cutPos_mem</code>
(A3) linearity of I	<code>I_add, I_smul</code>
(A4) positivity of I	<code>I_nonneg</code>
(A5) truncation limits	<code>cutNat_tendsto_raw, cutSmall_tendsto_raw</code>
(A6) constructive continuity	<code>continuity</code>

Mathematical object	Public alias	Implementation
Definition 3.3	—	<code>ConvergeInMeasureC</code>
Definition 3.4	—	<code>DominatedOnFullC</code>
Theorem 4.1	<code>bishop_cheng_dominated_convergence_propC</code>	<code>bishop_cheng_thm_4_15_propC</code>
Theorem 5.1	<code>profile_smooth_away_from_sequenceC</code>	<code>thm_3_5_smooth_aeC</code>
Lemma 3.4 (partition data)	<code>profile_partition_dataC</code>	<code>lemma_3_4DataC</code>
Theorem 5.2	<code>profile_level_sets_integrable_apart_globalC</code>	<code>thm_3_6_all_posC</code>
Theorem 3.6, interval-local core	<code>profile_level_sets_integrable_apartC</code>	—
Lemma 5.3	—	<code>Thm36B1ApartPointDataC</code> <code>thm36B1_apartPointDataC</code> <code>thm36B_smoothPointDataC_construct</code>
Proposition 5.4	<code>uniform_complement_from_profile_levelsC</code>	<code>lemma43UniformComplementData_of_majorantC</code> <code>lemma43UniformComplementData_of_majorantOnFullDataC</code>
Proof step (2)	—	<code>lemma43DyadicSmoothDataC_construct</code>
Proof step (4)	—	<code>lemma_4_14_local_two_epsilonC</code>
Proof step (5)	—	<code>integral_diff_lt_of_abs_error_integral_ltC</code>

Public aliases are in the namespace `ChoiceFreeMeasureDCT`; implementation declarations are in `BishopSec1P` or `BishopSec3P`. The complete list of public aliases, including the Type-valued route and the automatic wrappers of Section 7, is in `ChoiceFreeMeasureDCTPublic.lean`.

B Constructions specific to the formalization, and their mathematical status

In the spirit of the blueprint genre, but in the reverse direction: these sections state, as ordinary mathematics, what the formal development had to construct. The appendix has two purposes. The first is to write down, without Lean vocabulary, the constructions that the development did *not* receive from Bishop and Cheng but had to build in the course of the formalization. The second is to make explicit which parts of that work are routine, which are known, and which were the actual difficulties (Appendix B.11). Part I suffices for the mathematical content; this appendix discloses, in mathematical language, what the formalization concretely consisted of.

Throughout the appendix, presented reals are taken in the dyadic-gauge form used by the implementation: regularity reads $|x_m - x_n| \leq 2^{-m} + 2^{-n}$. Section 2.1 stated the $1/n$ form of the

Foundations; the two are translated by an explicit reindexing, and the translation is part of the cost accounted in Section 11.2.

B.1 The proof flow against the source

The conclusion first: the outline of the formal proof coincides with the source’s. The number of stages and their order are the same, and the differences concentrate in how each stage is *stated*. The single exception is countable avoidance, which in the source is not a separate stage at all: it lives inside the two transition sentences at the beginning of the proof of its Theorem 3.6, immediately after Theorem 3.5 is invoked (“all but countably many points t of (a, b) are smooth”; “let t be smooth”). Constructively that transition cannot be executed as written, and the development had to erect it as a theorem of its own (Appendix B.3).

The dependency structure of the proof, with the status of each stage relative to the source, is the following; arrows indicate constructed data or a theorem application, as in Section 6.

- Clauses (1)–(4) of Definition 1.1 $\Rightarrow$ Lemma 1.2 $\Rightarrow$ Corollary 1.3 at $k = 1 \Rightarrow$ positivity of I . Status: as in the source; that positivity is derivable, so that carrying it as an axiom is unnecessary, is an observation of this development (Appendix B.6).
- Coded profile and the partition data of Lemma 3.4 $\Rightarrow$ Theorem 3.5 (smoothness) together with the explicit exception sequence. Status: statements as in the source; the coded form and the sequence-returning form are formalization-specific (Appendices B.4, B.5).
- Presented-real order with cotransitivity as data $\Rightarrow$ countable avoidance by a two-point cut with a margin. Status: the statement is Bishop’s; the two-point cut is where the construction departs from the classical bisection (Appendices B.2, B.3).
- Avoidance applied to the exception sequence $\Rightarrow$ apart thresholds $\Rightarrow$ Theorem 3.6: the A pair and the B pair constructed independently, with the measure equality; the signed double series over a fixed shell enumeration is consumed here (Appendices B.8, B.9).
- Dyadic levels $\alpha_n \in (2^{-(n+1)}, 2^{-n})$ as data (source Lemma 4.3) $\Rightarrow$ uniform complement estimate returning (A, N, δ) (source: hypotheses of Lemma 4.14 and the construction inside Theorem 4.13) $\Rightarrow$ two-part estimate (Lemma 4.14) $\Rightarrow$ Theorem 4.1 = source Theorem 4.15, with full-set domination and the majorant $H = |g| + |f|$ (Appendix B.10).

The differences concentrate at four places.

(a) *From existence to data*. Where the source writes “there exists”, the development frequently writes a function returning the object: clauses (2) and (4) of Definition 1.1, the level sequence of Lemma 4.3, the triple (A, N, δ) in the hypotheses of Lemma 4.14. Under Bishop’s intended BHK reading of $\exists$ this is faithfulness rather than strengthening; as a Lean interface, however, it is a data refinement of the corresponding **Prop**-valued statement (Section 7.1) (Appendix B.12).

(b) *From “countably many” to an explicit sequence*. Theorem 3.5’s “all but countably many” cannot be handed to the avoidance construction as stated; only the form that returns an explicit sequence can be (Appendix B.5).

(c) *From a one-point bisection to a two-point cut*. Inside the avoidance, the classical bisection at a midpoint is undecidable. A margin is introduced, the interval is cut at two points, and the middle is discarded (Appendix B.3).

(d) *The B pair does not follow from the A pair*. Where the source dismisses the B side as essentially similar, the fact that $\leq$ is $\neg <$ forces an independent construction, with the monotonicity and the usable implications reversed (Appendix B.8).

None of the four is a claim that the source’s mathematics is wrong. The source is correct within constructive mathematics, and its omissions are the usual “essentially similar” economies of print. The four points say what is missing if one tries to execute the text directly as a construction, and the sections below supply exactly that.

B.2 The order on presented reals as data

Definition B.1 (Presented real, dyadic form). A *presented real* is a sequence $x = (x_n)_{n \geq 0}$ of rationals such that

$$|x_m - x_n| \leq 2^{-m} + 2^{-n} \quad (\forall m, n)$$

(a *regular sequence*). Two presented reals x, y are *equal* ($x \approx y$) if

$$\forall k \exists N \forall n \geq N : |x_n - y_n| \leq 2^{-k}.$$

A real is the regular sequence itself; no equivalence classes are taken. Below, $\mathbb{R}$ denotes the presented reals in this sense.

Definition B.2 (Order, and its data form). $x < y$ means that there exist k, N with

$$\forall n \geq N : y_n - x_n > 2^{-k}$$

(the development takes the gauge $2^{-(k+1)}$; nothing below depends on that choice). $x \leq y$ means $\neg(y < x)$, and x and y are *apart* when $0 < |x - y|$.

Data form: evidence for $x < y$ is a pair (k, N) together with the proof of its property. “To have $x < y$ as data” means holding this pair, as distinguished from the mere truth of $x < y$. The distinction runs through the entire appendix.

Fact B.3 (Cotransitivity—the constructive pivot). *Suppose $x < y$ is held as data. Then for every $z \in \mathbb{R}$ one can decide which of*

$$x < z \quad \text{or} \quad z < y$$

holds, obtaining evidence for the decided side.

Reason. The evidence (k, N) gives $y_n - x_n > 2^{-k}$ for $n \geq N$. Fix any $n_0 \geq \max(N, k + 4)$ and compare the rational z_{n_0} with $\frac{x_{n_0} + y_{n_0}}{2}$. Rational comparison is decidable, and the comparison selects one of the two cases. At n_0 the gap from the midpoint to either endpoint exceeds $2^{-(k+1)}$. Passing to a later index $n \geq n_0$ moves both members of the selected difference, so the regular-sequence error to subtract is $2(2^{-n} + 2^{-n_0}) \leq 4 \cdot 2^{-n_0} \leq 2^{-(k+2)}$; this still leaves the positive margin $2^{-(k+2)}$, so the selected side actually holds (the regularity error of the running index is absorbed by the eventual quantification, and the implementation realizes this extra unit of precision through the shifted subtraction index; the general mechanism is Appendix B.12). $\square$

The essential point is that *a gap is needed*. If $x < y$ were not held—if x and y might touch—the decision could not be made: for presented reals, the one-point dichotomy “ $z \leq m$ or $m \leq z$ ” is undecidable.

B.3 Countable avoidance by a two-point cut with a margin

This is the most characteristic construction of the development. The goal is the following theorem, which is exactly the assertion of Bishop’s well-known nested-interval method—the *statement* is known. Carrying it out as a construction over presented reals, however, cannot use the classical bisection.

Theorem B.4 (Countable avoidance; = Lemma 5.3). *Suppose $a < b$ is held as data, and let $(s_j)_{j \geq 0}$ be an arbitrary sequence of presented reals. Then one can construct $t \in (a, b)$ apart from every s_j :*

$$a < t < b, \quad 0 < |t - s_j| \quad (\forall j).$$

No choice principle is used.

Why the classical bisection fails. Classically one takes the midpoint $m = \frac{l+r}{2}$ of the current interval $[l, r]$ and keeps the right half if $s_j \leq m$, the left half otherwise. But as noted after Fact B.3, the position of s_j relative to the *single* point m cannot be decided, so this branch cannot be constructed.

The resolution: cut at two points and discard the middle. With two points separated by a gap, Fact B.3 applies as it stands.

Definition B.5 (Margin and cut points). For $d \in \mathbb{R}$ set the *margin*

$$\sigma(d) := \frac{1}{4} d$$

(the development implements it by halving twice). What is required of σ is exactly two properties: $0 < \sigma(w)$ and $2\sigma(w) < w$; the choice $\sigma(w) = \frac{1}{4}w$ is merely the simplest that satisfies them. For an interval $[l, r]$ with $l < r$ and width $w := r - l$, the *cut points* are

$$q_L := l + \sigma(w), \quad q_R := r - \sigma(w).$$

Lemma B.6. *If $l < r$ is held as data, then*

$$l < q_L < q_R < r, \quad q_R - q_L = \frac{1}{2} w,$$

and in particular $q_L < q_R$ is held as data.

Proof. From $w > 0$ we get $\sigma(w) = \frac{1}{4}w > 0$, hence $l < l + \sigma(w) = q_L$ and symmetrically $q_R < r$. Moreover

$$q_R - q_L = (r - \sigma(w)) - (l + \sigma(w)) = w - 2\sigma(w) = w - \frac{1}{2}w = \frac{1}{2}w > 0. \quad \square$$

Definition B.7 (Interval data). Fix $a < b$ and the sequence (s_j) . *Interval data at stage n* is a pair (l, r) of presented reals satisfying, with evidence, the four conditions: (i) $l < r$; (ii) $a < l$ and $r < b$; (iii) $r - l \leq (b - a) 2^{-n}$; (iv) for every $j < n$: $s_j < l$ or $r < s_j$. Condition (iv) is read “ s_j is avoided”.

Lemma B.8 (One step). *From interval data $I = (l, r)$ at stage n one can construct interval data $J = (l', r')$ at stage $n + 1$ with $l \leq l'$ and $r' \leq r$.*

Proof. Let $w := r - l$ and take q_L, q_R as in Definition B.5. By Lemma B.6, $q_L < q_R$ is held as data. Apply Fact B.3 to this pair and $z := s_n$, and distinguish the two cases it returns.

Case 1: $q_L < s_n$. Put $J := (l, q_L)$. (i) $l < q_L$ is Lemma B.6. (ii) $a < l$ by hypothesis, and $q_L < r < b$. (iii) The width is

$$q_L - l = \sigma(w) = \frac{1}{4}w \leq \frac{1}{4}(b - a)2^{-n} \leq (b - a)2^{-(n+1)}.$$

(iv) For $j < n$, avoidance is preserved because $J \subseteq I$ ($s_j < l = l'$ or $r' \leq r < s_j$). For $j = n$, the case hypothesis $q_L < s_n$, i.e. $r' < s_n$, is precisely the avoidance.

Case 2: $s_n < q_R$. Put $J := (q_R, r)$. Conditions (i)–(iii) follow symmetrically, the width being $r - q_R = \sigma(w)$. Condition (iv) for $j < n$ is as before, and for $j = n$ the case hypothesis $s_n < q_R = l'$ is the avoidance. $\square$

Remark B.9 (Discarding the middle). The surviving interval has $\frac{1}{4}$ of the original width, smaller than the $\frac{1}{2}$ of the classical bisection. This is not a benefit of faster convergence; it is the consequence of being forced to *abandon the central half*. The classical argument exhausts the interval by the two-way split “left or right”; constructively that split is undecidable, and only after surrendering the region between the two cut points does a decidable two-way split appear. This operation—replacing an undecidable dichotomy by a decidable one purchased with a margin—recurs throughout the development (Appendix B.12).

Proof of Theorem B.4. For the initial interval, apply the cut of Definition B.5 to (a, b) itself:

$$I_0 := (a + \sigma(b - a), b - \sigma(b - a))$$

(the artifact makes the same choice). By Lemma B.6, $a < \text{left endpoint} < \text{right endpoint} < b$, and the width is $\frac{1}{2}(b - a) \leq (b - a) \cdot 2^0$, so condition (iii) holds at $n = 0$; condition (iv) at $n = 0$ is vacuous. Iterating Lemma B.8 yields interval data $I_n = (l_n, r_n)$ for every n . By construction (l_n) is nondecreasing, (r_n) is nonincreasing, and

$$0 < r_n - l_n \leq (b - a) 2^{-n} \longrightarrow 0.$$

For $m \geq n$ we have $l_n \leq l_m \leq r_m \leq r_n$, hence $|l_m - l_n| \leq (b - a) 2^{-n}$: the left endpoints form a Cauchy sequence with an *explicit modulus*. The coefficient $b - a$ is itself a presented real; taking a rational Archimedean bound $b - a \leq K$, accuracy 2^{-k} is reached from index $n \geq k + \lceil \log_2 K \rceil$ on, a numerical modulus. Completeness of the presented reals (taking a diagonal sequence) constructs the limit $t := \lim_n l_n$, with $l_n \leq t \leq r_n$ for all n .

From $a < l_0 \leq t$ and $t \leq r_0 < b$ we get $a < t < b$ (composing a strict with a weak inequality is the standard transitivity lemma, itself a consequence of cotransitivity). For each j , condition (iv) of I_{j+1} gives $s_j < l_{j+1}$ or $r_{j+1} < s_j$; combined with $l_{j+1} \leq t \leq r_{j+1}$ this yields $s_j < t$ or $t < s_j$, that is, $0 < |t - s_j|$. $\square$

B.4 Coded profiles

Definition 3.5 presents a profile as a pair: a family F of ramp functions and a monotone functional λ on it. Constructively, the problem is that being handed an element of F “as a function” does not let one extract the witnesses of its behaviour that the proofs need.

Definition B.10 (Coded profile). A *coded profile* on $[a, b]$ consists of: a type C of *codes* with an interpretation $\text{ev} : C \times [a, b] \rightarrow \mathbb{R}$ and a subfamily $F \subseteq C$; boundedness, $0 \leq \text{ev}(f, x) \leq 1$ for $f \in F$ and $a \leq x \leq b$; codes for the two constants, with $\text{ev}(0, \cdot) \equiv 0$ and $\text{ev}(1, \cdot) \equiv 1$; *separation*: given $a \leq u$, $0 < v - u$ as data, and $v \leq b$, one can *construct* a code $f \in F$ whose interpretation is 0 on $[a, u]$ and 1 on $[v, b]$; and a value functional $\lambda : C \rightarrow \mathbb{R}$, monotone along pointwise comparison of interpretations.

The decisive point is that the separation clause reads “one can construct”—it *returns* a code. If F were an extensional set of functions, obtaining an actual separating function from its mere existence would require a choice principle; making codes first-class objects and defining λ on codes removes that extraction from the outset. Where the source silently “takes a ramp function”, the development holds a function that returns a code. The same shape—carry presentations, define functionals on presentations, never extract witnesses from extensional objects—is reusable at completions, at simple-function representations, and in the passage from measures to integrals.

B.5 The explicit exception sequence

Theorem B.11 (Explicit-sequence form of the smoothness theorem). *For an integrable h and an interval (a, b) , one constructs an explicit sequence $T = (T_n)_{n \geq 0}$ of presented reals such that every $t \in (a, b)$ apart from every T_n is a smooth level for the profile attached to h .*

Only in this form can Theorem B.4 be applied, with $(s_j) := T$. Two clarifications keep the claim exact. First, stating “the exceptions are countable” as “the exceptions are *this sequence*” is *not* a mathematical strengthening of the source: the source’s proof of its Theorem 3.5 itself constructs the finite stagewise families $\{t_1^k, \dots, t_{n(k)}^k\}$ and their union T . The work of the formalization is flattening the proof-internal data into a first-class $\mathbb{N}$ -indexed sequence, fixed as an interface that the avoidance theorem consumes directly. Second, the reason this does not collide with the no-go results is the confluence stated in Section 8, not the explicit-sequence form alone: over an abstract real-number object, a principle “from an explicit sequence, return a real apart from all its terms” would itself fall under the no-go; the avoidance construction of Appendix B.3 runs over presented-real order data, and that restriction is what carries the argument.

B.6 The truncation clause of Definition 1.1

The truncation demanded by clause (1) of the source’s Definition 1.1 is $\min\{f, 1\} \in L$ *only*; the source’s own remark recovers positive-integer truncation via $\min\{f, n\} = n \cdot \min\{n^{-1}f, 1\}$. The following four statements delimit what can and cannot be recovered.

Proposition B.12 (Positive constants). *If $0 < \alpha$ is held as data, then clause (1) yields $\min\{f, \alpha\} \in L$: indeed $\min\{f, \alpha\} = \alpha \cdot \min\{\alpha^{-1}f, 1\}$, and the right-hand side lies in L by $\min\{\cdot, 1\}$ and closure under scalars. Constructing α^{-1} requires evidence that α is apart from 0—exactly “ $0 < \alpha$ as data” in the sense of Definition B.2.*

Proposition B.13 ($\alpha = 0$). *Clause (1) yields $\min\{f, 0\} \in L$: indeed $\min\{f, 0\} = \frac{1}{2}(f - |f|)$, and the right-hand side lies in L by $|\cdot|$ and linear combination. The verification goes through the lattice identity $\min\{a, b\} = \frac{1}{2}(a + b - |a - b|)$, valid on presented reals without case analysis, at $b = 0$; a case split on the sign of $f(x)$ is not decidable, so the identity is the constructively correct route.*

Remark B.14 (What $0 \leq \alpha$ alone does not supply in the source route). Proposition B.12 consumes a positivity witness and Proposition B.13 exact zero; the split between the two cases is undecidable, so no uniform construction from $\neg(\alpha < 0)$ alone is obtained here. Its impossibility is *not* proved (that would require a model separation); the claim is only that the source’s route needs the witness, and that no other constructive route is evident. The source-derivable clause (Remark 2.2) is thus truncation at constants carrying a positivity witness, or at 0—the two kinds of constant the development in fact truncates at. The interface implements exactly this clause: truncation enters as a field only for constants carrying positivity data (`cutPos_mem`), truncation at 0 is the derived lemma of Proposition B.13 (`cutZero_mem`), and the witness type `CutConstWitnessC`, with a positive branch and a zero branch, carries the evidence to every use site.

Proposition B.15 (Negative constants fail in the non-compact canonical model). *For $\alpha < 0$, closure under $\min\{\cdot, \alpha\}$ fails in the non-compact instances of the source’s own canonical model. In the model $(X, C(X), \mu)$ of its Definition 7.1— X locally compact, $C(X)$ the continuous functions of compact support, μ a positive measure—one has $\min\{f, \alpha\} = \alpha \neq 0$ off the support of f , so for non-compact X the truncation has no compact support and leaves $C(X)$. $\square$*

Consequently an interface demanding truncation at arbitrary constants and the source’s Definition 1.1 would not be ordered by logical strength. With the witness-carrying clause (Remark B.14),

Definition 2.1 is strictly weaker, at the clause level, than the encoded Definition 1.1; the present proposition is the design reason for the witness-carrying form.

Proposition B.16 (Positivity of I is derivable). *Clause (2) implies: if $g \in L$ is nonnegative everywhere on its domain, then $I(g) \geq 0$.*

Proof. First, Corollary 1.3 of the source at $k = 1$: if $I(g) > 0$ then $g(x) > 0$ at some point x of its domain. Apply clause (2) to the sequence $z_n := 0 \cdot g$ (each $z_n \in L$ with the domain of g , pointwise nonnegative, $\sum_n I(z_n) = 0 < I(g)$); it returns a point x with $\sum_n z_n(x) = 0 < g(x)$. Now suppose $I(g) < 0$. Then $I(-g) = -I(g) > 0$ and $-g \in L$, so $-g(x) > 0$ at some point, i.e. $g(x) < 0$, contradicting $g \geq 0$. Since $I(g) \geq 0$ abbreviates $\neg(I(g) < 0)$, this is a constructively valid conclusion. $\square$

Hence the positivity axiom (A4) of Definition 2.1 need not be carried as a field; the derivation is machine-checked in the artifact.

B.7 The two limits of the point-evaluation model

Fix a point x_0 , let L be the partial functions defined at x_0 , and let $I(f) := f(x_0)$. Clauses (1)–(3) are immediate—the constant 1 lies in L with $I(1) = 1$, so the model is normalized—and clause (4) reduces to two statements about the single real $c := f(x_0)$.

Theorem B.17. *For every $c \in \mathbb{R}$, with explicit moduli: (a) $\min\{c, n\} \rightarrow c$; indeed with $N := 2^{m(c)}$, where $m(c)$ is the exponent of an Archimedean bound for c , every $n \geq N$ gives $\min\{c, n\} \approx c$ —the modulus is constant, and the convergence is exact from a finite stage on. (b) $\min\{|c|, 2^{-n}\} \rightarrow 0$; for gauge k , every $n \geq k + 2$ gives $|\min\{|c|, 2^{-n}\}| \leq 2^{-n} \leq 2^{-(k+2)} < 2^{-(k+1)}$.*

Proof. (a) Choose a natural number $N \geq c$ (the Archimedean property, in the explicit form $N = 2^m$); for $n \geq N$ we have $c \leq N \leq n$, hence $\min\{c, n\} = c$. (b) $0 \leq \min\{|c|, 2^{-n}\} \leq 2^{-n}$, and $2^{-n} \leq 2^{-(k+2)} < 2^{-(k+1)}$ for $n \geq k + 2$. $\square$

Remark B.18 (Where the proposition/data gap does actual work). For the Archimedean bound in (a), the development already possessed the *proposition* “there exists $N \in \mathbb{N}$ with $c \leq N$ ”. A modulus, however, is data, and cannot be extracted from that proposition; the witness $2^{m(c)}$ inside the existing proof had to be written out as a function. “The existence was known, but a witness was needed” is a concrete work item that prose mathematics does not register; the same phenomenon reappears at the uniform complement estimate (Appendix B.10).

B.8 The A pair and the B pair: two complemented sets at one level, neither yielding the other

The source’s Theorem 3.6 asserts that at a level t two complemented sets,

$$A = (\{h \geq t\}, \{h < t\}), \quad B = (\{h > t\}, \{h \leq t\}),$$

are integrable with the same measure. Classically each pair consists of a set together with its ordinary complement, and at a continuity level the two positive components differ only on the null boundary $\{h = t\}$, so their characteristic functions agree almost everywhere; the two pairs are not obtained from one another by swapping components. Constructively they are not.

Definition B.19 (The four sets). The value of an integrable h at a point x is given as data: the absolute convergence at x of a series representing h , together with the sum $s(x)$. On that basis,

$$A^1 := \{x : t \leq s(x)\}, \quad A^2 := \{x : s(x) < t\}, \quad B^1 := \{x : t < s(x)\}, \quad B^2 := \{x : s(x) \leq t\},$$

each set including the condition that the value exists.

That A is complemented—elements of A^1 and A^2 are distinct—follows because $t \leq s$ and $s < t$ are incompatible: $t \leq s$ is $\neg(s < t)$ by definition, so the incompatibility is definitional and requires no comparison of s with t . Similarly for B .

Remark B.20 (Why B is not a corollary of A). Classically the two components of B are ordinary complements, as are the two components of A ; but B is not obtained from A by swapping components. At a continuity level the difference between $\{h \geq t\}$ and $\{h > t\}$ is the null boundary $\{h = t\}$, and the integrability of B follows from that of A by that classical null-boundary argument. Constructively $\leq$ is the *negation* of $<$, and $\{h \geq t\}$ and $\{h > t\}$ are not each other's negations; passing between them is a double-negation elimination. The classical complement route is therefore closed; and on the data route, the public data of the A pair does not directly yield the data of the B pair either. The B pair was accordingly constructed independently. (Formal non-derivability—that no construction could exist—is not claimed.)

Remark B.21 (Why the mirror is not a copy: approximation from opposite sides). The ramp used to approximate the characteristic functions is

$$\text{ramp}_{u,v}(y) = \max\left(\min((v-u)^{-1}(y-u), 1), 0\right)$$

(0 for $y \leq u$, linear on $[u, v]$, 1 for $y \geq v$; the reciprocal $(v-u)^{-1}$ is constructed from the positivity evidence of $u < v$). The two pairs use it in opposite directions.

For the A pair, the *upper* end is fixed at $v = t$ and the lower end $u_n = t - \text{gap}_n$ rises to t from below. The resulting ramps form a *decreasing* sequence whose limit realizes $\chi_{\{h \geq t\}}$, and the directly usable implication is

$$\text{ramp}_{u_n,t}(y) < 1 \implies y < t.$$

For the B pair, the *lower* end is fixed at $u = t$ and the upper end $v_n = t + \text{gap}_n$ descends to t from above. The ramps form an *increasing* sequence whose limit realizes $\chi_{\{h > t\}}$, and what is usable is the *positive* evidence

$$\text{ramp}_{t,v_n}(y) > 0 \implies y > t,$$

not the implication used on the A side.

The two constructions thus approximate the characteristic function from opposite sides; the monotonicity of the ramp sequences is reversed, and the directly usable implications are different statements. This is why carrying out the B pair is not a renaming exercise: the chain—about 55 lemmas in the artifact—is laid out a second time with the monotonicity inverted, while the generic components, the signed double series and shell enumeration of Appendix B.9, are reused unchanged. On either side, the *strong* (narrow) set is reached by the direct implication and the *weak* (wide) set is supplied as its complemented partner; the strong side is $\{h < t\}$ for A and $\{h > t\}$ for B , and that is where the two constructions trade places.

B.9 Signed double series and the shell enumeration

Countable additivity arguments use totals of doubly indexed families (A_{ij}) both as iterated sums and as sums along a single enumeration. Classically, absolute convergence licenses free rearrangement; constructively, one must *fix* an enumeration before asserting anything.

Definition B.22 (Shell enumeration). Decompose each $m \in \mathbb{N}$ uniquely as $m = N^2 + t$ with $0 \leq t \leq 2N$, and set

$$\text{cell}(m) := \begin{cases} (N, t) & (t \leq N) \\ (t - N - 1, N) & (t > N). \end{cases}$$

Then $\text{cell} : \mathbb{N} \rightarrow \mathbb{N} \times \mathbb{N}$ is a bijection tracing, for $N = 0, 1, 2, \dots$, the L-shaped shells $\{(i, j) : \max(i, j) = N\}$.

Theorem B.23 (Signed double series along a fixed enumeration). *Suppose given as data: (i) for each i , the absolute row sum $R_i := \sum_j |A_{ij}|$ with modulus; (ii) the sum $\sum_i R_i$; (iii) the flattened sum $\Sigma_{\text{flat}} := \sum_k A_{\text{cell}(k)}$. Then the iterated sum $\Sigma_{\text{iter}} := \sum_i (\sum_j A_{ij})$ can be constructed, and $\Sigma_{\text{flat}} = \Sigma_{\text{iter}}$.*

Proof shape. Split $A_{ij} = A_{ij}^+ - A_{ij}^-$ with $A^+ := \max(A, 0)$ and $A^- := -\min(A, 0)$. Since $0 \leq A_{ij}^\pm \leq |A_{ij}|$, the comparison test transfers the summability data of (i)–(iii) to both halves; for nonnegative families the shell sum and the iterated sum agree by monotonicity of partial sums and squeezing; subtracting the two nonnegative results returns the signed case. $\square$

Three features carry the constructive content: the conclusion is an equality for one fixed bijection, not a rearrangement license; absolute convergence enters as modulus-carrying data, not as a bare assertion; and the sign decomposition meshes with the comparison test because max and min are controlled by absolute values.

B.10 The uniform complement estimate as data

The statement is Proposition 5.4; this subsection records why its conclusion is data and how the set A is built.

Why data. Classically “ $\forall \varepsilon \exists A, N, \delta$ ” would suffice. Here the sequel takes the set A in hand and performs the next construction on it, so an existence statement is not enough: the one-line “take such an A ” becomes a function returning the triple (A, N, δ) . This is the same phenomenon as Remark B.18.

How A is built. In each dyadic window $(2^{-(n+1)}, 2^{-n})$, choose a level α_n at which the profile of the majorant H is smooth—this is where the explicit exception sequence (Appendix B.5) and the avoidance construction (Appendix B.3) are consumed. Apply the level-set theorem at each α_n to obtain integrable $A_n = (\{H \geq \alpha_n\}, \{H < \alpha_n\})$ with its measure. Integrability of H controls $\mu(A_n)$, and for n large the integral of H off A_n is small; the full-set bounds $e_n \leq H$ transfer the estimate to the errors *uniformly in n* .

In order of application, the chain is: explicit exception sequence (Appendix B.5) $\rightarrow$ countable avoidance applied to it (Appendix B.3) $\rightarrow$ integrable level sets at apart thresholds (Appendix B.8) $\rightarrow$ dyadic level selection $\rightarrow$ uniform complement estimate (this subsection) $\rightarrow$ Theorem 4.1. If the first link were not constructive, everything downstream would degrade to existence statements.

B.11 Routine, known, and difficult: a classification

What was received, and what is routine:

Item	Status	Ground
The four clauses of Definition 1.1; Lemma 1.2; Corollary 1.3; the arcs of Theorems 3.5, 3.6, 4.15	known (source)	taken from Bishop–Cheng as stated
$\min\{f, n\} = n \cdot \min\{n^{-1}f, 1\}$	known (source)	the source’s own remark after Definition 1.1
The <i>idea</i> of countable avoidance by nested intervals	known (Bishop, standard)	statement and strategy classical
$\min\{f, 0\} = \frac{1}{2}(f - f)$; derivability of the positivity of $\bar{I}$	routine to known	elementary; that the positivity field is unnecessary is an observation of this development
Respect lemmas for Bishop equality; reading $\exists$ as data; convergence with moduli	formalization routine	not new mathematics, but obligatory

The difficulties:

Item	Status	Ground
Two-point cut with a margin (Appendix B.3)	difficulty	a one-point bisection is undecidable; a margin makes cotransitivity applicable, at the price of discarding the middle
Coded profiles (Appendix B.4)	difficulty, reusable	no witness extraction from extensional objects; the separation clause <i>returns</i> a code
Explicit exception sequence (Appendix B.5)	difficulty	“all but countably many” must become “away from this sequence” before avoidance applies
Mirror of the A pair and the B pair (Appendix B.8)	difficulty	$\leq$ is $\neg <$, so B is not a corollary of A ; approximation from opposite sides reverses monotonicity and usable implications
Signed double series with shell enumeration (Appendix B.9)	difficulty, reusable	an equality for a fixed bijection instead of a rearrangement license; absolute convergence as modulus-carrying data
Uniform complement estimate as data (Appendix B.10)	difficulty	the triple (A, N, δ) must be returned, not asserted
One-point certificates and strong gauge extraction (Appendix B.12)	known (Bishop) + difficulty, reusable	the one-point diagonal condition and its tail equivalence are Definition 3 and Lemma 2 of <i>Foundations</i> , Chapter 2; the difficulty is executing the round trip, under the proposition/data distinction, without choice

The observations recorded by this development:

Item	Status	Ground
Analysis of the truncation clause (Appendix B.6)	observation	a positivity witness is needed; the source’s canonical model refutes closure at negative constants; an arbitrary-constant clause would not be ordered against Definition 1.1
Resolution of the interface difference (the adapter)	result of this development	all fourteen fields follow from Definition 1.1 with no extra hypothesis, with clause (4) stated at the memoir’s own gauge and the bridge to the working dyadic gauge machine-checked
The proposition/data gap surfacing as work (Remark B.18)	methodological observation	an existential proposition does not yield a modulus

The rows labelled “difficulty” are difficulties of the *formalization*, not claims of mathematical novelty. The constructions of Appendices B.3, B.4, and B.5 rewrite, into constructively executable form, operations that the source’s proofs perform implicitly; none is a new theorem. What this formalization adds is that carrying them out determined *what is needed* to perform them—margins, codes, explicit sequences, positivity witnesses, one-point certificates. Those five items are what this appendix hopes to communicate to a mathematician.

B.12 One-point certificates and strong gauge extraction: a choice-free round trip between propositions and data

Section 8(i) said that the cotransitivity branch is returned as data from rational approximations; Section 9 said that every passage from propositions to data in the audited chain proceeds by explicit supply of data or by decidable search. This subsection writes out the decidable search. As Section 10 noted, unconditional extraction of data from an existential proposition is, in Lean, the territory of the choice axiom. Nevertheless, for the *strong order* on presented reals, one can pass between proposition and data without choice. This is a kind of statement for which classical published mathematics has no place—a conversion theorem between propositions and constructions—and which the formalization forced into statement form; type-theoretic literature does have statements of the same kind (Appendix B.13).

The one-point condition itself is not an invention of this development. Definition 3 in Chapter 2 of Bishop’s *Foundations* defines positivity of a real by exactly a one-point diagonal condition, $x_n > n^{-1}$ for some n , and its Lemma 2 supplies the equivalence with the tail form; the margin $2N^{-1}$ appearing there corresponds to the $2\varepsilon_n$ below, which is the dyadic-gauge version. What this subsection adds is the reading of that equivalence under the proposition/data distinction, and the verification that the round trip is executable, mechanically and without choice.

Definition B.24 (Tail positivity, in two forms). Write $\varepsilon_k := 2^{-k}$. For a regular sequence $x = (x_n)$ (so $|x_n - x_m| \leq \varepsilon_n + \varepsilon_m$), distinguish two forms of *tail positivity*: the *propositional form*, that there exist k and N such that $\varepsilon_k < x_n$ for all $n \geq N$; and the *data form*, the triple (k, N, proof) itself. The strong order $0 < x$ of Definition B.2 coincides by definition with tail positivity of x . The propositional form contains a universal quantifier inside the existential—each instance is a decidable rational comparison, but there are infinitely many—so it cannot be searched as it stands.

Definition B.25 (Strong one-point certificate: the diagonal gauge). For a regular sequence x put

$$W_x(n) : \Longleftrightarrow 2\varepsilon_n < x_n.$$

This is decidable by a single rational comparison. The point is that the threshold $2\varepsilon_n$ is tied to the gauge of the inspected index n itself: the certificate is *diagonal*.

Lemma B.26 (Self-propagation of the certificate). *If $W_x(n)$ holds, then $\varepsilon_{n+1} < x_M$ for every $M \geq n + 2$; hence $(k, N) := (n + 1, n + 2)$ is tail-positivity data.*

Proof. For $M \geq n + 2$ we have $\varepsilon_M \leq \varepsilon_{n+2}$. Regularity gives $x_M \geq x_n - (\varepsilon_n + \varepsilon_M)$, and the hypothesis gives $x_n > 2\varepsilon_n$, so

$$x_M > 2\varepsilon_n - \varepsilon_n - \varepsilon_M = \varepsilon_n - \varepsilon_M \geq \varepsilon_n - \varepsilon_{n+2} = 3\varepsilon_{n+2} > \varepsilon_{n+1}. \quad \square$$

The margin $2\varepsilon_n$ is engineered so that after regularity consumes its maximal budget $\varepsilon_n + \varepsilon_M$, one finer gauge ε_{n+1} still survives. It is the one-point version of the same idea as the σ -margin of Appendix B.3: *pay a margin in advance to make a comparison decidable*.

Lemma B.27 (Tails contain late certificates). *If x has tail positivity in the propositional form, with witnesses k, N , then $\exists n W_x(n)$. Indeed $n_0 := N + k + 2$ gives $2\varepsilon_{n_0} \leq 2\varepsilon_{k+1} = \varepsilon_k < x_{n_0}$. $\square$*

This inference goes from a proposition to a proposition, so opening the existential inside it is legitimate—under the BHK reading and in Lean alike. What has changed is essential: an existential whose body contains a universal has been replaced by the existential of a *decidable one-point predicate*. Lemmas B.26 and B.27 are the dyadic-gauge version of Lemma 2 of *Foundations*, Chapter 2.

Proposition B.28 (Choice-free round trip). *For a presented real x , the propositional form of $0 < x$ and the tail-positivity data are interconvertible without any choice principle.*

Proof. *Data to proposition is forgetting. For the converse: by Lemma B.27, pass from the propositional form to the existential of the decidable predicate W_x . Test $W_x(0), W_x(1), W_x(2), \dots$ in order and return the first success—a minimization search. Termination is guaranteed by the existential; the search itself is well-founded recursion over a decidable predicate and requires no choice. (In Lean it is minimization through the accessibility predicate, whose elimination into data for this purpose is part of the calculus and, as Carneiro’s axiomatization records, independent of the choice axiom; the axiom output of all constructions of this subsection lies within `{propext, Quot.sound}` and is covered by the audit of Section 10.) The certificate so found is converted to tail data by Lemma B.26. $\square$*

One reading deserves record. *Foundations* states that an element of $\mathbb{R}^+$ is properly a pair—a sequence together with a witness—and then declares that it will take “the position that the n was implicitly there all along”, adopting the loose usage. Proposition B.28 is the machine-checked statement that this looseness is harmless without choice: the witness said to exist implicitly can always be recomputed, on the spot, whenever it is needed.

Remark B.29 (Consequences, the dual certificate, and the limitation). (i) *Cotransitivity as data*. Once evidence for $a < b$ is at hand—a witness index and its slack—comparison with any c is decided by a single rational comparison at that index; this is the mechanism behind the branch of Appendix B.3, and behind Section 8(i).

(ii) *Reciprocals*. The reciprocal of a presented real is a partial function taking positivity data as an argument. Abstract constructive real-number interfaces *posit* the reciprocal, with its positivity witness, as an axiomatic field; the presented side first computes the witness by the round trip and then builds the reciprocal. This is the smallest unit of the “abstract versus presented” difference of Section 9.1.

(iii) *The dual: an upper one-point certificate.* The same one-point-plus-regularity budget works upward: from the zeroth approximation alone,

$$|x_n| \leq |x_0| + \varepsilon_n + \varepsilon_0 \leq |x_0| + 2 \quad (\text{all } n).$$

This is the faithful dyadic version of the standard bound K_x of *Foundations* (the least integer exceeding $|x_1| + 2$; the *Foundations* indexes its sequences from 1, so x_1 is the first approximation there, consistent with the gauge translation $x'_0 := x_{2^0} = x_1$), and it is why the Archimedean modulus in Appendix B.7 is constant. The positivity certificate from below and the boundedness certificate from above are two directions of one mechanism: regularity propagates one-point information to the whole sequence.

(iv) *Limitation.* The round trip exists only on the strong-order (apartness) side. The weak order $\leq$ is *defined* as $\neg <$ and has no one-point certificate; this is the root of the fact that the weak sides of the pairs in Appendix B.8 are handled only through negation. Finally, none of this is a new theorem in the classical sense: in classical mathematics the proposition/data distinction collapses, and a conversion theorem between the two has no place in the traditional literature. What is worth reporting is that in a setting where the distinction does not collapse, the round trip holds without choice, and that this fact is machine-checked.

B.13 A dictionary with Booi’s locators

The correspondence described here is conceptual; no locator type occurs in the artifact. Section and theorem numbers refer to Booi [26] in its arXiv version.

(i) *The dictionary.* Booi works in Martin-Löf type theory with propositional truncation $\|\cdot\|$, function extensionality, and propositional extensionality, and enforces the property/structure distinction down to the verbs (“there exists” against “has”, “one can find”, “one can construct”). Lean’s Prop/Type split is the counterpart of the $\|\cdot\|$ discipline; his decidable propositions correspond to decidability data; and his search engine—Escardó’s theorem as presented in his Theorem 3.5.1, that for a decidable predicate on $\mathbb{N}$ the truncated existence yields the untruncated witness pair, constructed as the *least* witness (his §3.4–3.6)—corresponds to the minimization search of Proposition B.28. The ambient foundations differ (his is compatible with univalence; Lean has definitional proof irrelevance with UIP, Remark 10.1), but Booi notes that univalence is not needed for this range, so the dictionary is sound across the foundations.

(ii) *Locators and their normal form.* A locator for a real x is $\ell : \Pi(q, r : \mathbb{Q}). q < r \rightarrow (q < x) + (x < r)$ (his Definition 3.1.1): locatedness with the disjunction replaced by an untruncated sum; one real may have many locators. His Lemma 3.4.2 shows a locator equivalent to a *decidable predicate* (“locates right”) together with two propositional soundness clauses. That is the same normal form as the one-point certificate: Definition B.25 is a decidable predicate, and Lemmas B.26–B.27 are its soundness; structure of this kind *is* a decidable predicate paired with propositional soundness.

(iii) *The same engine, asymmetric costs.* His central technique is to reason about locator structure through propositions and to construct unique answers by bounded search (his §3.4–3.5), obtaining rational bounds (Lemma 3.6.1) and Archimedean structure $x < y \rightarrow \Sigma(q : \mathbb{Q}). x < q < y$ (Lemma 3.7.2); Proposition B.28 is the same principle applied to positivity. The costs, however, are asymmetric: from a presentation, bounds can simply be read off (Remark B.29(iii): the first approximation and a constant modulus suffice), whereas a locator hides the sequence, so that even a bound needs a search over an enumeration of $\mathbb{Q}$; that a bound can be obtained from the locator alone at all is what Booi marks “perhaps surprisingly”. This is the plainest exhibition of what carrying representations buys.

(iv) *Where presented reals sit.* His Theorem 3.9.7, for mere existence: that a locator exists, that a signed-digit representation exists, that a rational Cauchy sequence with limit x exists, and that x is a Cauchy real, are equivalent; the corresponding statement at the level of structure is his Theorem 3.9.5, that locators and signed-digit representations are interdefinable. Presented reals carry modulus-bearing rational sequences by definition, so through this dictionary every object of the present development lives on the locator-carrying side. Indeed a presented real carries a canonical locator on paper: given $q < r$, pick n with $2\varepsilon_n < r - q$ and compare x_n with the midpoint $(q + r)/2$; the regularity margin decides $(q < x) + (x < r)$. His Theorem 3.10.4 states that, *presupposing the property* of being a Dedekind cut, a single locator suffices for the remaining structure (boundedness and roundedness in structural form) to follow—the same architecture as Remark B.29, where one certificate plus regularity yields bounds, comparisons, and moduli.

(v) *The same taboo, dual disciplines.* His Lemmas 3.10.1 and 3.10.3: a global assignment of locators to *all* Dedekind reals defines, through the observation “does the locator of x locate right at $0 < 1$ ”, a strongly nonconstant map $\mathbb{R}_{\mathbf{D}} \rightarrow \mathbf{2}$, and the constructive taboo WLPO follows. The same phenomenon has an appearance here: the least witness returned by the minimization search depends on the presentation and does *not* respect Bishop equality—two presentations of one real may return different least witnesses—and collapsing the witness to the Bishop-equality quotient corresponds exactly to truncating the locator to a property on his side. The two responses are dual. Booi keeps locators outside the definition of the real-number type, as structure carried alongside it, so that extensionality is preserved (his Theorem 3.10.5 and Corollary 3.9.8 analyse where the truncation may be placed: digit representations for all Dedekind reals are equivalent to the coincidence of the Cauchy and Dedekind reals, which is not constructively provable). The present development does not define its reals as the quotient—the carrier stays on the presentation side—so the collapsing operation never arises, and it pays instead the discipline of using extracted witnesses only inside Bishop-equality-invariant conclusions (the respect obligations of Section 11.2). These are dual answers, on the extensional side and on the presented side, to one and the same obstruction.

(vi) *The strategy, at both ends.* Booi obtains the exact intermediate value theorem—for which Bridges–Richman-style constructive analysis ordinarily consumes dependent choice—without choice principles, by carrying locators (his Theorem 4.3.5); and his Theorem 4.2.1 shows the Riemann integral preserves locators when a uniform-continuity modulus is carried as data, with locators for the endpoints and a structure lifting the function to locators also among the hypotheses. The present development does the corresponding thing at the other end, in the measure theory that Richman singled out as a principal obstacle (Section 8): the Bishop–Cheng profile route to dominated convergence, whose countable-avoidance step meets the choice obstruction discussed in Section 8 in unrestricted settings, is confined to $\{\text{propext}, \text{Quot.sound}\}$ by presenting the reals and carrying witnesses. The two are instances, at the elementary-analysis end and at the measure-theory end, of one strategy: *replace invocations of choice by carried structure*. His open problem—whether pointwise continuous functions lifting to locators must carry structural continuity—does not arise in the present design, where continuity and convergence moduli are data from the start; that is a consequence of the difference in starting points, not a superiority of either.

References

- [1] Errett Bishop. *Foundations of Constructive Analysis*. McGraw-Hill, New York, 1967.
- [2] Errett Bishop and Henry Cheng. *Constructive Measure Theory*. Memoirs of the American Mathematical Society, no. 116, American Mathematical Society, Providence, 1972.

- [3] Errett Bishop and Douglas Bridges. *Constructive Analysis*. Grundlehren der mathematischen Wissenschaften 279, Springer, 1985.
- [4] Percy J. Daniell. A general form of integral. *Annals of Mathematics*, 2nd ser., 19(4):279–294, 1918.
- [5] Yuen-Kwok Chan. *Foundations of Constructive Probability Theory*. Encyclopedia of Mathematics and its Applications, Cambridge University Press, 2021. Chapter 4, “Integration and measure”.
- [6] Radu Diaconescu. Axiom of choice and complementation. *Proceedings of the American Mathematical Society*, 51(1):176–178, 1975.
- [7] Fred Richman. Meaning and information in constructive mathematics. *The American Mathematical Monthly*, 89(6):385–388, 1982.
- [8] Douglas S. Bridges, Hajime Ishihara, Michael Rathjen, and Helmut Schwichtenberg (eds.). *Handbook of Constructive Mathematics*. Encyclopedia of Mathematics and its Applications 185, Cambridge University Press, 2023. DOI: 10.1017/9781009039888.
- [9] Yuen-Kwok Chan. A leisurely random walk down the lane of a constructive theory of stochastic processes. In [8], pp. 333–356.
- [10] Francesco Ciraulo. Subspaces in pointfree topology: towards a new approach to measure theory. In [8], pp. 426–444.
- [11] Fred Richman. Countable choice. In [8], pp. 515–524.
- [12] Maria Emilia Maietti and Giovanni Sambin. The Minimalist Foundation and Bishop’s constructive mathematics. In [8], pp. 525–563.
- [13] Robert S. Lubarsky. Inner and outer models for constructive set theories. In [8], pp. 584–635.
- [14] Helmut Schwichtenberg. Computational aspects of Bishop’s constructive mathematics. In [8], pp. 715–748.
- [15] Kenji Miyamoto. Application of constructive analysis in exact real arithmetic. In [8], pp. 749–776.
- [16] Mark Bickford. Efficient algorithms from proofs in constructive analysis. In [8], pp. 777–805.
- [17] Ulrich Berger and Monika Seisenberger. On the computational content of choice principles. In [8], pp. 806–825.
- [18] Fred Richman. The fundamental theorem of algebra: a constructive development without choice. *Pacific Journal of Mathematics*, 196(1):213–230, 2000.
- [19] Douglas Bridges, Fred Richman, and Peter Schuster. A weak countable choice principle. *Proceedings of the American Mathematical Society*, 128(9):2749–2752, 2000.
- [20] Fred Richman. Constructive mathematics without choice. In P. Schuster, U. Berger, and H. Osswald (eds.), *Reuniting the Antipodes—Constructive and Nonstandard Views of the Continuum*, pp. 199–205, Kluwer, 2001.

- [21] Robert S. Lubarsky. On the Cauchy completeness of the constructive Cauchy reals. *Electronic Notes in Theoretical Computer Science*, 167:225–254, 2007. DOI: 10.1016/j.entcs.2006.09.012.
- [22] Bas Spitters. Constructive algebraic integration theory without choice. In T. Coquand, H. Lombardi, and M.-F. Roy (eds.), *Mathematics, Algorithms, Proofs*, Dagstuhl Seminar Proceedings 05021, pp. 1–13, 2006. DOI: 10.4230/DagSemProc.05021.9.
- [23] Thierry Coquand and Erik Palmgren. Metric Boolean algebras and constructive measure theory. *Archive for Mathematical Logic*, 41:687–704, 2002. DOI: 10.1007/s001530100123.
- [24] Iosif Petrakis. Predicative Bishop–Cheng measure theory. Overview/lecture abstract, LMU München, 2021.
- [25] Iosif Petrakis and Max Zeuner. Pre-measure spaces and pre-integration spaces in predicative Bishop–Cheng measure theory. *Logical Methods in Computer Science*, 20(4):2:1–2:33, 2024. DOI: 10.46298/lmcs-20(4:2)2024.
- [26] Auke B. Booij. Extensional constructive real analysis via locators. *Mathematical Structures in Computer Science*, 31(1):64–88, 2021. DOI: 10.1017/S0960129520000171. Also [arXiv:1805.06781](#); the section and theorem numbers cited in this paper follow the arXiv version.
- [27] Herman Geuvers and Milad Niqui. Constructive reals in Coq: axioms and categoricity. In *Types for Proofs and Programs (TYPES 2000)*, LNCS 2277, pp. 79–95, Springer, 2002. DOI: 10.1007/3-540-45842-5_6.
- [28] Vincent Séméria. Nombres réels dans Coq. In *Actes des 31es Journées Francophones des Langages Applicatifs (JFLA 2020)*, pp. 104–111, 2020. HAL record [hal-02427360](#); archived paper PDF.
- [29] Claudio Sacerdoti Coen and Enrico Tassi. A constructive and formal proof of Lebesgue’s dominated convergence theorem in the interactive theorem prover Matita. *Journal of Formalized Reasoning*, 1(1):51–89, 2008. DOI: 10.6092/issn.1972-5787/1334.
- [30] Sylvie Boldo, François Clément, and Louise Leclerc. A Coq formalization of the Bochner integral. Inria Research Report RR-9456; [arXiv:2201.03242v2](#), 2022.
- [31] Leonardo de Moura and Sebastian Ullrich. The Lean 4 theorem prover and programming language. In *Automated Deduction (CADE 28)*, LNCS 12699, pp. 625–635, Springer, 2021. DOI: 10.1007/978-3-030-79876-5_37.
- [32] Mario Carneiro. *The Type Theory of Lean*. Master’s thesis, Carnegie Mellon University, 2019.
- [33] Freek Wiedijk. The de Bruijn factor. Poster, Theorem Proving in Higher Order Logics (TPHOLs 2000), 2000. [cs.ru.nl/~freek/factor](#).
- [34] The mathlib Community. The Lean mathematical library. In *Certified Programs and Proofs (CPP 2020)*, pp. 367–381, ACM, 2020. DOI: 10.1145/3372885.3373824.
- [35] The mathlib Community. `Mathlib.MeasureTheory.Integral.DominatedConvergence`. Source at the artifact’s pinned `mathlib` revision [c5ea00351c28e24afc9f0f84379aa41082b1188f](#); accessed 2026-07-09.

- [36] Vincent Séméria and the CoRN contributors. Bishop–Cheng constructive measure theory declarations in CoRN. Audited CoRN repository snapshot, commit `ada7c0b497ff15dd67cf7932c6f20e143a2aee2f`. Audited source file `CMTMeasurableFunctions.v`. Profile and countable-avoidance source `CMTprofile.v`.
- [37] Toshihisa Urahigashi. `choicelfree-bc-dct`: a Lean 4 artifact for choice-free Bishop–Cheng dominated convergence. Version 0.6.0. Software deposit, Zenodo. Concept DOI (resolves to the latest version): `10.5281/zenodo.21850965`. Version DOIs of earlier deposits: v0.4.1 `10.5281/zenodo.22137161`, v0.4.0 `10.5281/zenodo.21854936`, v0.3.0 `10.5281/zenodo.21850966`.