EXPLICIT INITIAL Θ-DATA AND THE TWO READINGS OF STEP (V) OF IUT IV, THEOREM 1.10: A CATALOGUE

TOSHIHISA URAHIGASHI

Abstract. We collect the explicit initial Θ-data, in the sense of Definition 3.1 of the first paper of Mochizuki's series on inter-universal Teichmüller theory, that the author has constructed over $\mathbb{Q}(\sqrt{5})$ and $\mathbb{Q}(i)$: three published earlier and four given here, one of them for fourteen values of ℓ. Each has a split rational prime above which the two places of the field of moduli are either one of good and one of multiplicative reduction, or both of multiplicative reduction with different orders of the q-parameter. For each datum we tabulate the local invariants and compare, in Step (v) of the proof of Theorem 1.10 in the fourth paper, the displayed comparison module with the hull component of the union of possible images. Under the reading in which the theta value stays at the label j, as in Step (v) and in what the public Lean formalization lana-agents/iut calls the reading of IUT IV, every comparison is a containment. Under the reading in which label permutations act before the hull is taken, which the same formalization calls the literal reading, and under stated membership hypotheses, every datum gives a non-containment for a single component; at primes with a good and a bad place the excess grows like j^2, and for three data the weighted average over the components also exceeds the bound displayed at the prime. No conclusion is drawn about Corollary 3.12, the final inequality of Theorem 1.10, or the abc conjecture.

1. Introduction

Explicit initial Θ-data in the sense of [IUT1, Def. 3.1] are needed whenever the estimates of [IUT4] are to be evaluated on actual numbers. Dupuy and Hilado computed one for the curve 11a1 over $\mathbb{Q}$ [DH1, §2.10], and Zhou constructed families from Frey–Hellegouarch curves [Zhou1, Zhou2]; in these constructions the field of moduli is $\mathbb{Q}$, so a rational prime carries a single place of it. The author has published three data over quadratic fields [First, Second, Note]. This note collects them with four further data, all over $\mathbb{Q}(\sqrt{5})$ or $\mathbb{Q}(i)$, and records for each the local invariants above one split prime and the comparisons in Step (v) of [IUT4, Thm. 1.10, proof].

The comparisons depend on how the label permutations contained in the indeterminacy (Ind1) enter the union of possible images of [IUT3, Cor. 3.12]. That they permute tensor factors, and that a simplified computation of the hull may omit this, is noted in [DH, §6.2, footnote 7]; procession automorphisms are described as permutations of tensor factors in [DH1, §4.6]. The public Lean repository [LANA-iut] formalizes two readings of Corollary 3.12, one with the theta value at the label j (its reading of IUT IV) and one with the theta value at any label (its literal reading), shows that the first implies the second, and records in prose that the two give the same regions when the orders of the q-parameters agree at all places above each prime, that two bad places with different orders already break this, and that at primes with a good and a bad place, which arise from the Belyi reduction, the literal reading is governed by the minimum rather than the average. The data below are explicit instances of these cases.

What is new here is the data themselves and the numbers: initial Θ-data whose field of moduli is not $\mathbb{Q}$ (we have not found such data in the literature; an explicit example of this kind was asked for in [MO]), data realizing both kinds of prime just described, and exact values of the comparisons of Step (v) on them. The distinctions between the readings, and the mechanism by which the readings differ, are not new; see §6.

Date: Version 1.1.0, 6 November 2026.

We do not claim that Corollary 3.12, the final inequality of Theorem 1.10 or the abc conjecture is false, that either reading is the one the source intends, or that the hypotheses of §2 identify the source's union of possible images.

2. Step (v) and the readings

We use the notation of [Note, §1]. For a rational prime p and a place v of F_{mod} above p let k_v be the completion of K at the selected place above v, e_v its ramification index over $\mathbb{Q}_p$, $\mathfrak{d}_v = (e_v - 1)/e_v$ the order of its different when $p \nmid e_v$, and $\rho(v) = \mathrm{ord}_p(q_v)$ if $v \in \mathbb{V}^{\mathrm{bad}}_{\mathrm{mod}}$, $\rho(v) = 0$ otherwise. For $j \in \{1, \dots, \ell^*\}$ and a tuple $\mathbf{v} = (v_0, \dots, v_j)$ of places above p, Step (v) compares the component at $\mathbf{v}$ of the union U_j of possible images with a module $M_{\mathbf{v}}$; when every $e_{v_i} \le p - 2$,

$$\mu^{\log}(M_{\mathbf{v}}) \le \Big(-j^2\rho(v_j) + \mathfrak{d}_I + 1\Big)\log p, \qquad \mathfrak{d}_I = \textstyle\sum_{i=0}^{j} \mathfrak{d}_{v_i} \tag{1}$$

[IUT4, Prop. 1.4(iii), p. 13], [Note, (1)]. All values below are in units of $\log p$.

We consider the following readings of [IUT3, Cor. 3.12]; none is proved here.

- (V) The theta value stays at the label j, as in Step (v) and in the reading of IUT IV of [LANA-iut]. (The file `Iut/Concrete/Ind1Invariance.lean` of [LANA-iut] also records that taking the hull for each image under (Ind1) separately gives the value of IUT IV.)
- (L) Label permutations fixing 0 act on possible-image regions before the hull is taken; this is (R) of [Note, §3]. (L^0) also admits permutations moving 0, as in [First-full, (SA2)].
- ($\mathrm{M_b}$) At a bad place the region of the Θ-pilot object is the theta value times a lattice containing 1, and its Kummer image is a possible image; this is (M) of [Note, §3], which gives [Note, Lemma 3].
- ($\mathrm{M_g}$) At a good place the region of the Θ-pilot object at a tuple whose factor of label j lies over it contains $1 \otimes \cdots \otimes 1$, the theta value being 1 there, and its Kummer image is a possible image; a source basis for this is given in [First-full, §7, (SA1)]. The proof of [Note, Lemma 3] then gives, at such a tuple $\mathbf{u}$, an element $X \in U_j$ with $\mathrm{pr}_{\mathbf{u}}(X) = \chi(1)$, $\chi = \otimes_i \chi_i$, $\chi_i \in \mathcal{A}(k_{u_i})$.

Under (V), for every datum and every tuple, the elements of [Note, Lemma 3] lie in $M_{\mathbf{v}}$: this is [Note, Prop. 6] with $b = v_j$, and at a good v_j the same argument with $\lambda = 0$ applies to $\chi(1)$.

Under (L), elements are moved by [Note, Lemma 4]. We use three kinds, with lower bounds for the hull component obtained from [Note, (1.1)–(1.3)] and the argument of [Note, Thm. 5] (if U_j is not relatively compact its hull has log-volume $+\infty$ and every finite bound fails).

- (BB) Two bad places b_s, b_l above p with $\rho(b_s) < \rho(b_l)$, $j \ge 2$, assuming ($\mathrm{M_b}$) and (L): the theta value of b_s is moved by a transposition $(t\ j)$, $1 \le t < j$, to a tuple with $v_t = b_s$ and $v_j = b_l$; the hull component is $\ge -\lceil j^2\rho(b_s)\rceil$.
- (GB) A good place g and a bad place b above p, assuming ($\mathrm{M_g}$) and (L) if $j \ge 2$, (L^0) if $j = 1$: the element $\chi(1)$ at a tuple with label j over g is moved to a tuple with label j over b; the hull component is ≥ 0.
- (GB_L) The variant of [First-full, Thm. 8.1], under its hypothesis (SA): the moved region contains the log-unit lattice, and the hull component is $\ge -\sum_i e_{v_i}^{-1}$.

Proposition 1. *Let $j \ge 1$ and let $\mathbf{v}$ range over the tuples of places above p. Among the elements just described, the largest excess of the hull component over the bound* (1) *is*

$$\begin{aligned}
&\textit{(BB)} && j^2\rho(b_l) - \lceil j^2\rho(b_s)\rceil - 1 - \mathfrak{d}_{b_s} - \mathfrak{d}_{b_l} - (j-1)\,\mathfrak{d}_{\min}, && \mathbf{v} = (b_*, b_s, b_*, \dots, b_*, b_l),\\
&\textit{(GB)} && j^2\rho(b) - \mathfrak{d}_b - 1, && \mathbf{v} = (g, \dots, g, b),\\
&\textit{(GB}_L\textit{)} && j^2\rho(b) - (j+2), && \textit{any } \mathbf{v} \textit{ with } v_j = b,
\end{aligned}$$

where $\mathfrak{d}_{\min}$ is the least $\mathfrak{d}_v$ over the places above p and b_* a place attaining it. A positive value is a non-containment of the component at $\mathbf{v}$ in $M_{\mathbf{v}}$.

Proof. The bound (1) at $\mathbf{v}$ is $-j^2\rho(v_j) + \sum_i \mathfrak{d}_{v_i} + 1$. In (BB) and (GB) the lower bound does not depend on the factors other than the moved one and the one of label j, so their $\mathfrak{d}$ is minimized;

in (GB) the good place has $\mathfrak{d}=0$. In (GB_L), $\sum_i (e_{v_i}^{-1}+\mathfrak{d}_{v_i}) = j+1$ for tame e_{v_i}. The values were also checked against an enumeration of all tuples for $\ell=7$ and $j\le 3$ (§7). □

3. The data

All data have $E_0\colon y^2=x(x-1)(x-a)$ over $F_0=F_{\mathrm{mod}}$, $F=F_0(\sqrt{-1},E_0[15])$, $K=F(E_0[\ell])$, and $\mathbb{V}_{\mathrm{mod}}^{\mathrm{bad}}$ the places of multiplicative reduction outside 2ℓ, as in [Note, §2]; π generates a prime of F_0 of degree one over $p=\mathrm{N}(\pi)$, and $\bar\pi$ is its conjugate. Table 1 lists them.

datum	π	p	a	ℓ	(e,ρ) above p	proof
D_{211}	$16+3\sqrt{5}$	211	π^{29}	7	$g\,(1,0)$, $b\,(105,\frac{29}{7})$	[First, Thm. 1]
D_{109}	$10+3i$	109	π^{29}	7	$g\,(1,0)$, $b\,(105,\frac{29}{7})$	[Second]
D_2	$14164+5825\sqrt{5}$	30965771	$\pi\bar\pi^6$	7	$b_1(105,\frac{1}{7})$, $b_2(35,\frac{6}{7})$	[Note, Prop. 2]
D_3	as D_2	30965771	$\pi^6\bar\pi^{-30}$	7	$b_1(35,\frac{6}{7})$, $b_2(7,\frac{30}{7})$	Prop. 6
D_{18}	as D_2	30965771	π^{18}	7	$g\,(1,0)$, $b\,(35,\frac{18}{7})$	Prop. 7
D_{15}	$-5932-919\sqrt{5}$	30965819	π^{15}	7	$g\,(1,0)$, $b\,(7,\frac{15}{7})$	Prop. 8
$D_{2357}^{(\ell)}$	$41+26i$	2357	π^{472}	$\ell\in\Lambda$	$g\,(1,0)$, $b\,(15\ell,\frac{472}{\ell})$	[Fourteen]

Table 1. The data; π lies in $\mathbb{Z}[\sqrt{5}]$ or $\mathbb{Z}[i]$, and $F_0=\mathbb{Q}(\pi)$. $\Lambda=\{89,97,101,103,107,109,113,127,131,137,139,149,151,157\}$. In every case $e\le p-2$ at both places, so (1) applies. The curve of $D_{2357}^{(\ell)}$ is that of [Second, §13]; the verification of [IUT1, Def. 3.1] for every $\ell\in\Lambda$ is in [Fourteen].

Proposition 2. *The data D_3, D_{18} and D_{15} extend to initial Θ-data in the sense of* [IUT1, Def. 3.1] *satisfying the hypotheses of* [IUT4, Thm. 1.10]*, with $F_{\mathrm{mod}}=F_0$ and the local data of Table 1; for each, $p>e^*_{\mathrm{mod}}\cdot\ell=7741440$.*

The proofs are in Appendix A; each is the proof of [Note, Prop. 2] with the changes stated there.

4. The comparisons

Theorem 3. *For each positive entry of Table 2, under the readings stated for its column, the component of U_j at the tuple of Proposition 1 is not contained in $M_{\mathbf{v}}$, and the log-volume of its hull exceeds the bound* (1) *by at least the value shown. Under (V) the elements considered lie in $M_{\mathbf{v}}$ for every datum and every tuple.*

Proof. Proposition 1 with the values of Table 1, and the remark on (V) in §2. □

datum	kind	$j=1$	$j=2$	$j=3$
D_2	(BB)	—	$-\frac{158}{105}$	$\frac{17}{21}$ ([Note, Thm. 5]: $\frac{27}{35}$)
D_3	(BB)	—	$\frac{331}{35}$	$\frac{911}{35}$
D_{211}, D_{109}	(GB) / (GB_L)	$\frac{226}{105}$ / $\frac{8}{7}$	$\frac{1531}{105}$ / $\frac{88}{7}$	$\frac{3706}{105}$ / $\frac{226}{7}$
D_{18}	(GB) / (GB_L)	$\frac{3}{5}$ / $-\frac{3}{7}$	$\frac{291}{35}$ / $\frac{44}{7}$	$\frac{741}{35}$ / $\frac{127}{7}$
D_{15}	(GB) / (GB_L)	$\frac{2}{7}$ / $-\frac{6}{7}$	$\frac{47}{7}$ / $\frac{32}{7}$	$\frac{122}{7}$ / $\frac{100}{7}$

Table 2. Largest excess of the hull component over (1), in units of $\log p$. (BB) assumes ($\mathrm{M_b}$) and (L); (GB) assumes ($\mathrm{M_g}$), and (L) for $j\ge 2$, (L^0) for $j=1$; (GB_L) assumes (SA) of [First-full]. For D_2, $j=3$, the tuple of Proposition 1 is (b_2,b_1,b_2,b_2); the tuple (b_1,b_1,b_1,b_2) of [Note, Thm. 5] gives $\frac{27}{35}$.

For $D_{2357}^{(\ell)}$ the values are $\frac{472j^2}{\ell}-2+\frac{1}{15\ell}$ (GB) and $\frac{472j^2}{\ell}-(j+2)$ (GB_L), positive for every $j\ge 1$ and $\ell\in\Lambda$; at $\ell=157$ they are $\frac{2371}{2355}$ and $\frac{1}{157}$ for $j=1$, and about 18288.8 and 18210.8 for $j=\ell^*=78$. At primes with a good and a bad place the excess grows like $j^2\rho(b)$, the order of the term $-j^2\rho(v_j)$ of (1) itself; at the bad-bad primes of D_2 and D_3 it is of the order $j^2(\rho(b_l)-\rho(b_s))$.

Theorem 4. *For D_2, D_3, D_{18} and D_{15} the two places above p have degree one, so the tuples have equal weights, and $p > e^*_{\mathrm{mod}}\ell$, so the terms with $\log(\mathfrak{s}_p^{\leq})$ vanish* [IUT4, pp. 23, 25]. *Under (M_{b}), (M_{g}) and (L), the weighted average over the tuples of the hull components at stage j is at least $L_j = -\mathbb{E}_{\mathbf{v}}\big[\min_{1\leq t\leq j} c(v_t)\big]$, where $c(b) = \lceil j^2\rho(b)\rceil$ for a bad place and $c(g) = 0$ for a good one. The weighted average that Step (v) displays at stage j is $B_j = (j+1)\bar{\mathfrak{d}} - j^2\bar{\rho} + 1$, and the final bound displayed at p is $F_p = \frac{\ell+1}{4}\big\{(1+\frac{4}{\ell})\bar{\mathfrak{d}} - \frac{\ell}{3}\bar{\rho} + \frac{4}{\ell}\big\}$* [IUT4, pp. 28–29], *where $\bar{\mathfrak{d}}$ and $\bar{\rho}$ are the averages over the two places. The values are in Table 3.*

Proof. For a tuple $\mathbf{v}$ and $t \in \{1,\dots,j\}$, [Note, Lemma 3] or ($\mathrm{M_g}$) at the tuple $\mathbf{v}\circ(t\ j)$ and [Note, Lemma 4] with the transposition $(t\ j)$, which fixes 0, give an element of the component at $\mathbf{v}$ whose factors have order at most $c(v_t)$ in total [Note, (1.1)–(1.3)]; so the hull component at $\mathbf{v}$ has log-volume at least $-\min_t c(v_t)$. For B_j: $\log(\mathfrak{d}_p^K) = \bar{\mathfrak{d}}\log p$, $\log(q_p) = 2\ell\bar{\rho}\log p$ and $\log(\mathfrak{s}_p^{\mathbb{Q}}) = \log p$ in the displays of [IUT4, pp. 28–29]. The values were computed exactly (§7). □

datum	$L_1 - B_1$	$L_2 - B_2$	$L_3 - B_3$	$\frac{1}{3}\sum_j L_j - F_p$
D_2	$-\frac{727}{210}$	$-\frac{517}{140}$	$-\frac{1333}{420}$	$-\frac{5477}{1470}$
D_3	$-\frac{114}{35}$	$-\frac{67}{70}$	$\frac{1851}{280}$	$\frac{3089}{5880}$
D_{18}	$-\frac{153}{70}$	$-\frac{9}{140}$	$\frac{197}{35}$	$\frac{2687}{2940}$
D_{15}	$-\frac{16}{7}$	$-\frac{1}{4}$	$\frac{31}{7}$	$\frac{251}{588}$

TABLE 3. Averaged comparisons at p (units of $\log p$). A positive entry means that the averaged lower bound for the hull exceeds the displayed bound. For D_2 no averaged bound is exceeded; its non-containment is in a single component.

Remark 5. Two variants change the numbers but not the signs of the last column. If the members of the poly-isomorphism over the transposition are taken to have identity components, as one may after identifying the members of the capsule with the model of [IUT1, Def. 4.1(iii)], the rounding $\lceil\cdot\rceil$ can be dropped; the last column becomes $\frac{241}{245}$, $\frac{1059}{980}$, $\frac{249}{392}$ for D_3, D_{18}, D_{15}. If permutations moving 0 are admitted as well (L^0), the minimum runs over $t \in \{0,\dots,j\}$, and without rounding the last column becomes $\frac{1217}{490}$, $\frac{4323}{1960}$, $\frac{1233}{784}$.

5. A global check for D_2

For D_2 the bad places are b_1, b_2 above p, the place above R (the large prime factor of $\mathrm{N}(1-a)$, see [Note, Prop. 2]) where $\mathrm{ord}(q) = 2$, and the two places above 11, where $\mathrm{ord}(q) = 2$ each; so $\log q = \frac{1}{2}(14\log p + 2\log R + 4\log 11) = 240.09\ldots$ in the normalization of [IUT4, Thm. 1.10]. The final inequality of Theorem 1.10 holds for D_2 with a wide margin, its left side $\frac{1}{6}\log q \approx 40.0$ being far below the constant $20e^*_{\mathrm{mod}}\ell = 154828800$ on its right. The inequality that [LANA-iut] formulates for Step (xi), $(s-1)|\log q| \leq \mu^{\log}(\mathrm{hull}(\mathrm{VC}(I)))$ with $s = \ell(\ell+1)/12$ and $|\log q| = \log q/2\ell$ (file `Iut/WallEquiv/README.md`), also holds for D_2: the left side is $62.88\ldots$, while by the formula $\sum_c w_c \sum_j \rho_{c_j}\log p$ for the prime components recorded there, with $\rho = 1 - 1/e$ at the tame places above p and R and the other components nonnegative, the right side is at least $3\big(\frac{1}{2}(\frac{104}{105} + \frac{34}{35})\log p + \frac{1}{2}\cdot\frac{104}{105}\log R\big) = 221.0\ldots$. The place above R, where q has order only 2, supplies most of this.

6. Earlier work and dates

- Explicit initial Θ-data: [DH1, §2.10] (11a1 over $\mathbb{Q}$, $\ell = 13$); [Zhou1, Zhou2] (families, $F_{\mathrm{mod}} = \mathbb{Q}$); [MFHMP, Def. 4.1] ($\mu_6$-initial Θ-data, no numerical example); an explicit example over an extension of $\mathbb{Q}(i)$ was asked for in [MO].
- The permutations of tensor factors in (Ind1): [DH1, §4] (2020); [DH, §6.2, footnote 7] (2020).

- The two readings, the agreement when the orders of q agree above each prime, the failure at two bad places with different orders, the minimum versus the average, and primes with a good and a bad place from the Belyi reduction: [LANA-iut], file `Iut/Concrete/Literal.lean`, first committed on 11 October 2026 (UTC); the per-image hull, file `Iut/Concrete/Ind1Invariance.lean`, the same day.
- The question of the source of $\lambda = \mathrm{ord}(q^{j^2})$ in Step (v): [Takkun] (July 2026). The last estimate of Step (v) restated without an independent proof: [JoshiIV, Prop. 6.10.9].
- A different loss in the hull of a tensor packet, from boxing the integral structure: [Zeng] (30 October 2026).
- The data of this note: D_{211} on 6 September 2026 [First], D_{109} on 7 September 2026 [Second], D_2 on 6 October 2026 [Note-v1]; D_3, D_{18}, D_{15} and the values of §4 are new here.

7. Reproducibility

The values of Tables 2 and 3 are computed with exact rational arithmetic by two standard-library Python programs: `catalogue.py` (Proposition 1, with an enumeration of all tuples for $\ell = 7$, $j \le 3$, and checks against the values $\frac{27}{35}$, $\frac{8}{7}$, $\frac{88}{7}$ recorded in [Note, First]) and `averaged.py` (Theorem 4). The arithmetic inputs of Proposition 2 are checked by the programs listed in Appendix A. All these programs, with their output, are in the archive deposited with this note.

Appendix A. Proofs for the new data

In each case the proof of [Note, Prop. 2] and of its Table 1 applies with the changes below; in particular [IUT4, Cor. 2.2] is not used, and conditions (d)–(f) are constructed directly from $j \notin \mathbb{Q}$, the split multiplicative reduction at the selected places and the image containing $\mathrm{SL}_2(\mathbb{F}_7)$, as there.

Proposition 6. *D_3: $a = \pi^6\bar\pi^{-30}$ with $\pi = 14164 + 5825\sqrt{5}$.*

Changes. Now $\mathrm{ord}_{b_1}(j) = -12$ and $\mathrm{ord}_{b_2}(j) = -60$, so $j \notin \mathbb{Q}$. At b_2, where a has a pole of even order, the change of variables $x = aX$, $y = a\sqrt{a}\,Y$, with $\sqrt{a} = \pi^3\bar\pi^{-15} \in F_0$, gives $Y^2 = X(X-1)(X-a^{-1})$ with $\mathrm{ord}_{b_2}(a^{-1}) = 30$. The numerator of $1-a$ is $-\prod_{d\mid 6}\Phi_d(\pi,\bar\pi^5)$, with the homogeneous cyclotomic forms Φ_d, no two of which have a common prime divisor outside $2, 3$. None of the four norms $\mathrm{N}(\Phi_d(\pi,\bar\pi^5))$ (38, 38, 75 and 75 digits) is divisible by the seventh power of an odd prime (checked by a segmented sieve up to the seventh roots and independently by a complete factorization with primality certificates), $1-a$ is a unit at the place above 3 and has order 5 at the place above 2. At the place above 7, a and $1-a$ are units. For the image, the place of degree one above 41 with $\sqrt{5} \equiv 13$ gives $a_{41} = -6$, and $X^2+6X+41$ is irreducible over $\mathbb{F}_7$; the transvection comes from b_1, where $\mathrm{ord}(q) = 12$. At b_1 and b_2 the orders of q are 12 and 60, so $e = 35$ and 7, the differents have orders 34 and 6, and $\rho = \frac{6}{7}$ and $\frac{30}{7}$. □

Proposition 7. *D_{18}: $a = \pi^{18}$ with $\pi = 14164 + 5825\sqrt{5}$.*

Changes. The six Legendre orbit comparisons of a fail in $\mathbb{Z}[\sqrt{5}]$, so $j \notin \mathbb{Q}$. We have $a - 1 = \prod_{d\mid 18}\Phi_d(\pi)$; no norm of a factor is divisible by the seventh power of a prime (a sieve up to the seventh roots of the norms, at most 2634850), two factors meet only above 2 and 3, which are inert, the order of $1-a$ at the place above 3 is 0, and the place above 2 is not selected. So every positive order of $1-a$ at a selected place is at most 6, the only positive order of a is 18, at v_π, and 7 divides neither $\mathrm{N}(a) = p^{18}$ nor $\mathrm{N}(1-a)$. For the image, the place of degree one above 31 with $\sqrt{5} \equiv 6$ gives $a_{31} = -8$, and $a_{31}^2 - 4\cdot 31 \equiv 3 \pmod 7$ is not a square; the transvection comes from v_π, as $7 \nmid 36$. Above p, $\bar\pi$ gives a good place g with $e = 1$; at $b = v_\pi$ the curve is split Tate over $\mathbb{Q}_p(\sqrt{-1})$ with $\mathrm{ord}_p(q) = 36$, so $e = 105/\gcd(105, 36) = 35$, the different has order 34 and $\rho = \frac{18}{7}$. □

Proposition 8. *D_{15}: $a = \pi^{15}$ with $\pi = -5932 - 919\sqrt{5}$, $p = 30965819$.*

Changes. p is prime, $p \equiv 4 \pmod 5$ and $p \equiv 3 \pmod 4$; $p = \pi\bar{\pi}$ splits in F_0 and both places have degree one. At $b = v_\pi$, $\mathrm{ord}(a) = 15$; at $g = v_{\bar\pi}$, a is a unit and $a \not\equiv 1$; $1 - a$ is a unit at both, so $\mathrm{ord}_b(j) = -30$, $\mathrm{ord}_g(j) \geq 0$ and $j \notin \mathbb{Q}$. The content of $a - 1$ in $\mathbb{Z}[(1 + \sqrt{5})/2]$ is 2, and $5 \nmid \mathrm{N}(a - 1)$; hence at a split odd prime at most one place divides $a - 1$, no inert odd prime and not the ramified prime 5 does, and at the place above 2 (inert) $1 - a$ has order 1, where $\mathrm{ord}(j) = 6 \geq 0$. We have $\pi^{15} - 1 = \Phi_1(\pi)\Phi_3(\pi)\Phi_5(\pi)\Phi_{15}(\pi)$; the norms of the blocks are pairwise coprime and none has a seventh-power prime factor other than at 2 and 7 (trial division up to the seventh roots, at most 363942203, by two methods with positive and negative controls), and $a - 1$ has order 0 at the places above 3 and 5. So the orders of $1 - a$ at selected places lie in $\{1, \ldots, 6\}$, the order of a at b is 15, and 7 divides neither $\mathrm{N}(a)$ nor $\mathrm{N}(1 - a)$. For the image, at a place of degree one above 11 the reduction is good with $a_{11} = 0$, and $X^2 + 4$ is irreducible over $\mathbb{F}_7$; the transvection comes from b, where $\mathrm{ord}_w(q) = 30e(w|b)$ is prime to 7. At b the curve is non-split multiplicative over $\mathbb{Q}_p$ and split over $\mathbb{Q}_p(\sqrt{-1})$, with $\mathrm{ord}_p(q) = 30$, so $e = 7$, the different has order 6 and $\rho = \frac{15}{7}$; at g, $e = 1$. □

The arithmetic inputs were checked by two programs for each datum, the second written independently from the specification:

- D_3: `certify_wide.py`, `independent_check_wide.py`;
- D_{18}: `certify_datum.py`, `independent_check.py`;
- D_{15}: `certify_mixed15.py`, `independent_check_mixed15.py`.

For D_3, the complete factorization of the four norms, with primality certificates, is checked by `check_pratt_factorizations.py`.

Use of AI tools. Generative AI tools (large language models: Claude by Anthropic, and ChatGPT and Codex by OpenAI) were used extensively in this work, under the author's direction, in finding the data and the proofs, in the computations and in the writing. The author takes full responsibility for the contents of this note.

References

[DH] Taylor Dupuy and Anton Hilado, *Probabilistic Szpiro, baby Szpiro, and explicit Szpiro from Mochizuki's Corollary 3.12*, 2020, arXiv:2004.13108v2 [math.NT].

[DH1] Taylor Dupuy and Anton Hilado, *The statement of Mochizuki's Corollary 3.12, initial theta data, and the first two indeterminacies*, 2020, arXiv:2004.13228 [math.NT]; §4 and §4.6 refer to version 1, §2.10 to version 2 (2025).

[First] Toshihisa Urahigashi, *An explicit initial Θ-datum with mixed reduction at a split prime*, version 0.1.7, Zenodo, doi:10.5281/zenodo.22537342.

[First-full] Toshihisa Urahigashi, *An explicit initial Θ-datum with mixed reduction over a split prime, and the local estimates of inter-universal Teichmüller theory IV*, version 0.1.7, in the deposit of [First]; its §§7–9 record a reading whose application to the source was withdrawn in that version.

[Fourteen] Toshihisa Urahigashi, *Fourteen initial Θ-data on one curve over* $\mathbb{Q}(i)$, version 1.0.0, Zenodo, 6 November 2026, doi:10.5281/zenodo.23990717.

[IUT1] Shinichi Mochizuki, *Inter-universal Teichmüller theory I: construction of Hodge theaters*, Publ. Res. Inst. Math. Sci. **57** (2021), 3–207.

[IUT3] Shinichi Mochizuki, *Inter-universal Teichmüller theory III: canonical splittings of the log-theta-lattice*, Publ. Res. Inst. Math. Sci. **57** (2021), 403–626.

[IUT4] Shinichi Mochizuki, *Inter-universal Teichmüller theory IV: log-volume computations and set-theoretic foundations*, Publ. Res. Inst. Math. Sci. **57** (2021), 627–723; page numbers refer to the preprint version of April 2020.

[JoshiIV] Kirti Joshi, *Construction of arithmetic Teichmüller spaces IV*, arXiv:2403.10430v2 [math.NT].

[LANA-iut] lana-agents, *iut*, Lean repository, `https://github.com/lana-agents/iut`, consulted at commit `ecb3c86` (1 November 2026).

[MFHMP] Shinichi Mochizuki, Ivan Fesenko, Yuichiro Hoshi, Arata Minamide, and Wojciech Porowski, *Explicit estimates in inter-universal Teichmüller theory*, Kodai Math. J. **45** (2022), no. 2, 175–236.

[MO] *Explicit example of elliptic curve of the kind needed for IUTT*, MathOverflow question 299845 (2018).

[Note] Toshihisa Urahigashi, *A local non-containment in Step (v) of IUT IV, Theorem 1.10*, version 3.0.0, Zenodo, 1 November 2026, doi:10.5281/zenodo.23852752.

[Note-v1] Toshihisa Urahigashi, *A local non-containment in Step (v) of IUT IV, Theorem 1.10*, version 1.0.0, Zenodo, 6 October 2026, doi:10.5281/zenodo.23166433.

[Second] Toshihisa Urahigashi, *An initial Θ-datum over an imaginary quadratic field, and numerical values of the local quantities of inter-universal Teichmüller theory IV*, version 1.0.5, Zenodo, doi:10.5281/zenodo.22601333.

[Takkun] Takkun-kohinata, *IUT_LEAN*, GitHub repository, `https://github.com/Takkun-kohinata/IUT_LEAN` (July 2026).

[Zeng] Chunping Zeng, *An integral-structure inflation in the hull operation of IUT Corollary 3.12*, Zenodo, 30 October 2026, doi:10.5281/zenodo.23811410.

[Zhou1] Zhong-Peng Zhou, *The inter-universal Teichmüller theory and new Diophantine results over the rational numbers. I*, arXiv:2503.14510 [math.NT], 2025.

[Zhou2] Zhong-Peng Zhou, *The inter-universal Teichmüller theory and new Diophantine results over rational numbers. II*, arXiv:2510.05448 [math.NT], 2025.

Email address: `123026.urahigashi.toshihisa@gmail.com`