

# Data-dependent PAC-Bayes priors via differential privacy

Speaker: Yi-Shan Wu

Institute of Information Science

Academia Sinica

Taiwan

May 25, 2018

- 1 data-dependent PAC-bound
- 2 Weak convergence suffice

# Notations

- observe  $\mathcal{S} \sim \mathcal{D}^m$  :  $m$  i.i.d. samples from  $\mathcal{D}$
- parameter of NN :  $\mathbf{w} \in \mathbb{R}^p$
- $L_{\mathcal{D}}(\mathbf{w}) = \mathbb{E}_{z \sim \mathcal{D}} [\ell(\mathbf{w}, z)]$
- $L_{\mathcal{D}}(\mathcal{Q}) = \mathbb{E}_{\mathbf{w} \sim \mathcal{Q}} [L_{\mathcal{D}}(\mathbf{w})] = \mathbb{E}_{\mathbf{w} \sim \mathcal{Q}} [\mathbb{E}_{z \sim \mathcal{D}} [\ell(\mathbf{w}, z)]]$
- $\hat{L}_{\mathcal{S}}(\mathcal{Q}) = \frac{1}{m} \sum_{i=1}^m \mathbb{E}_{\mathbf{w} \sim \mathcal{Q}} [\ell(\mathbf{w}, z_i)]$
- generalization error :  $L_{\mathcal{D}}(\mathcal{Q}) - \hat{L}_{\mathcal{S}}(\mathcal{Q})$

# PAC-Bayes bounds

**Theorem 7 (PAC-Bayes; Maurer 2004, Thm. 5)** Under bounded loss  $\ell \in [0, 1]$ , for every  $\delta > 0$ ,  $m \in \mathbb{N}$ , distribution  $\mathcal{D}$  on  $Z$ , and distribution  $P$  on  $\mathbb{R}^p$ ,

$$\mathbb{P}_{S \sim \mathcal{D}^m} \left( (\forall Q) \text{KL}(\hat{L}_S(Q) \| L_{\mathcal{D}}(Q)) \leq \frac{\text{KL}(Q \| P) + \log 2\sqrt{m}}{m} + \frac{1}{m} \log \frac{1}{\delta} \right) \geq 1 - \delta. \quad (2)$$

General PAC-Bayes bound, where the prior  $\mathcal{P}$  is chosen before observing samples.

Typically the **KL term is the dominant quantity** in the bound and analysis is constrained by the need to choose  $Q$  such that  $\text{KL}(Q \| \mathcal{P})$  is not large. If  $\mathbf{w}^*$  receives low probability from  $\mathcal{P}$ , ( $\mathcal{P}$  is chosen badly), the bound is obvious bad.

# Prior results by Lever[2013]

Catoni [2007] started studying the problem on finding bounds for **data-distribution-dependent prior**. Lever [2013] studied the randomized classifier with  $Q(S) \propto \exp(-\tau \hat{L}_S)$ ,  $\mathcal{P}(S) \propto \exp(-\tau L_{\mathcal{D}})$ . They are able to bound  $KL(Q||\mathcal{P})$  independent of  $\mathcal{D}$ , yielding

**Theorem 1 (Lever, Laviolette, and Shawe-Taylor 2013)** For  $S \in Z^m$ , let  $Q(S) = P_{\exp(-\tau \hat{L}_S)}$  be a Gibbs posterior with respect to some measure  $P$  on  $\mathbb{R}^p$  and some bounded (surrogate) risk  $\tilde{L}_S$ . Under bounded loss  $\ell \in [0, 1]$ , for every  $\delta > 0$ ,  $m \in \mathbb{N}$ , distribution  $\mathcal{D}$  on  $Z$ ,

$$\mathbb{P}_{S \sim \mathcal{D}^m} \left( KL(\hat{L}_S(Q(S)) || L_{\mathcal{D}}(Q(S))) \leq \frac{1}{m} \left( \tau \sqrt{\frac{2}{m} \log \frac{2\sqrt{m}}{\delta} + \frac{\tau^2}{2m} + \log \frac{2\sqrt{m}}{\delta}} \right) \right) \geq 1 - \delta. \quad (1)$$

However, the bound can be still loose, since it allows classifier overfits on some distribution.

# data-dependent PAC-Bayes bound (Goal)

Trivial but bad approach to consider several  $\mathcal{P}$ s : by union bound.  
Fix a distribution  $\mathcal{D}$  on  $\mathcal{Z}$ ,

$$R(\epsilon) = \{\mathcal{S} \in \mathcal{Z}^m : (\exists Q) KL(\hat{L}_{\mathcal{S}}(Q) \| L_{\mathcal{D}}(Q)) > \epsilon\} \text{ --- before}$$

$$R(\epsilon_p) = \{\mathcal{S} \in \mathcal{Z}^m : (\exists Q) KL(\hat{L}_{\mathcal{S}}(Q) \| L_{\mathcal{D}}(Q)) > \epsilon_p\} \text{ --- now}$$

If we union  $M$  kinds of  $\mathcal{P}$ , then the probability becomes  $1 - M\delta$ .

**Theorem 8** Under bounded loss  $\ell \in [0, 1]$ , for every  $\delta > 0$ ,  $m \in \mathbb{N}$ , distribution  $\mathcal{D}$  on  $Z$ , and  $\epsilon$ -differentially private data-dependent prior  $\mathcal{P}: Z^m \rightsquigarrow \mathcal{M}_1(\mathbb{R}^p)$ ,

$$\mathbb{P}_{S \sim \mathcal{D}^m} \left( (\forall Q) KL(\hat{L}_S(Q) \| L_{\mathcal{D}}(Q)) \leq \frac{KL(Q \| \mathcal{P}(S)) + \ln 2\sqrt{m}}{m} + \max \left\{ \frac{2}{m} \ln \frac{3}{\delta}, 2\epsilon^2 \right\} \right) \geq 1 - \delta.$$

# differential privacy

Since now our goal is to obtain a "data-dependent prior  $\mathcal{P}$ ", bad  $\mathcal{S}$  may lead to bad prior  $\mathcal{P}$ , and bad prior lead to bad  $R(\epsilon_p)$ . However, If  $\mathcal{P}_{\mathcal{S}}$  is generated by some "stable" mechanism, then maybe we don't have to union too much of them.

## differential private algorithm

**Definition 3** A randomized algorithm  $\mathcal{A}: Z^m \rightsquigarrow T$  is  $(\epsilon, \delta)$ -differentially private if, for all pairs  $S, S' \in Z^m$  that differ at only one coordinate, and all measurable subsets  $B \subseteq T$ , we have  $\mathbb{P}\{\mathcal{A}(S) \in B\} \leq e^\epsilon \mathbb{P}\{\mathcal{A}(S') \in B\} + \delta$ .

Under such intuition, we have the following theorems,

# differential privacy

**Theorem 6 (Dwork et al. 2015b, Thm. 11)** Let  $m \in \mathbb{N}$ , let  $\mathcal{A}: Z^m \rightsquigarrow T$ , let  $\mathcal{D}$  be a distribution over  $Z$ , let  $\beta \in (0, 1)$ , and, for each  $t \in T$ , fix a set  $R(t) \subseteq Z^m$  such that  $\mathbb{P}_{S \sim \mathcal{D}^m}\{S \in R(t)\} \leq \beta$ . If  $\mathcal{A}$  is  $\epsilon$ -differentially private for  $\epsilon \leq \sqrt{\ln(1/\beta)/(2m)}$ , then  $\mathbb{P}_{S \sim \mathcal{D}^m}\{S \in R(\mathcal{A}(S))\} \leq 3\sqrt{\beta}$ .

(for each  $\mathcal{P} \in \mathcal{M}_1(\mathbb{R}^p)$ , fix a set  $R(\mathcal{P}) = \{S \in Z^m : \text{conditions}\}$  such that  $\mathbb{P}_{S \sim \mathcal{D}^m}\{S \in R(\mathcal{P})\} \leq \beta$ )

That is, with a stable algorithm, the probability only enlarge to  $3\sqrt{\beta}$ .

**Theorem 8** Under bounded loss  $\ell \in [0, 1]$ , for every  $\delta > 0$ ,  $m \in \mathbb{N}$ , distribution  $\mathcal{D}$  on  $Z$ , and  $\epsilon$ -differentially private data-dependent prior  $\mathcal{P}: Z^m \rightsquigarrow \mathcal{M}_1(\mathbb{R}^p)$ ,

$$\mathbb{P}_{S \sim \mathcal{D}^m} \left( (\forall Q) \text{KL}(\hat{L}_S(Q) \| L_{\mathcal{D}}(Q)) \leq \frac{\text{KL}(Q \| \mathcal{P}(S)) + \ln 2\sqrt{m}}{m} + \max \left\{ \frac{2}{m} \ln \frac{3}{\delta}, 2\epsilon^2 \right\} \right) \geq 1 - \delta.$$

If we have probability  $\geq 1 - \beta$  in Theorem 7, then  $\delta = 3\sqrt{\beta}$



# Ideally

For convenience, choose  $\mathcal{P}_{\mathbf{w}} = \mathcal{N}(\mathbf{w}, \Sigma)$ , where  $\mathbf{w}_0 \in \mathbb{R}^p$  is chosen by private mechanism and approximately minimizing  $\hat{L}_S$ .  
(Ideal prior : easy to calculate & data dependent)

**Theorem 10** *Let  $\tau > 0$  and assume loss is bounded  $\ell \in [0, 1]$ . One sample from the Gibbs posterior  $P_{\exp(-\tau \hat{L}_S)}$  is  $\frac{2\tau}{m}$ -differentially private.*

However,

- Gibbs distribution is intractable.
- We can only approximate it by producing a sequence of samples that converges in distribution to the target Gibbs distribution.  
Weak convergence, however, does not yield privacy guarantees.

- 1 data-dependent PAC-bound
- 2 Weak convergence suffice

# Weak convergence suffice

**Theorem 16 (Weak convergence suffices)** *Let  $\tau > 0$  and let  $\Sigma \in \mathbb{R}^{p \times p}$  be positive definite. For every  $\mathbf{w} \in \mathbb{R}^p$  and  $S \in Z^m$ , let  $P_{\mathbf{w}} = \mathcal{N}(\mathbf{w}, \Sigma)$ ,  $Q_{\mathbf{w}}^S = (P_{\mathbf{w}})_{\exp(-\tau \tilde{L}_S)}$ , where  $\tilde{L}_S(\cdot)$  is bounded (surrogate) risk. Then, for every  $\epsilon' > 0$  and  $\delta, \delta' \in (0, 1)$ , with probability at least  $1 - \delta - \delta'$  over  $S \sim \mathcal{D}^m$  and a sequence  $\mathbf{w}_1, \mathbf{w}_2, \dots$  that, conditional on  $S$ , converges in distribution a.s. to an  $\epsilon$ -differentially private mean  $\mathbf{w}^*(S)$ , there exists  $N \in \mathbb{N}$ , such that, for all  $n > N$ ,*

$$\text{KL}(\hat{L}_S(Q_{\mathbf{w}_n}^S) || L_{\mathcal{D}}(Q_{\mathbf{w}_n}^S)) \leq \frac{\text{KL}(Q_{\mathbf{w}_n}^S || P_{\mathbf{w}_n}) + \ln 2\sqrt{m}}{m} + \max\left\{\frac{2}{m} \ln \frac{3}{\delta}, \epsilon^2\right\} + \epsilon'.$$

Here, the sequence of  $\mathbf{w}_i$  are obtained by SGLD that SGLD converges weakly to Gibbs distribution.

# Some prior works

## Lemma 15

Using bounded cross-entropy. Define  $\mathcal{P}_{\mathbf{w}} = \mathcal{N}(\mathbf{w}, \Sigma)$  and let  $G, G_1, G_2, \dots \in \mathcal{M}_1(\mathbb{R}^p)$ , where  $G_n \rightarrow G$  weakly. Then  $\forall \epsilon, \delta, \exists N \in \mathbb{N}$  such that  $\forall n > N$ , w.h.p. over  $(\mathbf{w}_n, \mathbf{w}_n^*) \sim (G_n, G)$  that

$$KL(\hat{L}_{\mathcal{S}}(\mathcal{Q}_{\mathbf{w}_n}) \| L_{\mathcal{D}}(\mathcal{Q}_{\mathbf{w}_n})) \leq KL(\hat{L}_{\mathcal{S}}(\mathcal{Q}_{\mathbf{w}_n^*}) \| L_{\mathcal{D}}(\mathcal{Q}_{\mathbf{w}_n^*})) + \epsilon$$

$$KL(\mathcal{Q}_{\mathbf{w}_n^*} \| \mathcal{P}_{\mathbf{w}_n^*}) \leq KL(\mathcal{Q}_{\mathbf{w}_n} \| \mathcal{P}_{\mathbf{w}_n}) + \epsilon$$

- ❶ Let  $(\mathbf{w}_n, \mathbf{w}_n^*) \sim (G_n, G)$ . There exists a sequence of random variables  $\mathbf{w}^*, \mathbf{w}_1, \mathbf{w}_2, \dots$  such that  $\mathbf{w}_n \sim G_n$  for all  $n \in \mathbb{N}$ ,  $\mathbf{w}^* \sim G$  and  $\mathbf{w}_n \sim \mathbf{w}^*$  a.s.
- ❷ If  $\mathcal{P}_{\mathbf{w}} = \mathcal{N}(\mathbf{w}, \Sigma)$ , then  $\mathcal{P}_{\mathbf{w}_n} \rightarrow \mathcal{P}$  and  $KL()$  have these inequalities