


DiTox-3 *Plus*


BINDS & ELIMINATE TOXIN





Is an activated, most effective mycotoxin adsorbent, hydrated sodium calcium aluminosilicate (HSCAS) is specially formulated to adsorb & eliminate all major mycotoxins affecting poultry health & productivity.

EFFECT OF MYCOTOXINS ON POULTRY




PURPOSE : To bind and eliminate mycotoxins & their secondary metabolites (chemicals).

COMPOSITION

Natural dipolar phyllosilicates, Activated carbon, Mannan oligosaccharides (MOS), Neem powder, Liver & Kidney rejuvenators.

DiTox-3^{Plus}

BINDS & ELIMINATE TOXIN



ACTIVATED CHARCOAL

Used in DiTox-3 plus having properties of high adsorption, low contents of ash, iron, acid soluble, water soluble, and free from heavy metals making it suitable for high rate of mycotoxin adsorption. Activated Charcoal was found to reduce ochratoxin and is efficacious in preventing T2 toxicosis.

NATURAL DIPOLAR PHYLLOSILICATES

Natural dipolar phyllosilicates, are effective toxin binders used in poultry feed to mitigate the harmful effects of mycotoxins like aflatoxins and zearalenone. Their layered structure and dipolar nature enable strong adsorption of polar toxins, preventing their absorption in the gut.

MANNAN OLIGOSACCHARIDES (MOS)

Mannan oligosaccharides (MOS) are natural toxin binders derived from yeast cell walls, known for their dual role in poultry feed. They effectively bind mycotoxins, particularly zearalenone and ochratoxins, reducing their absorption in the gut. MOS also support gut health by inhibiting the binding of harmful pathogens, such as E. coli and Salmonella, to intestinal walls.

NEEM POWDER

Antibacterial, antiviral, antifungal, antiprotozoal, hepatoprotective, immunomodulator.

LIVER AND KIDNEY REJUVENATORS

Helps to keep healthy liver & kidney during detoxifying the toxins

Guideline limits for mycotoxins in Poultry feeds to reduce negative effects on health and performance:

	Maximum tolerance limit*
Aflatoxin	20 ppb
Ochratoxin A	40 ppb
Citrinin toxin	100 ppb
Zearalenone	400 ppb
DON (Vomitoxin)	5 ppm
T2 toxin	200 ppb

Relative toxicity of different mycotoxins on different livestock species


(★ mild toxicity ★★ moderate toxicity ★★★ high toxicity)

Toxin	Poultry	Swine	Ruminant
Aflatoxins	★★★	★★	★
Ochratoxins	★★★	★	★
T2 toxin	★★	★★★	★★★
Deoxynivalenol	★	★★	★★
Zearalenone	★	★★★	★★
Fumonisin	★	★★★	★

- DiTox-3 plus have a high affinity for aflatoxin B1 forming a complex which was stable at temperatures of 25 and 37°C. in a pH range of 2-10.
- DiTox-3 plus is used to eliminate Chemical Toxins.

DiTox-3^{Plus}

BINDS & ELIMINATE TOXIN



BENEFITS

1. Binds and neutralizes toxins, ensuring safer and higher-quality feed for poultry.
2. Mannan oligosaccharides (MOS) attracts and bind pathogens and their toxins in the gastrointestinal tract.
3. Phyllosilicates adsorb toxins effectively due to their high surface area and layered structure, allowing them to trap and neutralize harmful substances within their interlayer spaces.
4. Improves Liver and Kidney functions.

MAIN EFFECT OF DITOX-3 plus

- Effective degradation, adsorption of mold & its metabolites.
- Optimize the intestinal metabolism, improve digestion.
- Stimulate the body's immune response, enhance immunity, reduce the incidence of a disease.
- Improve bird's production & reproductive performance
- Improving economic efficiency

DOSAGE

Regular : 1 kg/tonne of feed

High moisture and toxin suspected feed : 2 kg/tonne of feed

or As advised by the Veterinarian.

PRESENTATION

25 KG bag

SHELF LIFE

24 months from the date of manufacture.



(An ISO 9001 : 2015 Certified Company)



GMP & HACCP Certified

Manufactured and Marketed By :

ZENMAK NUTRIGENCIES AND HEALTH PVT. LTD.

Regd. Off: No. F2, Sand Stone Apartments,
14/5, Main Road, Velacherry, Chennai - 600 042.

E-mail : info@zenmakglobal.com

Contact : 93459 13689, 93449 31464

Web : www.zenmakglobal.com

