The relationship between a set of variables and miles per gallon

Tushar Mahajan

21-05-2020

Executive Summary

We are looking at a data set of a collection of cars, and are interested in exploring the relationship between a set of variables and miles per gallon. In particularly the following two questions:

- "Is an automatic or manual transmission better for MPG"
- "Quantify the MPG difference between automatic and manual transmissions"

We'll take the following steps:

- Process the data, for use of this project
- Explore the data, especially focussing on the two paramaters we are interested in (Transmission and MPG)
- Model selection, where we try different models to help us answer our questions
- Model examination, to see wether our best model holds up to our standards
- A Conclusion where we answer the questions based on the data

Processing

First change 'am' to factor (0 = automatic, 1 = manual) And make cylinders a factor as well (since it is not continious)

```
library(ggplot2)
library(dglyr)
library(ggfortify)
data(mtcars)
mtcarsFactors <- mtcars
mtcarsFactors$am <- as.factor(mtcarsFactors$am)
levels(mtcarsFactors$am) <- c("automatic", "manual")
mtcarsFactors$cyl <- as.factor(mtcarsFactors$cyl)
mtcarsFactors$gear <- as.factor(mtcarsFactors$gear)
mtcarsFactors$vs <- as.factor(mtcarsFactors$vs)
levels(mtcarsFactors$vs) <- c("V", "S")</pre>
```

Exploratory data analyses

Look at the dimensions & head of the dataset to get an idea

```
# Res 1
dim(mtcarsFactors)
```


```
## [1] 32 11
```

```
# Res 2
head(mtcarsFactors)
```

```
##
                      mpg cyl disp hp drat
                                                                    am gear carb
                                                    qsec vs
## Mazda RX4
                                160 110 3.90 2.620 16.46
                     21.0
                                                                manual
                               160 110 3.90 2.875 17.02
## Mazda RX4 Wag
                     21.0
                                                                manual
                                                                                4
## Datsun 710
                     22.8
                                     93 3.85 2.320 18.61
                                                                manual
                                                                                1
## Hornet 4 Drive
                     21.4
                             6
                                258 110 3.08 3.215 19.44
                                                           S automatic
                                                                           3
                                                                                1
## Hornet Sportabout 18.7
                                360 175 3.15 3.440 17.02
                                                                           3
                                                                                2
                             8
                                                           V automatic
## Valiant
                     18.1
                                225 105 2.76 3.460 20.22
                                                           S automatic
                                                                                1
```

Now just take a look at the realtionship between the two parameters we are intereseted in.

```
# Figure 1
library(ggplot2)
p <- ggplot(mtcarsFactors, aes(am, mpg))
p + geom_boxplot(aes(fill = am))</pre>
```




Even this shows clearly that the manual transmissions have higher mpg's there could be a bias in the dataset that we are overlooking. Before creating a model we should look at which parameters to include besides 'am'. So we look at all correlations of parameters and take only those higher then the 'am' correlation.

```
# Res 3
cors <- cor(mtcars$mpg, mtcars)
orderedCors <- cors[,order(-abs(cors[1,]))]
orderedCors</pre>
```

```
##
                        cyl
                               disp
                                                drat
                wt
                                         hp
                                                          ٧S
                                                                  am
                                                                         carb
       mpg
##
  ##
      qsec
  0.4186840
##
# Res 4
amPos <- which(names(orderedCors)=="am")</pre>
subsetColumns <- names(orderedCors)[1:amPos]</pre>
subsetColumns
## [1] "mpg"
          "wt"
               "cyl" "disp" "hp"
                               "drat" "vs"
```

```
# Figure 2
mtcarsFactors[,subsetColumns] %>%
    ggpairs(
    mapping = ggplot2::aes(color = am),
    upper = list(continuous = wrap("cor", size = 3)),
    lower = list(continuous = wrap("smooth", alpha=0.4, size=1), combo = wrap("dot"))
)
```



Model selection

Now that we have seen that mpg has many other (stronger) correlations than just 'am' we can guess that a model predicting the mpg solely on this parameter will not be the most accurate model. Let's check this out.

First we start with the basic model

```
# Res 5
basicFit <- lm(mpg ~ am, mtcarsFactors)
summary(basicFit)</pre>
```

```
##
## Call:
## lm(formula = mpg ~ am, data = mtcarsFactors)
##
## Residuals:
##
      Min
               1Q Median
                               3Q
                                      Max
## -9.3923 -3.0923 -0.2974 3.2439 9.5077
##
## Coefficients:
##
              Estimate Std. Error t value Pr(>|t|)
## (Intercept)
                17.147
                            1.125 15.247 1.13e-15 ***
                 7.245
                            1.764
                                    4.106 0.000285 ***
## ammanual
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 4.902 on 30 degrees of freedom
## Multiple R-squared: 0.3598, Adjusted R-squared: 0.3385
## F-statistic: 16.86 on 1 and 30 DF, p-value: 0.000285
```

The p-values are actually quite low, the R-squared is problematic however. Now go to the other side of the spectrum by fitting all parameters of mtcars

```
# Res 6
totalFit <- lm(mpg ~ ., mtcarsFactors)
summary(totalFit)

##
## Call:
## lm(formula = mpg ~ ., data = mtcarsFactors)
##
## Residuals:
## Min    1Q Median    3Q Max
## -3.2015 -1.2319    0.1033    1.1953    4.3085
##
## Caefficients:</pre>
```

```
## Coefficients:
               Estimate Std. Error t value Pr(>|t|)
## (Intercept) 15.09262
                          17.13627
                                     0.881
                                             0.3895
               -1.19940
                                    -0.502
## cyl6
                           2.38736
                                             0.6212
## cyl8
                3.05492
                           4.82987
                                     0.633
                                             0.5346
                                     0.708
## disp
               0.01257
                           0.01774
                                             0.4873
               -0.05712
                           0.03175
                                    -1.799
## hp
                                             0.0879
## drat
                0.73577
                           1.98461
                                     0.371
                                             0.7149
                                    -1.857
## wt
               -3.54512
                           1.90895
                                             0.0789 .
               0.76801
                           0.75222
                                     1.021
                                             0.3201
## qsec
## vsS
                2.48849
                           2.54015
                                     0.980
                                             0.3396
## ammanual
                3.34736
                           2.28948
                                     1.462
                                             0.1601
## gear4
               -0.99922
                           2.94658
                                    -0.339
                                             0.7382
## gear5
                1.06455
                           3.02730
                                     0.352
                                             0.7290
## carb
                0.78703
                           1.03599
                                     0.760
                                             0.4568
## ---
## Signif. codes: 0 '*** 0.001 '** 0.01 '* 0.05 '.' 0.1 ' 1
##
```

Residual standard error: 2.616 on 19 degrees of freedom

```
## Multiple R-squared: 0.8845, Adjusted R-squared: 0.8116
## F-statistic: 12.13 on 12 and 19 DF, p-value: 1.764e-06
```

The R-squared has improved, but the p-values hardly show any significance anymore. Perhaps this is due to overfitting. We now have to meet somewhere in the middle. Let's iterare using the step method

```
# Res 7
bestFit <- step(totalFit,direction="both",trace=FALSE)
summary(bestFit)</pre>
```


```
##
## Call:
## lm(formula = mpg ~ wt + qsec + am, data = mtcarsFactors)
##
## Residuals:
##
      Min
                1Q Median
                                3Q
                                       Max
  -3.4811 -1.5555 -0.7257
                           1.4110
                                   4.6610
##
## Coefficients:
##
              Estimate Std. Error t value Pr(>|t|)
                9.6178
                            6.9596
                                     1.382 0.177915
## (Intercept)
                            0.7112 -5.507 6.95e-06 ***
                -3.9165
## wt
                1.2259
                            0.2887
                                     4.247 0.000216 ***
## qsec
                2.9358
                                     2.081 0.046716 *
## ammanual
                            1.4109
## Signif. codes: 0 '*** 0.001 '** 0.01 '* 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 2.459 on 28 degrees of freedom
## Multiple R-squared: 0.8497, Adjusted R-squared: 0.8336
## F-statistic: 52.75 on 3 and 28 DF, p-value: 1.21e-11
```

Model examination

The resulting best model mpg ~ wt + qsec + am is actually dependent on the transmission (am), but also weight (wt) and 1/4 mile time (qsec). All have significant p-values The R-squared is pretty good to (0.85)

Now let's look (amongst others) at the Residuals vs Fitted

```
# Figure 3
autoplot(bestFit)
```



The 'Normal Q-Q' plot looks ok, but the 'Residuals vs Fitted' and 'Scale-Location' both show worrysome trends

Conclusion

The question "Is an automatic or manual transmission better for MPG" can be answered because all models (#Res5, #Res6 and #Res7) show that, holding all other paramters constant, manual transmission will increase your MPG.

The question "Quantify the MPG difference between automatic and manual transmissions" is harder to answer.

Based on the 'bestFit' (#Res7) model mpg ~ wt + qsec + am we could conclude that (with a p < 0.05 confidence) cars with manual transmission have 2.9358 (say 3) more miles per gallon than automatic transmissions. The model seems clean with a p < 0.05 and R squared of 0.85

The residuals vs fitted chart however warns us that there is something missing in our model. The real problem I think is that we only have 32 observations to train on (#Res1) and that observations hardly have overlap on the parameters 'wt' and 'qsec' (amongst others) if we look at the diagonal in the matrix chart (#Figure2)

Although the conclusion of ca. 3 mpg better performance on manual transmissions seems feasible, I cannot with confidence conclude that this model will fit all future observations.